

Search for Neutral Resonances Decaying into a Z Boson and a Pair of b Jets or Tau Leptons postprint

Authors: Khachatryan, Vardan and others

Date: 2016-08-22T00:00:00+00:00

Abstract

A search is performed for a new resonance decaying into a lighter resonance and a Z boson. Two channels are studied, targeting the decay of the lighter resonance into either a pair of oppositely charged tau leptons or a b-bbar pair. The Z boson is identified via its decays to electrons or muons. The search exploits data collected by the CMS experiment at a centre-of-mass energy of 8 TeV, corresponding to an integrated luminosity of 19.8 inverse femtobarns. No significant deviations are observed from the standard model expectation and limits are set on production cross sections and parameters of two-Higgs-doublet models.

Full Text

Preamble

Search for neutral resonances decaying into a Z boson and a pair of b jets or τ leptons The CMS Collaboration*

Abstract

A search is performed for a new resonance decaying into a lighter resonance and a Z boson. Two channels are studied, targeting the decay of the lighter resonance into either a pair of oppositely charged τ leptons or a bb pair. The Z boson is identified via its decays to electrons or muons. The search exploits data collected by the CMS experiment at a centre-of-mass energy of 8 TeV, corresponding to an integrated luminosity of 19.8 fb⁻¹. No significant deviations are observed from the standard model expectation and limits are set on production cross sections and parameters of two-Higgs-doublet models.

Submitted to Physics Letters B c(13) 2016 CERN for the benefit of the CMS Collaboration. CC-BY-3.0 license *See Appendix A for the list of collaboration members

1 Introduction

The observation of a new particle with a mass of approximately 125 GeV was reported by the ATLAS and CMS experiments at the CERN LHC in the WW , ZZ and $\gamma\gamma$ final states [1–3]. Evidence of the decay of the particle to pairs of fermions ($\tau\tau$ and $b\bar{b}$) has also been reported in Refs. [4, 5]. The measurements of branching fractions, production rates, spin and parity are all consistent with the predictions for the standard model (SM) Higgs boson [6], wherein a single doublet of Higgs fields is present. However, additional Higgs bosons are expected in simple extensions of the SM scalar sector, such as models with two Higgs-boson doublets (2HDMs) [7]. These models predict five physical Higgs particles that arise as a consequence of the electroweak symmetry-breaking mechanism: two neutral CP-even scalars (h , H), one neutral CP-odd pseudoscalar (A), and two charged scalars (H_{\pm}).

An important motivation for 2HDMs is that such models provide a way to accommodate the asymmetry between matter and antimatter observed in the universe [7, 8]. An extension of the SM scalar sector with two Higgs boson doublets would also naturally arise in supersymmetry [9, 10], which requires a scalar structure more complex than a single doublet. Axion models [11] provide a strong interaction that does not violate CP symmetry and give rise to an effective low-energy theory with two Higgs doublets. Finally, it has recently been noted [12] that certain realisations of 2HDMs can accommodate the muon $g-2$ anomaly [13] without violating present theoretical and experimental constraints.

In the most general case, 14 parameters describe the scalar sector of a 2HDM [7]. Only six free parameters remain once the experimental observations are included by imposing the so-called Z_2 symmetry to suppress flavour changing neutral currents, and by fixing both the values of the mass of the recently discovered SM-like Higgs boson (125 GeV) [14] and the electroweak vacuum expectation value (246 GeV). The compatibility of a SM-like Higgs boson with 2HDMs is possible in the so-called alignment limit. The alignment limit is reached when $\cos(\beta - \alpha) \rightarrow 0$, where $\tan \beta$ is the ratio of the vacuum expectation values and α is the mixing angle of the two Higgs doublets. In such a regime, one of the CP-even scalars, h or H , is identified with the SM-like Higgs boson. A recent theoretical study [8] has shown that, in this limit, a large mass splitting (>100 GeV) between the A and H bosons would favour the electroweak phase transition that would be at the origin of baryogenesis in the early universe, satisfying thereby the currently observed matter-antimatter asymmetry. In this context, the most frequent decay mode of the pseudoscalar A boson would be $A \rightarrow ZH$. Since the analysis strategy presented in this paper is independent of the assumed model and parity of the resonance, the results can also be interpreted in the reversed topology $H \rightarrow ZA$, where the expected 2HDM mass hierarchy is

inverted and the mass of A is expected to be light [15]. For both topologies, the lighter scalar resonance (A or H) is not identified with the SM-like Higgs boson.

This paper describes the first CMS search for a new resonance decaying into a lighter resonance and a Z boson. Two searches are performed, targeting the decay of the lighter resonance into either a pair of oppositely charged τ leptons or a bb pair. In both cases, the Z boson is identified via its decay into a pair of oppositely charged electrons or muons (light leptons), labelled in the text by the symbol l . The choice of bb and $\tau\tau$ final states is motivated by the large branching fractions predicted in most of the 2HDM phase space [16]. For the $\tau\tau$ channel, the following $\tau\tau$ final states are considered: $e\mu$, $e\tau h$, $\mu\tau h$, and τh τh , where τh indicates the decays $\tau \rightarrow \text{hadrons} + \tau$. Given its sensitivity to a different region of the 2HDM parameter space, the search presented in this paper is complementary to other related searches performed in the same final state by the ATLAS and CMS collaborations [17, 18].

2 The CMS detector

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Located in concentric layers within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter (HCAL), each composed of one barrel and two endcap sections. These layers provide coverage up to a pseudorapidity $|\eta| = 2.5$. Extensive forward calorimetry complements are provided by the endcap detectors for $|\eta| < 5.2$. Combining the energy measurement in the ECAL with the measurement in the tracker, the momentum resolution for electrons with $p_T \sim 45$ GeV from $Z \rightarrow ee$ decays ranges from 1.7% for nonshowering electrons in the barrel region to 4.5% for showering electrons in the endcaps [19]. Muons are measured in gas-ionisation detectors embedded in the steel flux-return yoke outside the solenoid. They cover the pseudorapidity range $|\eta| < 2.4$, with detection planes made using three technologies: drift tubes, cathode strip chambers, and resistive plate chambers. Matching muons to tracks measured in the silicon tracker results in a relative transverse momentum resolution for muons with $20 < p_T < 100$ GeV of 1.3–2.0% in the barrel and better than 6% in the endcaps [20]. The first level of the CMS trigger system uses information from the calorimeters and muon detectors to select the most interesting events. A high-level trigger processor farm decreases the event rate from approximately 100 kHz to 600 Hz before data storage. A more detailed description of the CMS detector, together with a definition of the coordinate system and kinematic variables, can be found in Ref. [21].

3 Data and simulated samples

The data used for this search were collected by the CMS experiment at $\sqrt{s} = 8$ TeV, and correspond to a total integrated luminosity of 19.8 fb $^{-1}$. The

average number of interactions per bunch crossing (pileup) in the data was 21 [22]. Events were selected using dielectron and dimuon triggers [19, 20]. These triggers have p_T thresholds of 17 and 8 GeV for the leading and subleading lepton respectively, and require relatively loose reconstruction and identification criteria.

The main SM background processes giving rise to prompt leptons are W/Z +jets, $t\bar{t}$ +jets, tW , and diboson production (WW , ZZ , and WZ). The SM background contribution from ZZ is generated at next-to-leading order (NLO) with POWHEG 1.0 [23] for the $\tau\tau$ channel and using the leading-order (LO) MADGRAPH 5.1 Monte Carlo (MC) program [24], matched to PYTHIA 6.4 [25] for the parton showering and hadronization, for the bb channel. Single top quark events are generated at NLO using POWHEG 1.0. Simulated events for other samples are obtained using the MADGRAPH 5.1 MC matched to PYTHIA 6.4. The PYTHIA parameters affecting the description of the underlying event are set to those of the $Z2^*$ tune [26]. All generators used for processes including τ leptons in the final state are interfaced with TAUOLA 2.4 [27] for the simulation of the τ decays. The detector response is simulated using a detailed description of CMS, based on the GEANT4 toolkit [28]. The simulated samples account for contributions from pileup collisions that reflect the distributions observed in data. The trigger and reconstruction efficiency in the simulation is rescaled by as much as 2% in order to match that measured in the data [22].

Two benchmark 2HDM processes are considered as signal: $H \rightarrow ZA$ and $A \rightarrow ZH$, where the lightest boson (pseudoscalar or scalar, according to the process) can decay to $\tau\tau$ or bb , and the Z decays to $\tau\tau$. The MADGRAPH 5.1 generator, interfaced to PYTHIA 6.4 and TAUOLA 2.4, was used to generate signal samples corresponding to different values of A and H masses (m_A and m_H , respectively). The same properties of the SM Higgs boson are assigned to the lightest scalar boson, h , and its mass m_h is fixed at 125 GeV. The identification of the observed Higgs boson, together with all its measured properties, with the scalar h constrains the phenomenologically reliable parameter space regions to not depart from the SM-like condition $\cos(\beta - \alpha) \approx 0$. This corresponds to the so-called alignment limit [29]. Considering the parameter space still allowed by direct searches [10], the chosen values for $\cos(\beta - \alpha)$ and $\tan\beta$ are 0.01 and 1.5, respectively, and type-II Yukawa couplings are assumed for the benchmark processes.

The masses of the charged Higgs bosons (m_{H^\pm}) are kept equal to the highest mass involved in the signal process (m_H or m_A) to preserve the degeneracy $m_{2H^\pm} = m_{2H/A}$ [15], denoting with $m_{H/A}$ the mass of the scalar H or the mass of the pseudoscalar A . The value of the m_{12} parameter, the soft $Z2$ symmetry breaking mass, was set to $m_{2H^\pm} \tan\beta / (1 + \tan^2\beta)$, according to the minimal supersymmetric standard model (MSSM) parametrisation [9]. The value of the complex couplings μ_6 and μ_7 in this parametrisation are set to zero, in order to avoid tree-level CP violation. The production cross sections, used

for the normalisation of the signal samples, are computed using the SUSHI 1.4 program [30], which provides next-to-next-to-leading-order (NNLO) predictions. The branching fraction for the heavy and light Higgs bosons are obtained using the 2HDMC 1.6 program [31], following the guidelines in Refs. [32, 33].

The signal benchmark where the light boson decays into $\tau\tau$ is simulated for values of mH/A and mA/H varying in the ranges 200–1000 and 15–900 GeV, respectively, with the constraint $mH/A > mA/H + mZ$. For the bb analysis the lower bound for the invariant mass mA/H goes down to 10 GeV. The region where mH is smaller than m_h is not pertinent in this model.

4 Event reconstruction and selection

Event reconstruction is based on the particle-flow algorithm [34, 35], which exploits information from all the CMS subdetectors to identify and reconstruct individual particles in the event: muons, electrons, photons, charged and neutral hadrons. Such particles are algorithmically combined to form the jets, the τh candidates, the missing transverse momentum p_{miss}^T , defined as the projection on the plane perpendicular to the beams of the negative vector sum of the particles momenta and its magnitude, denoted as E_{miss}^T . To minimise the contributions from pileup interactions, charged tracks are required to originate from the primary vertex (reconstructed using the deterministic annealing algorithm [36]), which is the one characterised by the largest p^T sum of its associated tracks.

Electrons are identified by combining information from tracks and ECAL clusters, including energy depositions from final-state radiation [19]. Muons are identified through a combined fit to position measurements from both the inner tracker and the muon detectors [20]. The τh objects are identified and reconstructed using the “hadron-plus-strips” algorithm [37], which uses charged hadrons and photons to reconstruct the main hadronic decay modes of the τ lepton: one charged hadron, one charged hadron and photons, and three charged hadrons. Electrons and muons can be misidentified as hadronic taus if produced in jets or if close-by activity from pile-up or bremsstrahlung is present. These misidentifications are suppressed using dedicated criteria based on the consistency between the measurements in the tracker, the calorimeters, and the muon detector [37]. To reject nonprompt or misidentified leptons, requirements are imposed on the isolation criteria, based on the sum of deposited energies. The absolute lepton isolation I_{abs} is defined by the scalar sum of the p^T of the charged particles from the primary vertex, neutral hadrons, and photons in an isolation cone of size $\Delta R = \sqrt{(\Delta \eta)^2 + (\Delta \phi)^2} = 0.4$ ($\Delta R = 0.3$ for electrons), centred around the lepton direction. To reduce the effect from pileup, the energy deposit released in the isolation cone by charged particles not associated with the primary vertex is subtracted from the neutral particles p^T scalar sum. For electrons and muons the relative isolation, defined as $I_{\text{rel}} = I_{\text{abs}}/p^T$, is used.

Jets are clustered using the anti- k^T algorithm [38], with a distance parame-

ter of 0.5, as implemented in the FASTJET software package [39]. Charged particles not associated with the primary vertex are excluded by means of the charged-hadron subtraction technique [40]. The remaining energy originating from pileup interactions, including the neutral components, is subtracted based on the median energy density in the detector computed through the effective jet area technique [41]. The identification of b quark initiated jets is achieved through the combined secondary vertex (CSV) algorithm [42], which exploits observables related to the long lifetime of B hadrons.

4.1 Selection for $Z \rightarrow$

In selecting $b\bar{b}$ and $\tau\tau$ events, the leptons from Z boson decay are required to be well within the CMS trigger and detector acceptance of $p_T > 20$ GeV and $|\eta| < 2.5$ for electrons, and $p_T > 20$ GeV, $|\eta| < 2.4$ for muons. Muon momentum-scale [20] and electron energy corrections [19] are applied to recover the global shift of the scale observed between data and simulation. The requirement on the relative isolation for the leptons is set to $I_{rel} < 0.15$ for electrons and $I_{rel} < 0.2$ for muons in selecting $b\bar{b}$ events. For the leptons from the Z boson, in the case of $\tau\tau$ events, the required relative isolation is $I_{rel} < 0.3$. The presence of two reconstructed same-flavour, oppositely charged lepton candidates forming a pair with invariant mass in the range of 76–106 GeV is required to suppress contamination of non-resonant Drell–Yan+jets and $t\bar{t}$ processes. In events where multiple Z candidates are present, the lepton pair with the invariant mass closest to the nominal Z boson mass [43] is chosen.

4.2 Event selection for $b\bar{b}$

For the $b\bar{b}$ search, the jets are selected to be in the kinematic region $p_T > 30$ GeV and $|\eta| < 2.4$. At least two CSV b-tagged jets are required to be present in the event, to reduce the contribution of Z+light-parton jets (originating from gluons or u, d, or s quarks) events. The threshold on the b tagging discriminator corresponds to a b tagging efficiency greater than 65% and to a misidentification probability for light-parton jets of 1% [42]. The two b-tagged jets with highest values of the CSV discriminant are considered as candidate decay products of the new light resonance.

The EmissT significance [44, 45] is used to suppress background events originating from $t\bar{t}$ processes. This variable provides an event-by-event assessment of the likelihood that the observed missing transverse energy is consistent with zero given the reconstructed content of the event and known measurement resolutions. This variable is a stronger discriminant against $t\bar{t}$ background than EmissT alone and also provides smaller systematic uncertainties. The distribution of the $t\bar{t}$ component motivates the requirement on the EmissT significance to be smaller than 10.

4.3 Event selection for $\tau\tau$

To increase the signal sensitivity in the high $\tau\tau$ mass region, the $\tau\tau$ event selection includes the requirement of a transversely boosted Z boson ($p_T > 20$ GeV), together with a large (>1.5 rad) azimuthal angle between the Z boson flight direction and $p_{\text{miss}T}$, particularly effective in suppressing the Z+jets background. In addition to the two light leptons required to reconstruct the Z boson, two additional oppositely charged and different-flavor leptons (e, μ , and τh) are used to reconstruct the A or H boson candidate. The requirements on the pseudorapidity for light leptons are the same as for the Z decay leptons, with the p_T threshold lowered to 10 GeV.

The τh candidates are required to have $p_T > 20$ GeV and $|\eta| < 2.3$. The relative isolation for electrons and muons, and the absolute isolation for τ leptons are required to be smaller than 0.3 and 2 GeV, respectively. Since the Z+jets background is characterised by a softer lepton transverse momentum spectrum than the signal one, this background is reduced by selecting events with high LT, where LT indicates the scalar sum of the visible p_T of the decay products from a $\tau\tau$ pair. Both the isolation requirements and the value of the LT threshold are determined as a result of an optimisation procedure that maximises the expected significance of the searched signal. The optimal requirement on the LT quantity is found by scanning the threshold between 20 and 200 GeV, at intervals of 20 GeV.

Jets are required to have $p_T > 30$ GeV and $|\eta| < 4.7$. To reduce the large $t\bar{t}$ background, all events with at least one jet with $p_T > 20$ GeV and $|\eta| < 2.4$, reconstructed as a jet originating from a b quark according to the output of the CSV discriminator used for tagging, are vetoed.

To calculate the $\tau\tau$ invariant mass, the secondary-vertex fit algorithm (SVFIT) [46] is used, a likelihood-based method that combines the reconstructed $p_{\text{miss}T}$ and its resolution with the momentum of the visible τ decay products to obtain an estimator of the mass of the parent particle.

5 Modelling of the background

5.1 The bb channel

The relevant sources of background for the bb final state originate from Z+jets processes, $t\bar{t}$ and tW production, diboson production, and vector boson production in association with a SM Higgs boson. The contributions of Z+jets and $t\bar{t}$ backgrounds are measured by means of a data-based method, the diboson and tW backgrounds are normalised to the CMS measurements. For these backgrounds, the shapes are taken from MC, while the normalisations are extracted from data. The vector boson production in association with a SM Higgs boson is normalised to the theoretical prediction.

The comparison of data and predictions after the selection of events for the bb final state shows the importance of an accurate theoretical calculation of the

Z+jets production rate. In particular, in the 400-700 GeV range of the m_{bb} distribution, the data is found to exceed the LO prediction by up to two standard deviations, depending on the considered mass. This excess is no longer significant when NLO QCD corrections, as implemented in aMC@NLO [47], are included in the modelling of the Z+jets process. For this reason, the LO predictions are corrected using a reweighting technique, in order to account for NLO QCD effects. To this end, it becomes necessary to apply the reweighting according to the parton (or hadron) flavour of the jets in the generated event. The ratio NLO/LO of the light- and heavy-flavour components of the m_{jj} distribution is each fitted with a third-order polynomial and a separate reweighting of the shape of the light and heavy flavour components of m_{jj} is applied, resulting in better agreement with the data.

To determine the Z+jets and $t\bar{t}$ normalization, a data-based method is exploited. Data-derived correction factors for simulation are obtained after an additional categorisation of the Z+jet background events, based on the flavour (b jet or not) and multiplicity (exactly two jets or three or more jets) of the reconstructed jets. These categories are sensitive to NLO effects related to the modelling of extra jets [48]. Scale factors (SFs) are introduced for the $t\bar{t}$ background and the light and heavy flavour components of Z+jets background. These are left free to float in a two-dimensional fit of the distributions predicted by the simulation to the data. The distributions used as input are the product of the CSV discriminants of the two selected jets, and the invariant mass of the lepton pair from the Z boson decay in the range $60 < m_{ll} < 120$ GeV. The first observable is sensitive to the contribution from non-b jets, whereas the second one is sensitive to the contribution of the $t\bar{t}$ production process. The fit is performed simultaneously in four different categories: electrons, muons, exactly two jets, and more than two jets. The SF for the $t\bar{t}$ is found to be very close to the unity, while for the Z+jets process the SFs depart from unity by as much as 1.3 for the light flavour component.

The overall yields from diboson and tW processes are normalised to the CMS measurements [49–52]. The associated production of a Z boson together with the Higgs-like scalar boson (Zh) is also accounted for as background, and normalised to the expected theoretical cross section [53].

5.2 The $\tau\tau$ channel

Methods based on both data and simulation are used to estimate the residual background after event selection. Normalisations and mass distributions in the ZZ, Zh, as well as for the minor fully leptonic WWZ, WZZ, ZZZ and $t\bar{t}Z$ backgrounds are estimated from simulation. The Z+jets and WZ+jets contributions are measured by means of a data-based method.

Production of Z+jets and WZ+jets constitutes the main source of background when at least one lepton is misidentified. Misidentified light leptons arise from semileptonic decays of heavy-flavour quarks, decays in flight of hadrons, and

photon conversions, while jets originating from quarks or gluons can be misidentified as τh . Backgrounds with at least one misidentified lepton are estimated from control samples in data starting from the estimation of the lepton misidentification probabilities. The lepton misidentification probability is defined as the probability that a genuine jet, satisfying loose lepton identification criteria (which refer to the so-called “loose” lepton), also passes the identification criteria required for a lepton candidate in the signal region (so-called “tight” lepton). This probability is measured for each lepton flavour using a data sample where a Z candidate is selected, and an additional single lepton (electron, muon, or τh) passes the loose identification requirements. Counting the fraction of such loose leptons that also pass the tight lepton identification criteria in the p_T bins of the reconstructed jet closest, in ΔR , to the loose lepton, yields the misidentification probability f as a function of p_T . The contribution from genuine leptons arising from the WZ and ZZ production are subtracted.

Once the misidentification probabilities are computed, three control regions (CR) are defined with a Z candidate and two opposite-sign leptons, as follows: the CR00 wherein both leptons pass loose identification criteria but not the tight ones; CR10 region, wherein one lepton passes tight identification requirements, the other only loose criteria, and the loose lepton is the τh with lower p_T in the $\tau h \tau h$ channel, the light lepton in the τh channels, and the electron in the $e\mu$ channel; the CR01 region, which is similar to CR10 but the loose lepton is the τh with higher p_T in the $\tau h \tau h$ channel, the τh in the τh channels, and the muon in the $e\mu$ channel. The estimated N_{misID} of the background with at least one misidentified lepton from a pair of closest-jet p_T bins is given by:

$$N_{\text{misID}} = N_{10}/(1 - f_1) + N_{01}/(1 - f_2) - N_{00} f_1 f_2/(1 - f_1)(1 - f_2)$$

where N_{00} , N_{01} , and N_{10} denote the number of events from the CR00, CR01, and CR10 control regions, respectively, with closest jets in the considered p_T bins, and f_1 and f_2 indicate the misidentification probabilities associated with the two different flavor (except for the $\tau h \tau h$ final state) loose leptons in the p_T bins. The expression in Eq. (1) takes into account both the background with two misidentified leptons (mostly from Z+jets) and that from only one misidentified lepton (primarily from WZ+jets).

The contamination from genuine leptons in the control regions from the SM Zh, WWZ, WZZ, ZZZ, $t\bar{t}Z$, and ZZ processes is estimated from simulation, and subtracted from N_{00} , N_{01} , and N_{10} . The total background in the signal region is obtained by summing the contributions from all pairs of p_T bins.

6 Systematic uncertainties

The systematic uncertainties are reported in the following paragraphs and summarised in Tab. 1.

The uncertainty on the integrated luminosity recorded by CMS is estimated to be 2.6% [54]. The systematic uncertainties associated with the lepton efficiency

SFs, used to correct the simulation and derived from studies at the Z peak using the tag-and-probe (T&P) method [19, 20], are approximately 1% for muons and 2% for electrons, and affect both signal and background processes in the same way. Also, the uncertainties on the double muon and double electron trigger efficiencies are evaluated to be 1% from similar studies at the Z peak [22].

The uncertainty on the jet energy scale is derived from the method of Ref. [55] and the parameters describing the shape of the energy distribution are varied by one standard deviation (SD). The effect is estimated separately on the background and on the signal, resulting in a 3–5% variation, depending on the p_T and η of the jets. The uncertainty on signal and background yields induced by the imperfect knowledge of the jet energy resolution is estimated to be 3% [55].

The uncertainties affecting b tagging efficiencies are p_T -dependent, and vary from 3% to 12% (for $p_T > 30$ GeV) [42]. The impact of these uncertainties on the normalisation of signal is 5% for background and 4–6% for signal in the $b\bar{b}$ analysis, and about 1% in the $\tau\tau$ analysis. The uncertainty in the mistagging rate is found to have a negligible impact.

The systematic uncertainty on the signal is evaluated by varying the set of parton distribution functions (PDFs) according to the PDF4LHC prescriptions [56–58] and the factorisation and renormalisation scales by varying their values by a factor one half and two. An effect of 5–6% is estimated for the entire mass range for both $\tau\tau$ and $b\bar{b}$ final states.

Finally, an 11% uncertainty is assigned to the ZZ normalisation from the cross section measured by CMS [49].

For the $b\bar{b}$ final state, the uncertainty on the SFs used for normalisation of Z+jets and $t\bar{t}$ backgrounds is derived from the statistical uncertainty resulting from the fit used to derive these SFs and it is estimated to be $<8\%$. An additional systematic uncertainty associated with the $m_{b\bar{b}}$ spectrum correction, described in Sec. 5, ranges from 5% for $m_{b\bar{b}}$ below 700 GeV to 30% for masses at the TeV scale. An uncertainty of 8% is assigned to the normalisation of the WW process, corresponding to the uncertainties in the cross section measured by CMS [50]. A similar uncertainty is assigned also to the WZ process, which shares the same sources of uncertainties in the cross section measurement. For the minor tW background, the uncertainty is estimated as 23%, also based on the measured cross section [52]. The uncertainty in the Zh process is based on the theoretical cross section and found to be 7% [53].

In order to interpolate smoothly the signal efficiency across the parameter space, additional mass points for the $b\bar{b}$ final state are processed using a parametric simulation [59], tuned for delivering a realistic approximation of the CMS response in the reconstruction of physics objects used in this search. For this reason, an additional source of uncertainty is introduced for the SF applied to these samples to reproduce the efficiency measured with the full simulation. This is measured for the different signal points in the m_H - m_A plane and it is

close to 3% in most of the phase space, but rises to 50% at the boundaries of the sensitivity region.

In the $\tau\tau$ final state, the uncertainty of 6% [37] in the τh identification efficiency, which has been determined using a T&P method, has been taken into account. The τh energy scale uncertainty is within 3% [37] and only affects the shapes of the $\tau\tau$ mass distributions. The systematic uncertainties estimated for e , μ , τh and jet energy scales are propagated to $p_{\text{miss}T}$ and to the mass distributions. The propagation to $p_{\text{miss}T}$ involves a sum of the energies of each object first and a consequent subtraction of such contributions once the nominal energy scales (or resolutions) are varied up and down by one SD (for e , μ , τh , and jets). One of the main systematic uncertainties is related to the nonprompt background estimation. This uncertainty has been evaluated using simulation by comparing the direct estimate of the backgrounds with that obtained using the procedure adopted in the analysis, but applied to simulated events. The discrepancy between the two estimates never exceeds 40%. This value is thus considered as the uncertainty on the estimates of the reducible background yield for all channels and all LT thresholds.

7 Results

The analysis searches for new resonance decays by comparing data to simulation in the two-dimensional plane defined by the four-body (m_{bb} or $m_{\tau\tau}$) and two-body (m_{bb} or $m_{\tau\tau}$) invariant masses. The numerical values for the upper limits or the significance of a local excess are obtained using the asymptotic method described in Ref. [60]. The CLs method [61, 62] is used to determine the 95% confidence level (CL) upper limits on the excluded signal cross section.

For both final states, the limits in the lower-right triangle of the mass plane, which corresponds to the process $A \rightarrow ZH$, are obtained by mirroring the results obtained in the upper-left triangle, since the signal efficiencies for $H \rightarrow ZA$ and $A \rightarrow ZH$ are equal for the same masses of the heavy and light Higgs bosons in the two processes.

7.1 The bb channel

For the bb final state, results are obtained using a counting approach, which can be reinterpreted in other theoretical models with the same final state. Results are reported in bins of m_{bb} and m_{bb} masses, in the range from 10 GeV to 1 TeV for m_{bb} , and from 140 GeV to 1 TeV for m_{bb} . To define the proper granularity of the binning, a study is performed using signal benchmark points and evaluating the width of the m_{bb} and m_{bb} peaks in the considered mass range. The average reconstructed width, defined as one SD, for m_{bb} and m_{bb} is found to be approximately 15% of the considered mass. The bin widths have been chosen to be ± 1.5 SD around each considered mass point.

The efficiency, defined as the fraction of generated signal events reconstructed after the final selection, is calculated with the full CMS simulation and recon-

struction software at 13 representative signal points in the mH - mA mass plane. The signal efficiencies for the rest of the plane are obtained by interpolating the ratio between the full simulation and the parametric simulation (typically 0.9), calculated in each of the 13 signal mass points, and scaling the efficiencies calculated using the parametric simulation by this interpolated ratio. The resulting signal efficiency ranges from 8% at $(mA, mH) = (100, 300)$ GeV to 13% at $(300, 600)$ GeV.

Figure 1 [Figure 1: see original paper] shows the observed upper limits on the product of the cross section (σ) and branching fraction (B) for the $b\bar{b}$ final state in the mH - mA plane. The achieved sensitivity provides an exclusion limit at 95% CL of approximately 10 fb for a large fraction of the two-dimensional mass plane. In particular, the observed limit ranges from just above 1 fb for mH close to 1 TeV to 100 fb for $mH < 300$ GeV. The validity of these results is applicable to models allowing the existence of both A and H bosons with a natural width smaller than 15% of their masses.

Two moderate excesses are observed for the $b\bar{b}$ channel in the regions around $(m_{bb}, m_{bb}) = (95, 285)$ GeV and $(575, 660)$ GeV. According to the procedure described at Ref. [63], they have local significances of 2.6 and 2.85 SD respectively, which become globally 1.6 and 1.9 SD, once accounting for the look-elsewhere effect [64]. The low-mass excess is more compatible with the signal hypothesis, both in terms of yield and width. The reconstructed invariant mass distributions for the $b\bar{b}$ and $b\bar{b}$ systems, in the regions around this excess, are reported in Fig. 2 [Figure 2: see original paper] and compared with the expectations from background processes. A 2HDM type-II benchmark signal at $mH = 270$ GeV and $mA = 104$ GeV, normalised to the NNLO SUSHI prediction, is also superimposed.

7.2 The $\tau\tau$ channel

In the context of the $\tau\tau$ analysis, a search based on the $m_{\tau\tau}$ distribution is performed. For every considered pair of H and A mass values, the search is performed in eight $\tau\tau$ SVFIT binned mass distributions, each corresponding to one of the eight considered final states. Variable bin widths are adopted in order to account for the mass resolution. A simultaneous likelihood fit to the observed distributions is performed with the expected distributions from the background-only and signal plus background hypotheses. The normalisation of the signal distribution is a free parameter in the fit. No significant deviations are observed in data from the SM expectation. The SVFIT mass distributions of the $\tau\tau$ pair in the eight different final states are shown in Fig. 3 [Figure 3: see original paper]. The chosen signal corresponds to $mH = 350$ GeV and $mA = 90$ GeV, which is the one closest to the centre of the bin in which the highest excess is observed in the $b\bar{b}$ channel. The shown shapes correspond to $LT > 40$ GeV for $e\mu$, $LT > 60$ GeV for $e\tau h$ and $\mu\tau h$, and $LT > 80$ GeV for $\tau h\tau h$.

Figure 4 [Figure 4: see original paper] shows the limit on σB for the $\tau\tau$ final

state in the mH - mA plane. Signal cross sections of about 5–10 fb are excluded in most of the mH - mA plane ($500 < mH/A < 1000$ GeV and $90 < mA/H < 400$ GeV).

7.3 Combination in the context of 2HDM

Observed and expected upper limits on the signal cross section modifier $\mu = \sigma/\sigma_{\text{th}}$ are also derived and reported in Fig. 5 [Figure 5: see original paper], where σ_{th} is the theory cross section of the 2HDM benchmark used in this analysis. The results are obtained from the combination of the bb and $\tau\tau$ final states. This search is not able to exclude the high-mass regions where $mH > 300$ GeV and $mA > 300$ GeV, due to the drop in the signal cross section, where the $A/H \rightarrow t\bar{t}$ channel opens up for $mH > 2m_t$, where m_t is the top quark mass [16]. Furthermore, in the region where highly-boosted topologies start contributing ($mH \sim 10 mA$), the sensitivity is lower relative to the rest of the plane, primarily caused by the inefficiency in reconstructing signal decay products in such a regime. Still, a significant portion of the benchmark masses is excluded for a 2HDM type-II scenario with $\tan \beta = 1.5$ and $\cos(\beta - \alpha) = 0.01$, delimited by the solid contour in Fig. 5. The observed 95% CL exclusion region is localised in the range $mH = 200$ – 700 GeV and $mA = 20$ – 270 GeV for the decay $H \rightarrow ZA$, and similarly in the range $mH = 120$ – 270 GeV and $mA = 200$ – 700 GeV for the $A \rightarrow ZH$ decay. The region where $|mH - mA| < mZ$ is kinematically inaccessible.

The limits on μ can be also visualised as a function of the 2HDM parameters $\tan \beta$ and $\cos(\beta - \alpha)$ for a given pair of mH and mA , from the combination of bb and $\tau\tau$ final states. Results are given in Fig. 6 [Figure 6: see original paper], where the exclusion limits on the parameters are shown for $mH = 378$ GeV and $mA = 188$ GeV, a mass pair chosen to be within the exclusion region of Fig. 5. The area contained within the solid line shows the parameter space excluded for the chosen mass pair, where $\tan \beta$ lies between 0.5 and 2.3 and $\cos(\beta - \alpha)$ between -0.7 and 0.3 .

8 Summary

The paper describes the first CMS search for a new resonance decaying into a lighter resonance and a Z boson. Two searches have been performed, targeting the decay of the lighter resonance into either a pair of oppositely charged τ leptons or a bb pair. The Z boson is identified via its decays to electrons or muons. The search is based on data corresponding to an integrated luminosity of 19.8 fb^{-1} in proton–proton collisions at $\sqrt{s} = 8$ TeV. Deviations from the SM expectations are observed with a global significance of less than 2 SD and upper limits on the product of cross section and branching fraction are set. The search excludes σB as low as 5 fb and 1 fb for the bb and $\tau\tau$ final states, respectively, depending on the light and heavy resonance mass values.

Limits are also set on the mass parameters of the type-II 2HDM model that

predicts the processes $H \rightarrow ZA$ and $A \rightarrow ZH$, where H and A are CP-even and CP-odd scalar bosons, respectively. Combining the bb and $\tau\tau$ final states, the specific model corresponding to the parameter choice $\cos(\beta - \alpha) = 0.01$ and $\tan \beta = 1.5$ is excluded for m_H in the range 200–700 GeV and m_A in the range 20–270 GeV with $m_H > m_A$, or alternatively for m_A in the range 200–700 GeV and m_H in the range 120–270 GeV with $m_A > m_H$.

Limits on the signal cross section modifier are also derived as a function of $\tan \beta$ and $\cos(\beta - \alpha)$ parameters. As a result, for specific m_H - m_A mass values, a fairly large region in the parameter space $\tan \beta$ vs. $\cos(\beta - \alpha)$ is excluded. This covers a region unexplored so far, that cannot be probed by studying production and decay modes of the SM-like Higgs boson. In particular, for $m_H = 378$ GeV and $m_A = 188$ GeV, a range where $\tan \beta$ lies between 0.5 and 2.3 and $\cos(\beta - \alpha)$ between -0.7 and 0.3 is excluded, after the combination of the bb and $\tau\tau$ final states.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centres and personnel of the Worldwide LHC Computing Grid for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC and the CMS detector provided by the following funding agencies: BMWFW and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, and FAPESP (Brazil); MES (Bulgaria); CERN; CAS, MoST, and NSFC (China); COLCIENCIAS (Colombia); MSES and CSF (Croatia); RPF (Cyprus); MoER, ERC IUT and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRT (Greece); OTKA and NIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); LAS (Lithuania); MOE and UM (Malaysia); CINVESTAV, CONACYT, SEP, and UASLP-FAI (Mexico); MBIE (New Zealand); PAEC (Pakistan); MSHE and NSC (Poland); FCT (Portugal); JINR (Dubna); MON, RosAtom, RAS and RFBR (Russia); MESTD (Serbia); SEIDI and CPAN (Spain); Swiss Funding Agencies (Switzerland); MST (Taipei); ThEPCenter, IPST, STAR and NSTDA (Thailand); TUBITAK and TAEK (Turkey); NASU and SFFR (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie programme and the European Research Council and EPLANET (European Union); the Leventis Foundation; the A. P. Sloan Foundation; the Alexander von Humboldt Foundation; the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium);

the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Council of Science and Industrial Research, India; the HOMING PLUS programme of the Foundation for Polish Science, cofinanced from European Union, Regional Development Fund; the OPUS programme of the National Science Center (Poland); the Compagnia di San Paolo (Torino); MIUR project 20108T4XTM (Italy); the Thalís and Aristeia programmes cofinanced by EU-ESF and the Greek NSRF; the National Priorities Research Program by Qatar National Research Fund; the Rachadapisek Sompot Fund for Postdoctoral Fellowship, Chulalongkorn University (Thailand); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project (Thailand); and the Welch Foundation, contract C-1845.

References

- [1] ATLAS Collaboration, “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC”, *Phys. Lett. B* 716 (2012) 1, doi:10.1016/j.physletb.2012.08.020, arXiv:1207.7214.
- [2] CMS Collaboration, “Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC”, *Phys. Lett. B* 716 (2012) 30, doi:10.1016/j.physletb.2012.08.021.
- [3] CMS Collaboration, “Observation of a new boson with mass near 125 GeV in pp collisions at $\sqrt{s} = 7$ and 8 TeV”, *JHEP* 06 (2013) 081, doi:10.1007/JHEP06(2013)081, arXiv:1303.4571.
- [4] CMS Collaboration, “Evidence for the direct decay of the 125 GeV Higgs boson to fermions”, *Nature Phys.* 10 (2014) 557, doi:10.1038/nphys3005.
- [5] ATLAS Collaboration, “Search for the $b\bar{b}$ decay of the Standard Model Higgs boson in associated (W/Z)H production with the ATLAS detector”, *JHEP* 01 (2015) 069, doi:10.1007/JHEP01(2015)069, arXiv:1409.6212.
- [6] CMS Collaboration, “Precise determination of the mass of the Higgs boson and tests of compatibility of its couplings with the standard model predictions using proton collisions at 7 and 8 TeV”, *Eur. Phys. J. C* 75 (2015) 212, doi:10.1140/epjc/s10052-015-3351-7, arXiv:1412.8662.
- [7] G. C. Branco et al., “Theory and phenomenology of two-Higgs-doublet models”, *Phys. Rep.* 516 (2012) 1, doi:10.1016/j.physrep.2012.02.002, arXiv:1106.0034.
- [8] G. Dorsch, S. Huber, K. Mimasu, and J. No, “Echoes of the Electroweak Phase Transition: Discovering a second Higgs doublet through $A0 \rightarrow ZH0$ ”, *Phys. Rev. Lett.* 113 (2014) 211802, doi:10.1103/PhysRevLett.113.211802, arXiv:1405.5537.
- [9] S. P. Martin, “A Supersymmetry Primer”, *Adv. Ser. Direct. High Energy Phys.* 18 (1998) 1, doi:10.1142/9789812839657_0001, arXiv:hep-ph/9709356.

- [10] E. Bagnaschi et al., “Benchmark scenarios for low $\tan \beta$ in the MSSM”, Technical Report LHCHXSWG-2015-002, CERN, Geneva, 2015.
- [11] J. E. Kim, “Light Pseudoscalars, Particle Physics and Cosmology”, Phys. Rep. 150 (1987) 1, doi:10.1016/0370-1573(87)90017-2.
- [12] A. Broggio et al., “Limiting two-Higgs-doublet models”, JHEP 11 (2014) 058, doi:10.1007/JHEP11(2014)058, arXiv:1409.3199.
- [13] F. Jegerlehner and A. Nyffeler, “The Muon $g-2$ ”, Phys. Rep. 477 (2009) 1, doi:10.1016/j.physrep.2009.04.003, arXiv:0902.3360.
- [14] ATLAS, CMS Collaboration, “Combined Measurement of the Higgs Boson Mass in pp Collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS and CMS Experiments”, Phys. Rev. Lett. 114 (2015) 191803, doi:10.1103/PhysRevLett.114.191803, arXiv:1503.07589.
- [15] J. M. Gerard and M. Herquet, “A Twisted custodial symmetry in the two-Higgs-doublet model”, Phys. Rev. Lett. 98 (2007) 251802, doi:10.1103/PhysRevLett.98.251802, arXiv:hep-ph/0703051.
- [16] B. Grinstein and P. Uttayarat, “Carving Out Parameter Space in Type-II Two Higgs Doublets Model”, JHEP 06 (2013) 094, doi:10.1007/JHEP06(2013)094, arXiv:1304.0028. [Erratum: JHEP 09 (2013) 110].
- [17] CMS Collaboration, “Search for a pseudoscalar boson decaying into a Z boson and the 125 GeV Higgs boson in $+ -bb$ final states”, Phys. Lett. B 748 (2015) 221, doi:10.1016/j.physletb.2015.07.010, arXiv:1504.04710.
- [18] ATLAS Collaboration, “Search for a CP-odd Higgs boson decaying to Zh in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Lett. B 744 (2015) 163, doi:10.1016/j.physletb.2015.03.054, arXiv:1502.04478.
- [19] CMS Collaboration, “Performance of electron reconstruction and selection with the CMS detector in proton-proton collisions at $\sqrt{s} = 8$ TeV”, J. Instrum. 10 (2015) P06005, doi:10.1088/1748-0221/10/06/P06005, arXiv:1502.02701.
- [20] CMS Collaboration, “Performance of CMS muon reconstruction in pp collision events at $\sqrt{s} = 7$ TeV”, J. Instrum. 7 (2012) P10002, doi:10.1088/1748-0221/7/10/P10002.
- [21] CMS Collaboration, “The CMS experiment at the CERN LHC”, J. Instrum. 3 (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.
- [22] CMS Collaboration, “Measurement of inclusive W and Z boson production cross sections in pp collisions at $\sqrt{s} = 8$ TeV”, Phys. Rev. Lett. 112 (2014) 191802, doi:10.1103/PhysRevLett.112.191802, arXiv:1402.0923.
- [23] S. Alioli, P. Nason, C. Oleari, and E. Re, “A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX”, JHEP 06 (2010) 043, doi:10.1007/JHEP06(2010)043, arXiv:1002.2581.

- [24] J. Alwall et al., “MadGraph5: going beyond”, JHEP 06 (2011) 128, doi:10.1007/JHEP06(2011)128, arXiv:1106.0522.
- [25] T. Sjöstrand, S. Mrenna, and P. Skands, “PYTHIA 6.4 physics and manual”, JHEP 05 (2006) 026, doi:10.1088/1126-6708/2006/05/026, arXiv:hep-ph/0603175.
- [26] CMS Collaboration, “Measurement of the Underlying Event Activity at the LHC with $\sqrt{s} = 7$ TeV and Comparison with $\sqrt{s} = 0.9$ TeV”, JHEP 09 (2011) 109, doi:10.1007/JHEP09(2011)109, arXiv:1107.0330.
- [27] S. Jadach, Z. Was, R. Decker, and J. H. Kuhn, “The tau decay library TAUOLA: Version 2.4”, Comput. Phys. Commun. 76 (1993) 361, doi:10.1016/0010-4655(93)90061-G.
- [28] GEANT4 Collaboration, “GEANT4—a simulation toolkit”, Nucl. Instrum. Meth. A 506 (2003) 250, doi:10.1016/S0168-9002(03)01368-8.
- [29] J. F. Gunion and H. E. Haber, “The CP conserving two Higgs doublet model: The approach to the decoupling limit”, Phys. Rev. D 67 (2003) 075019, doi:10.1103/PhysRevD.67.075019, arXiv:hep-ph/0207010.
- [30] R. V. Harlander, S. Liebler, and H. Mantler, “SusHi: A program for the calculation of Higgs production in gluon fusion and bottom-quark annihilation in the Standard Model and the MSSM”, Comput. Phys. Commun. 184 (2013) 1605, doi:10.1016/j.cpc.2013.02.006, arXiv:1212.3249.
- [31] D. Eriksson, J. Rathsman, and O. Stal, “2HDMC: Two-Higgs-Doublet Model Calculator Physics and Manual”, Comput. Phys. Commun. 181 (2010) 189, doi:10.1016/j.cpc.2009.09.011, arXiv:0902.0851.
- [32] R. Harlander et al., “Interim recommendations for the evaluation of Higgs production cross sections and branching ratios at the LHC in the Two-Higgs-Doublet Model”, (2013). arXiv:1312.5571.
- [33] B. Hespel, F. Maltoni, and E. Vryonidou, “Higgs and Z boson associated production via gluon fusion in the SM and the 2HDM”, JHEP 06 (2015) 065, doi:10.1007/JHEP06(2015)065, arXiv:1503.01656.
- [34] CMS Collaboration, “Particle-Flow Event Reconstruction in CMS and Performance for Jets, Taus, and MET”, CMS Physics Analysis Summary CMS-PAS-PFT-09-001, 2009.
- [35] CMS Collaboration, “Commissioning of the Particle-flow Event Reconstruction with the first LHC collisions recorded in the CMS detector”, CMS Physics Analysis Summary CMS-PAS-PFT-10-001, 2010.
- [36] CMS Collaboration, “Description and performance of track and primary-vertex reconstruction with the CMS tracker”, J. Instrum. 9 (2014) P10009, doi:10.1088/1748-0221/9/10/P10009.

- [37] CMS Collaboration, “Reconstruction and identification of τ lepton decays to hadrons and τ at CMS”, *J. Instrum.* (2016) P01019.
- [38] M. Cacciari, G. P. Salam, and G. Soyez, “The anti-kt jet clustering algorithm”, *JHEP* 04 (2008) 063, doi:10.1088/1126-6708/2008/04/063, arXiv:0802.1189.
- [39] M. Cacciari, G. P. Salam, and G. Soyez, “FastJet user manual”, *Eur. Phys. J. C* 72 (2012) 1896, doi:10.1140/epjc/s10052-012-1896-2, arXiv:1111.6097.
- [40] CMS Collaboration, “Pileup Removal Algorithms”, CMS Physics Analysis Summary CMS-PAS-JME-14-001, 2014.
- [41] M. Cacciari and G. P. Salam, “Pileup subtraction using jet areas”, *Phys. Lett. B* 659 (2008) 119, doi:10.1016/j.physletb.2007.09.077, arXiv:0707.1378.
- [42] CMS Collaboration, “Identification of b-quark jets with the CMS experiment”, *J. Instrum.* 8 (2013) 04013, doi:10.1088/1748-0221/8/04/P04013, arXiv:1211.4462.
- [43] Particle Data Group Collaboration, “Review of Particle Physics”, *Chin. Phys. C* 38 (2014) 090001, doi:10.1088/1674-1137/38/9/090001.
- [44] CMS Collaboration, “Performance of the CMS missing transverse momentum reconstruction in pp data at $\sqrt{s} = 8$ TeV”, *J. Instrum.* 10 (2015) P02006, doi:10.1088/1748-0221/10/02/P02006, arXiv:1411.0511.
- [45] CMS Collaboration, “Missing transverse energy performance of the CMS detector”, *J. Instrum.* 6 (2011) P09001, doi:10.1088/1748-0221/6/09/P09001.
- [46] L. Bianchini, J. Conway, E. K. Friis, and C. Veelken, “Reconstruction of the Higgs mass in $H \rightarrow \tau\tau$ Events by Dynamical Likelihood techniques”, *J. Phys. Conf. Ser.* 513 (2014) 022035, doi:10.1088/1742-6596/513/2/022035.
- [47] J. Alwall et al., “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations”, *JHEP* 07 (2014) 079, doi:10.1007/JHEP07(2014)079, arXiv:1405.0301.
- [48] CMS Collaboration, “Measurement of the production cross sections for a Z boson and one or more b jets in pp collisions at $\sqrt{s} = 7$ TeV”, *JHEP* 06 (2014) 120, doi:10.1007/JHEP06(2014)120, arXiv:1402.1521.
- [49] CMS Collaboration, “Measurement of the pp $\rightarrow ZZ$ production cross section and constraints on anomalous triple gauge couplings in four-lepton final states at $\sqrt{s} = 8$ TeV”, *Phys. Lett. B* 740 (2015) 250, doi:10.1016/j.physletb.2014.11.059, arXiv:1406.0113.
- [50] CMS Collaboration, “Measurement of the $W+W-$ cross section in pp collisions at $\sqrt{s} = 8$ TeV and limits on anomalous gauge couplings”, (2015). arXiv:1507.03268. Submitted to *Eur. Phys. J. C*.

- [51] CMS Collaboration, “Measurement of WZ and ZZ production in pp collisions at $\sqrt{s} = 8$ TeV in final states with b-tagged jets”, *Eur. Phys. J. C* 74 (2014) 2973, doi:10.1140/epjc/s10052-014-2973-5.
- [52] CMS Collaboration, “Observation of the associated production of a single top quark and a W boson in pp collisions at $\sqrt{s} = 8$ TeV”, *Phys. Rev. Lett.* 112 (2014) 231802, doi:10.1103/PhysRevLett.112.231802, arXiv:1401.2942.
- [53] CMS Collaboration, “Search for the standard model Higgs boson produced in association with a W or a Z boson and decaying to bottom quarks”, *Phys. Rev. D* 89 (2014) 012003, doi:10.1103/PhysRevD.89.012003, arXiv:1310.3687.
- [54] CMS Collaboration, “CMS Luminosity Based on Pixel Cluster Counting - Summer 2013 Update”, CMS Physics Analysis Summary CMS-PAS-LUM-13-001, 2013.
- [55] CMS Collaboration, “Determination of Jet Energy Calibration and Transverse Momentum Resolution in CMS”, *J. Instrum.* 6 (2011) P11002, doi:10.1088/1748-0221/6/11/P11002, arXiv:1107.4277.
- [56] M. Botje et al., “The PDF4LHC Working Group Interim Recommendations”, (2011). arXiv:1101.0538.
- [57] S. Alekhin et al., “The PDF4LHC Working Group Interim Report”, (2011). arXiv:1101.0536.
- [58] NNPDF Collaboration, “Parton distributions with LHC data”, *Nucl. Phys. B* 867 (2013) 244, doi:10.1016/j.nuclphysb.2012.10.003, arXiv:1207.1303.
- [59] DELPHES 3 Collaboration, “DELPHES 3, A modular framework for fast simulation of a generic collider experiment”, *JHEP* 02 (2014) 057, doi:10.1007/JHEP02(2014)057, arXiv:1307.6346.
- [60] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, “Asymptotic formulae for likelihood-based tests of new physics”, *Eur. Phys. J. C* 71 (2011) 1554, doi:10.1140/epjc/s10052-011-1554-0, arXiv:1007.1727. [Erratum: *Eur. Phys. J. C* 73 (2013) 7].
- [61] T. Junk, “Confidence level computation for combining searches with small statistics”, *Nucl. Instrum. Meth. A* 434 (1999) 435, doi:10.1016/S0168-9002(99)00498-2, arXiv:hep-ex/9902006.
- [62] A. L. Read, “Presentation of search results: The CL(s) technique”, *J. Phys. G* 28 (2002) 2693, doi:10.1088/0954-3899/28/10/313.
- [63] ATLAS Collaboration, CMS Collaboration, LHC Higgs Combination Group Collaboration, “Procedure for the LHC Higgs boson search combination in Summer 2011”, Technical Report CMS-NOTE-2011-005, ATL-PHYS-PUB-2011-11, 2011.
- [64] E. Gross and O. Vitells, “Trial factors or the look elsewhere effect in high

energy physics”, Eur. Phys. J. C 70 (2010) 525, doi:10.1140/epjc/s10052-010-1470-8, arXiv:1005.1891.

Appendix A: The CMS Collaboration

[The full author list appears in the original document]

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv — Machine translation. Verify with original.