

A New Measurement of Antineutrino Oscillation with the Full Detector Configuration at Daya Bay postprint

Authors: An, F. P. and others

Date: 2016-09-09T00:00:00+00:00

Abstract

We report a new measurement of electron antineutrino disappearance using the fully-constructed Daya Bay Reactor Neutrino Experiment. The final two of eight antineutrino detectors were installed in the summer of 2012. Including the 404 days of data collected from October 2012 to November 2013 resulted in a total exposure of 6.9×10^5 GWth-ton-days, a 3.6 times increase over our previous results. Improvements in energy calibration limited variations between detectors to 0.2%. Removal of six ^{241}Am - ^{13}C radioactive calibration sources reduced the background by a factor of two for the detectors in the experimental hall furthest from the reactors. Direct prediction of the antineutrino signal in the far detectors based on the measurements in the near detectors explicitly minimized the dependence of the measurement on models of reactor antineutrino emission. The uncertainties in our estimates of $\sin^2 2\theta_{13}$ and $|\Delta m_{2ee}^2|$ were halved as a result of these improvements. Analysis of the relative antineutrino rates and energy spectra between detectors gave $\sin^2 2\theta_{13} = 0.084 \pm 0.005$ and $|\Delta m_{2ee}^2| = (2.42 \pm 0.11) \times 10^{-3} \text{ eV}^2$ in the three-neutrino framework.

Full Text

Preamble

A new measurement of antineutrino oscillation with the full detector configuration at Daya Bay

F. P. An, A. B. Balantekin, H. R. Band, M. Bishai, S. Blyth, I. Butorov, G. F. Cao, J. Cao, W. R. Cen, Y. L. Chan, J. F. Chang, L. C. Chang, Y. Chang, H. S. Chen, Q. Y. Chen, S. M. Chen, Y. X. Chen, Y. Chen, J. H. Cheng, J. Cheng, Y. P. Cheng, J. J. Cherwinka, M. C. Chu, J. P. Cummings, J. de Arcos, Z. Y. Deng, X. F. Ding, Y. Y. Ding, M. V. Diwan, E. Draeger, D. A. Dwyer, W. R. Edwards, S. R. Ely, R. Gill, M. Gonchar, G. H. Gong, H. Gong, M. Grassi, W.

Q. Gu, M. Y. Guan, L. Guo, X. H. Guo, R. W. Hackenburg, R. Han, S. Hans, M. He, K. M. Heeger, Y. K. Heng, A. Higuera, Y. K. Hor, Y. B. Hsiung, B. Z. Hu, L. M. Hu, L. J. Hu, T. Hu, W. Hu, E. C. Huang, H. X. Huang, X. T. Huang, P. Huber, G. Hussain, D. E. Jaffe, P. Jaffke, K. L. Jen, S. Jetter, X. P. Ji, X. L. Ji, J. B. Jiao, R. A. Johnson, L. Kang, S. H. Kettell, M. Kramer, K. K. Kwan, M. W. Kwok, T. Kwok, T. J. Langford, K. Lau, L. Lebanowski, J. Lee, R. T. Lei, R. Leitner, A. Leung, J. K. C. Leung, C. A. Lewis, D. J. Li, F. Li, G. S. Li, Q. J. Li, S. C. Li, W. D. Li, X. N. Li, X. Q. Li, Y. F. Li, Z. B. Li, H. Liang, C. J. Lin, G. L. Lin, P. Y. Lin, S. K. Lin, J. J. Ling, J. M. Link, L. Littenberg, B. R. Littlejohn, D. W. Liu, H. Liu, J. L. Liu, J. C. Liu, S. S. Liu, C. Lu, H. Q. Lu, J. S. Lu, K. B. Luk, Q. M. Ma, X. Y. Ma, X. B. Ma, Y. Q. Ma, D. A. Martinez Caicedo, K. T. McDonald, R. D. McKeown, Y. Meng, I. Mitchell, J. Monari Kebwaro, Y. Nakajima, J. Napolitano, D. Naumov, E. Naumova, H. Y. Ngai, Z. Ning, J. P. Ochoa-Ricoux, A. Olshevski, J. Park, S. Patton, V. Pec, J. C. Peng, L. E. Pilonen, L. Pinsky, C. S. J. Pun, F. Z. Qi, M. Qi, X. Qian, N. Raper, B. Ren, J. Ren, R. Rosero, B. Roskovec, X. C. Ruan, B. B. Shao, H. Steiner, G. X. Sun, J. L. Sun, W. Tang, D. Taychenachev, H. Themann, K. V. Tsang, C. E. Tull, Y. C. Tung, N. Viaux, B. Viren, V. Vorobel, C. H. Wang, M. Wang, N. Y. Wang, R. G. Wang, W. Wang, W. W. Wang, X. Wang, Y. F. Wang, Z. Wang, Z. M. Wang, H. Y. Wei, L. J. Wen, K. Whisnant, C. G. White, L. Whitehead, T. Wise, H. L. H. Wong, S. C. F. Wong, E. Worcester, Q. Wu, D. M. Xia, J. K. Xia, X. Xia, Z. Z. Xing, J. Y. Xu, J. L. Xu, J. Xu, Y. Xu, T. Xue, J. Yan, C. G. Yang, L. Yang, M. S. Yang, M. T. Yang, M. Ye, M. Yeh, Y. S. Yeh, B. L. Young, G. Y. Yu, Z. Y. Yu, S. L. Zang, L. Zhan, C. Zhang, H. H. Zhang, J. W. Zhang, Q. M. Zhang, Y. M. Zhang, Y. X. Zhang, Y. M. Zhang, Z. J. Zhang, Z. Y. Zhang, Z. P. Zhang, J. Zhao, Q. W. Zhao, Y. F. Zhao, Y. B. Zhao, L. Zheng, W. L. Zhong, L. Zhou, N. Zhou, H. L. Zhuang, and J. H. Zou

(The Daya Bay Collaboration)

1Institute of Modern Physics, East China University of Science and Technology, Shanghai 2University of Wisconsin, Madison, Wisconsin, USA 3Department of Physics, Yale University, New Haven, Connecticut, USA 4Brookhaven National Laboratory, Upton, New York, USA 5Department of Physics, National Taiwan University, Taipei 6National United University, Miao-Li 7Joint Institute for Nuclear Research, Dubna, Moscow Region 8Institute of High Energy Physics, Beijing 9Chinese University of Hong Kong, Hong Kong 10Institute of Physics, National Chiao-Tung University, Hsinchu 11Shandong University, Jinan 12Department of Engineering Physics, Tsinghua University, Beijing 13North China Electric Power University, Beijing 14Shenzhen University, Shenzhen 15Siena College, Loudonville, New York, USA 16Department of Physics, Illinois Institute of Technology, Chicago, Illinois, USA 17Lawrence Berkeley National Laboratory, Berkeley, California, USA 18Department of Physics, University of Illinois at Urbana-Champaign, Urbana, Illinois, USA 19Shanghai Jiao Tong University, Shanghai 20Beijing Normal University, Beijing 21Department of Physics, University of Houston, Houston, Texas, USA 22Center for Neutrino Physics, Virginia Tech, Blacksburg, Virginia, USA 23China Institute of Atomic

Energy, Beijing 24School of Physics, Nankai University, Tianjin 25Department of Physics, University of Cincinnati, Cincinnati, Ohio, USA 26Dongguan University of Technology, Dongguan 27Department of Physics, University of California, Berkeley, California, USA 28Department of Physics, The University of Hong Kong, Pokfulam, Hong Kong 29Charles University, Faculty of Mathematics and Physics, Prague 30University of Science and Technology of China, Hefei 31Sun Yat-Sen (Zhongshan) University, Guangzhou 32Joseph Henry Laboratories, Princeton University, Princeton, New Jersey, USA 33California Institute of Technology, Pasadena, California, USA 34College of William and Mary, Williamsburg, Virginia, USA 35Xi'an Jiaotong University, Xi'an 36Department of Physics, College of Science and Technology, Temple University, Philadelphia, Pennsylvania, USA 37Instituto de Física, Pontificia Universidad Católica de Chile, Santiago, Chile 38Nanjing University, Nanjing 39Department of Physics, Applied Physics, and Astronomy, Rensselaer Polytechnic Institute, Troy, New York, USA 40China General Nuclear Power Group 41College of Electronic Science and Engineering, National University of Defense Technology, Changsha 42Iowa State University, Ames, Iowa 43Chongqing University, Chongqing

(Dated: September 11, 2015)

Abstract

We report a new measurement of electron antineutrino disappearance using the fully-constructed Daya Bay Reactor Neutrino Experiment. The final two of eight antineutrino detectors were installed in the summer of 2012. Including the 404 days of data collected from October 2012 to November 2013 resulted in a total exposure of 6.9×10^5 GWth-ton-days, a 3.6 times increase over our previous results. Improvements in energy calibration limited variations between detectors to 0.2%. Removal of six ^{241}Am - ^{13}C radioactive calibration sources reduced the background by a factor of two for the detectors in the experimental hall furthest from the reactors. Direct prediction of the antineutrino signal in the far detectors based on the measurements in the near detectors explicitly minimized the dependence of the measurement on models of reactor antineutrino emission.

The uncertainties in our estimates of $\sin^2 2\theta_{13}$ and $|\Delta m_{ee}^2|$ were halved as a result of these improvements. Analysis of the relative antineutrino rates and energy spectra between detectors gave $\sin^2 2\theta_{13} = 0.084 \pm 0.005$ and $|\Delta m_{ee}^2| = (2.42 \pm 0.11) \times 10^{-3} \text{ eV}^2$ in the three-neutrino framework.

Neutrino flavor oscillation due to the mixing angle θ_{13} has been observed using reactor antineutrinos [?, ?, ?] and accelerator neutrinos [?, ?]. The Daya Bay experiment previously reported the discovery of a non-zero value of $\sin^2 2\theta_{13}$ by observing the disappearance of reactor antineutrinos over kilometer distances [?, ?, ?], and the first measurement of the effective mass splitting $|\Delta m_{ee}^2|$ [?] via the distortion of the $\bar{\nu}_e$ energy spectrum [?]. Here we present new results with significant improvements in energy calibration and background reduction.

Installation of the final two detectors and a tripling of operation time provided a total exposure of 6.9×10^5 GWth-ton-days, 3.6 times more than reported in our previous publication [?]. With these improvements the precision of $\sin^2 2\theta_{13}$ was enhanced by a factor of two compared to the world's previous best estimate.

The precision of $|\Delta m_{ee}^2|$ was equally enhanced, and is now competitive with the precision of $|\Delta m_{32}^2|$ measured via accelerator neutrino disappearance [?, ?].

Experimental Setup and Data Collection

The Daya Bay experiment started collecting data on 24 December 2011 with six antineutrino detectors (ADs) located in three underground experimental halls (EHs). Three ADs were positioned in two near halls at short distances from six nuclear reactor cores (two ADs in EH1 and one in EH2), and three ADs were positioned in the far hall, EH3. Data taking was paused on 28 July 2012 while two new ADs were installed, one in EH2 and the other in EH3. During the installation, a broad set of calibration sources were deployed into the two ADs of EH1 using automated calibration units [?] and a manual calibration system [?]. Operation of the full experiment with all eight ADs started on 19 October 2012.

This Letter presents results based on 404 days of data acquired in the 8-AD period combined with all 217 days of data acquired in the 6-AD period. A blind analysis strategy was implemented by concealing the baselines and target masses of the two new ADs, as well as the operational data of all reactor cores for the new data period.

Each of the three Daya Bay experimental halls hosts functionally identical ADs inside a muon detector system. The muon detector system consists of a two-zone pure water Cherenkov detector, referred to as the inner and outer water shields (IWS and OWS), covered on top by an array of resistive plate chambers (RPCs). Each AD consists of nested cylindrical vessels. The inner vessel is filled with 0.1% gadolinium-doped liquid scintillator (Gd-LS), which constitutes the primary antineutrino target. The vessel surrounding the target is filled with undoped LS, increasing the efficiency of detecting gamma rays produced in the target. The outermost vessel is filled with mineral oil. A total of 192 20-cm photomultiplier tubes (PMTs) are radially positioned in the mineral-oil region of each AD. Further details on the experimental setup are contained in Refs. [?, ?, ?, ?].

Reactor antineutrinos are detected via the inverse β -decay (IBD) reaction, $\bar{\nu}_e + p \rightarrow e^+ + n$. The gamma rays (totaling 8 MeV) generated from the neutron capture on Gd with a mean capture time of 30 μ s form a delayed signal and enable powerful background suppression. The light from the e^+ gives an estimate of the incident $\bar{\nu}_e$ energy, $E_{\bar{\nu}_e} \approx E_p + E_n + 0.78$ MeV, where E_p is the prompt energy including the positron kinetic and annihilation energy, and E_n is the average neutron recoil energy (10 keV).

Energy Calibration and Reconstruction

The estimated energy response of the detectors to positrons, including both kinetic and annihilation gamma energy, is shown in Fig. 2 [Figure 2: see original paper] (red solid curve). The prominent nonlinearity below 4 MeV was attributed to scintillator light yield (from ionization quenching and Cherenkov light production) and the charge response of the electronics. Gamma rays from both deployed and intrinsic sources as well as spallation ^{12}B decay determined the model, and provided an envelope of curves consistent with the data within a 68.3% C.L. (grey band). An independent estimate using the beta+gamma energy spectra from ^{212}Bi , ^{214}Bi , ^{208}Tl , as well as the 53-MeV edge in the Michel electron spectrum gave a similar result (blue dashed line), albeit with larger systematic uncertainties.

Nonlinearity in the energy response of an AD originated from two dominant sources: particle-dependent nonlinear light yield of the scintillator and charge-dependent nonlinearity in the PMT readout electronics. Each effect was at the level of 10%. We constructed a semi-empirical model that predicted the reconstructed energy for a particle assuming a specific energy deposited in the scintillator. The model contained four parameters: Birks' constant, the relative contribution to the total light yield from Cherenkov radiation, and the amplitude and scale of an exponential correction describing the non-linear electronics response. This exponential form for the electronics response was motivated by Monte Carlo simulations and confirmed with an independent FADC measurement.

The nominal parameter values were obtained from an unconstrained χ^2 fit to various AD calibration datasets, comprising twelve gamma lines from both deployed and naturally occurring sources as well as the continuous β -decay spectrum of ^{12}B produced by muon spallation inside the Gd-LS volumes. The nominal positron response derived from the best fit parameters is shown in Fig. 2 [Figure 2: see original paper]. The depicted uncertainty band represents other response functions consistent with the fitted calibration data within a 68.3% C.L. This χ^2 -based approach to obtain the energy response resulted in $<1\%$ uncertainties of the absolute energy scale above 2 MeV. The uncertainties of the positron response were validated using the 53 MeV cutoff in the Michel electron spectrum from muon decay at rest and the continuous β^+ spectra from natural bismuth and thallium decays. These improvements added confidence in the characterization of the absolute energy response of the detectors, although they resulted in negligible changes to the measured mixing parameters.

PMT gains were calibrated continuously using uncorrelated single electrons emitted by the photocathode. The signals of 0.3% of the PMTs were discarded due to abnormal hit rates or charge distributions. The detector energy scale was calibrated using Am-C neutron sources [?] deployed at the detector center, with the 8 MeV peaks from neutrons captured on Gd aligned across all eight detectors. The time variation and the position dependence of the energy scale were

corrected using the 2.506 MeV gamma-ray peak from ^{60}Co calibration sources.

The reconstructed energies of various calibration reference points in different ADs are compared in Fig. 1 [Figure 1: see original paper]. The spatial distribution of each calibration reference varies, incorporating deviations in spatial response between detectors. Measurements of ^{68}Ge , ^{60}Co and Am-C calibration sources when placed at the center of each detector are presented. Neutrons from IBD and muon spallation that were captured on gadolinium were distributed nearly uniformly throughout the Gd-LS region. Those neutrons that were captured on ^1H , intrinsic particles from polonium and radon decays, and gammas from ^{40}K and ^{208}Tl decays were distributed inside and outside of the target volume. All of these events were selected within the Gd-LS region based on their reconstructed vertices. Systematic variations between detectors for all calibration references were $<0.2\%$. The 8 MeV n-Gd capture gamma peaks from Am-C sources were used to define the energy scale of each detector, and hence show zero deviation.

The uncorrelated relative uncertainty of the energy scale is thus determined to be 0.2%. This reduction of 43% compared to the previous publication [?] was enabled by improvements in the correction of position and time dependence, and enhanced the precision of $|\Delta m_{ee}^2|$ by 9%. The reduction was confirmed by an alternative method which used the n-Gd capture of muon-induced spallation neutrons to calibrate the scale, time dependence, and spatial dependence of the detector energy response.

Background Reduction and Event Selection

IBD candidates were selected using the same criteria discussed in Ref. [?]. Noise introduced by PMT light emission in the voltage divider, called flashing, was efficiently removed using the techniques of Ref. [?]. We required $0.7 \text{ MeV} < E_p < 12.0 \text{ MeV}$, $6.0 \text{ MeV} < E_d < 12.0 \text{ MeV}$, and $1 \mu\text{s} < \Delta t < 200 \mu\text{s}$, where E_d is the delayed energy and $\Delta t = t_d - t_p$ is the time difference between the prompt and delayed signals. To suppress cosmogenic products, candidates were rejected if their delayed signal occurred (i) within a $(-2 \mu\text{s}, 600 \mu\text{s})$ time window with respect to an IWS or OWS trigger with a PMT multiplicity > 12 , (ii) within a $(-2 \mu\text{s}, 1000 \mu\text{s})$ time window with respect to triggers in the same AD with reconstructed energy $> 20 \text{ MeV}$, or (iii) within a $(-2 \mu\text{s}, 1 \text{ s})$ time window with respect to triggers in the same AD with reconstructed energy $> 2.5 \text{ GeV}$. To select only definite signal pairs, we required the signal to have a multiplicity of 2: no other $> 0.7 \text{ MeV}$ signal occurred within a $(t_p - 200 \mu\text{s}, t_d + 200 \mu\text{s})$ time window.

Estimates for the five major sources of background for the new data sample are improved with respect to Ref. [?]. The background produced by the three Am-C neutron sources inside the automated calibration units contributed significantly to the total systematic uncertainty of the correlated backgrounds in the 6-AD period. Because of this, two of the three Am-C sources in each AD in EH3 were

removed during the 2012 summer installation period. As a result, the average correlated Am-C background rate in the far hall decreased by a factor of 4 in the 8-AD period. As in previous publications [?, ?], this rate was determined by monitoring the single neutron production rate from the Am-C sources. Removal of these Am-C sources had negligible consequences for our calibration.

Energetic, or fast, neutrons of cosmogenic origin produced a correlated background for this study. Relaxing the prompt-energy selection to (0.7–100) MeV revealed the fast-neutron background spectrum above 12 MeV. Previously we deduced the rate and spectrum of this background using a linear extrapolation into the IBD prompt signal region. Here we used a background-enhanced dataset to improve the estimate. We found 6043 fast neutron candidates with prompt energy from 0.7 to 100 MeV in the 200 μ s following cosmogenic signals only detected by the OWS or RPC. The energy spectrum of these veto-tagged signals was consistent with the spectrum of IBD-like candidate signals above 12 MeV, and was used to estimate the rate and energy spectrum for the fast neutron background from 0.7 to 12 MeV. The systematic uncertainty was estimated from the difference between this new analysis and the extrapolation method previously employed, and was determined to be half of the estimate reported in Ref. [?].

The methods used in Refs. [?, ?] to estimate the backgrounds from the uncorrelated prompt-delayed pairs (i.e. accidentals), the correlated β -n decays from cosmogenic ${}^9\text{Li}$ and ${}^8\text{He}$, and the ${}^{13}\text{C}(\text{n}){}^{16}\text{O}$ reaction, were extended to the current 6+8 AD data sample. The decrease in the single-neutron rate from the Am-C sources reduced the average rate of accidentals in the far hall by a factor of 2.7. As a result, the total backgrounds amount to about 3% (2%) of the IBD candidate sample in the far (near) hall(s). The systematic uncertainties in the ${}^{13}\text{C}(\text{n}){}^{16}\text{O}$ cross section and in the transportation of the β particles were reassessed through a comparison of experimental results and simulation packages, respectively [?]. The estimation of ${}^9\text{Li}/{}^8\text{He}$ now dominated the background uncertainty in both the near and far halls.

The estimated signal and background rates, as well as the efficiencies of the muon veto, ϵ_μ , and multiplicity selection, ϵ_m , are summarized in Table I. A detailed treatment of the absolute and relative efficiencies using the first six ADs was reported in Refs. [?, ?]. The uncertainties of the absolute efficiencies are correlated among the ADs and thus play a negligible role in the relative measurement of $\bar{\nu}_e$ disappearance. The performance of the two new ADs was found to be consistent with the other detectors. Estimates of two prominent uncorrelated uncertainties, the delayed-energy selection efficiency and the fraction of neutrons captured on Gd, were confirmed for all eight ADs using improved energy reconstruction and increased statistics.

Oscillation Analysis

Oscillation was measured using the L/E-dependent disappearance of $\bar{\nu}_e$, as given by the survival probability

$$P = 1 - \cos^4 \theta_{13} \sin^2 2\theta_{12} \sin^2 \left(\frac{1.267 \Delta m_{21}^2 L}{E} \right) - \sin^2 2\theta_{13} \sin^2 \left(\frac{1.267 \Delta m_{ee}^2 L}{E} \right)$$

where E is the energy in MeV of the $\bar{\nu}_e$, L is the distance in meters from its production point, θ_{12} is the solar mixing angle, and Δm_{21}^2 is the mass-squared difference of the first two neutrino mass eigenstates in eV^2 .

Recent precise measurements of the IBD positron energy spectrum disagree with models of reactor $\bar{\nu}_e$ emission [?, ?, ?]. The characteristics of the signals in this energy range are consistent with reactor antineutrino emission, and disfavor background or detector response as possible origins for the discrepancy. A separate manuscript, in preparation, will present the evidence in detail and provide the necessary data to allow detailed comparison of our measurement with existing and future models.

Given these discrepancies between measurements and models, here we present a technique for predicting the signal in the far hall based on measurements obtained in the near halls, with minimal dependence on models of the reactor antineutrinos. In our previous measurements [?], model-dependence was limited by allowing variation of the predicted $\bar{\nu}_e$ flux within model uncertainties, while the technique here provides an explicit demonstration of the negligible model dependence.

A χ^2 was defined as

$$\chi^2 = \sum_{i,j} (N_i^f - w_i \cdot N_j^n) (V^{-1})_{ij} (N_j^f - w_j \cdot N_j^n)$$

where N_i is the observed number of events after background subtraction in the i -th bin of reconstructed positron energy E_{rec} . The superscript f (n) denotes a far (near) detector. The symbol V represents a covariance matrix that includes known systematic and statistical uncertainties. The quantity w_i is a weight that accounts for the differences between near and far measurements.

For the case of a single reactor, the weight w_i can be simply calculated from the ratios of detector mass, distance to the reactor, efficiency, and antineutrino oscillation probability, as given by the relation:

$$w_i = \left(\frac{T_f}{T_n} \right) \left(\frac{\epsilon_f}{\epsilon_n} \right) \left(\frac{L_n}{L_f} \right)^2 \left(\frac{P_f}{P_n} \right) \left(\frac{\phi_f}{\phi_n} \right)$$

where T is the number of target protons, ϵ is the efficiency, and L is the distance to the reactor for a given detector. P_i is the oscillation probability for the i -th reconstructed energy bin and ϕ is the reactor antineutrino flux (which cancels from w_i). With P_i calculated in reconstructed positron energy, the detector response introduces small ($<0.2\%$ above 2 MeV) calculable deviations from Eq. 1.

For multiple reactor cores, the weight w_i was modified:

$$w_i = \left(\frac{T_f}{T_n}\right) \left(\frac{\epsilon_f}{\epsilon_n}\right) \sum_j P(E_{\text{true}}|E_{\text{rec}})_j r_j$$

The probability distribution $P(E_{\text{true}}|E_{\text{rec}})$ accounts for the energy transfer from the $\bar{\nu}_e$ to the e^+ and imperfections in the detector energy response (loss in non-active elements, non-linearity, and resolution). The extrapolation factor r_j was calculated as

$$r_j = \frac{\sum_{\text{cores}} P(E_{\text{true},j}, L_f) \phi_{jk} / L_f^2}{\sum_{\text{cores}} P(E_{\text{true},j}, L_n) \phi_{jk} / L_n^2}$$

where P is given by Eq. 1, L_f (L_n) is the distance between a far (near) detector and core k , and ϕ_{jk} is the predicted antineutrino flux from core k for the j -th true energy bin. In the single-reactor core case, the antineutrino flux ϕ cancels in the expression for r_j and Eq. 4 reduces to Eq. 3. Although the cancellation is not exact for multiple cores, the impact of the uncertainty in reactor antineutrino flux was found to be 0.1%.

The covariance matrix element V_{ij} was the sum of a statistical term, calculated analytically, and a systematic term determined by Monte Carlo calculation using

$$V_{ij} = \frac{1}{N} \sum_{k=1}^N (S_{f,i}^k - w_i \cdot S_{n,i}^k) (S_{f,j}^k - w_j \cdot S_{n,j}^k)$$

where N is the number of simulated experiments generated with energy spectra S , including systematic variations of detector response, $\bar{\nu}_e$ flux, and background. The choice of reactor antineutrino model [?, ?, ?, ?, ?] in calculating the covariance had negligible ($<0.2\%$) impact on the determination of the oscillation parameters.

Without loss of sensitivity, we summed the IBD signal candidates of the ADs within the same hall, accounting for small differences of target mass, detection efficiency, background and baseline. We considered the 6-AD and 8-AD periods separately in order to properly handle correlations in reactor antineutrino flux, detector exposure, and background. This means that i and j in the above

equations ran over the 37 reconstructed energy bins for the two near/far combinations and for the two periods considered ($37 \times 2 \times 2 = 148$). More details of this method are described in Ref. [?].

Using this method, we found $\sin^2 2\theta_{13} = 0.084 \pm 0.005$ and $|\Delta m_{ee}^2| = (2.42 \pm 0.11) \times 10^{-3} \text{ eV}^2$, with $\chi^2/\text{NDF} = 134.6/146$ (see the Supplemental Material [?]). While we use $\sin^2 2\theta_{12} = 0.857 \pm 0.024$ and $\Delta m_{21}^2 = (7.50 \pm 0.20) \times 10^{-5} \text{ eV}^2$ from Ref. [?], our result was largely independent of these values. Consistent results were obtained when our previous methods [?, ?] were applied to this larger dataset. Under the normal (inverted) hierarchy assumption, $|\Delta m_{32}^2| = (2.47 \pm 0.11) \times 10^{-3} \text{ eV}^2$ ($|\Delta m_{32}^2| = (2.37 \pm 0.11) \times 10^{-3} \text{ eV}^2$). This result was consistent with and of comparable precision to measurements obtained from accelerator ν_μ and $\bar{\nu}_\mu$ disappearance [?, ?]. Using only the relative rates between the detectors and $|\Delta m_{32}^2|$ from Ref. [?] we found $\sin^2 2\theta_{13} = 0.085 \pm 0.006$, with $\chi^2/\text{NDF} = 1.37/3$.

The reconstructed positron energy spectrum observed in the far site is compared in Fig. 3 [Figure 3: see original paper] with the expectation based on the near-site measurements. The 68.3%, 95.5% and 99.7% C.L. allowed regions in the $|\Delta m_{ee}^2| - \sin^2 2\theta_{13}$ plane are shown in Fig. 4 [Figure 4: see original paper]. The spectral shape from all experimental halls is compared in Fig. 5 [Figure 5: see original paper] to the electron antineutrino survival probability assuming our best estimates of the oscillation parameters.

The total uncertainties of both $\sin^2 2\theta_{13}$ and $|\Delta m_{ee}^2|$ are dominated by statistics. The most significant systematic uncertainties for $\sin^2 2\theta_{13}$ are due to the relative detector efficiency, reactor power, relative energy scale and ${}^9\text{Li}/{}^8\text{He}$ background. The systematic uncertainty in $|\Delta m_{ee}^2|$ is dominated by uncertainty in the relative energy scale.

Summary

Enhanced measurements of $\sin^2 2\theta_{13}$ and $|\Delta m_{ee}^2|$ have been obtained by studying the energy-dependent disappearance of the electron antineutrino interactions recorded in a 6.9×10^5 GWth-ton-days exposure. Improvements in calibration, background estimation, as well as increased statistics allow this study to provide the most precise estimates to date of the neutrino mass and mixing parameters $|\Delta m_{ee}^2|$ and $\sin^2 2\theta_{13}$.

Acknowledgments

Daya Bay is supported in part by the Ministry of Science and Technology of China, the U.S. Department of Energy, the Chinese Academy of Sciences, the CAS Center for Excellence in Particle Physics, the National Natural Science Foundation of China, the Guangdong provincial government, the Shenzhen municipal government, the China General Nuclear Power Group, Key Laboratory of Particle and Radiation Imaging (Tsinghua University), the Ministry of Educa-

tion, Key Laboratory of Particle Physics and Particle Irradiation (Shandong University), the Ministry of Education, Shanghai Laboratory for Particle Physics and Cosmology, the Research Grants Council of the Hong Kong Special Administrative Region of China, the University Development Fund of The University of Hong Kong, the MOE program for Research of Excellence at National Taiwan University, National Chiao-Tung University, and NSC fund support from Taiwan, Alfred P. Sloan Foundation, the Ministry of Education, Youth, and Sports of the Czech Republic, the Joint Institute of Nuclear Research in Dubna, Russia, the NSFC-RFBR joint research program, the National Commission of Scientific and Technological Research of Chile, and the Tsinghua University Initiative Scientific Research Program. We acknowledge Yellow River Engineering Consulting Co., Ltd., and China Railway 15th Bureau Group Co., Ltd., for building the underground laboratory. We are grateful for the ongoing cooperation from the China General Nuclear Power Group and China Light and Power Company.

Appendix: Why Δm_{ee}^2 is used by Daya Bay

This section describes the advantages of reporting the Daya Bay measurement of electron antineutrino disappearance in terms of an effective mass-squared difference Δm_{ee}^2 , which is independent of the unknown ordering of neutrino masses and future improvements in our knowledge of the solar oscillation parameters.

In the three-flavor framework, the survival probability of electron antineutrino is given by

$$P(\bar{\nu}_e \rightarrow \bar{\nu}_e) = 1 - \cos^4 \theta_{13} \sin^2 2\theta_{12} \sin^2 \Delta_{21} - \sin^2 2\theta_{13} (\cos^2 \theta_{12} \sin^2 \Delta_{31} + \sin^2 \theta_{12} \sin^2 \Delta_{32}) \quad (7)$$

where $\Delta_{ji} \equiv 1.267 \Delta m_{ji}^2 L/E$ and Δm_{ji}^2 is the mass-squared difference between eigenstates ν_j and ν_i in eV^2 . The three mass-squared differences are subject to the constraint $|\Delta m_{31}^2| = |\Delta m_{32}^2| \pm |\Delta m_{21}^2|$, where “+” (“-”) is for the normal (inverted) mass ordering. Therefore, determination of $|\Delta m_{32}^2|$ (or $|\Delta m_{31}^2|$) depends on knowledge of the mass ordering and solar oscillation parameters.

The Daya Bay experiment reports a precise measurement of the effective mass splitting Δm_{ee}^2 , which is independent of our knowledge of the ordering and solar parameters. In this approach, we approximate the survival probability using

$$P(\bar{\nu}_e \rightarrow \bar{\nu}_e) \approx 1 - \cos^4 \theta_{13} \sin^2 2\theta_{12} \sin^2 \Delta_{21} - \sin^2 2\theta_{13} \sin^2 \Delta_{ee} \quad (8)$$

Despite the advantage of using Δm_{ee}^2 for the measurement, it has the disadvantage of not being a fundamental parameter. Therefore, we must determine a relation between Δm_{ee}^2 and Δm_{32}^2 given knowledge of the mass ordering and solar oscillation parameters.

Mathematical Derivation

Using the relation $|\Delta m_{31}^2| = |\Delta m_{32}^2| \pm |\Delta m_{21}^2|$, Eq. 7 can be written as

$$P(\bar{\nu}_e \rightarrow \bar{\nu}_e) = 1 - 2s_{12}^2 \sin^2 \Delta_{21} - 4c_{12}^4 s_{13}^2 - 4s_{12}^2 c_{12}^2 s_{13}^2 \cos(2\Delta_{32} \pm \phi) \quad (9)$$

where $c_x = \cos \theta_x$, $s_x = \sin \theta_x$, and $\phi = \arctan\left(\frac{\sin 2\Delta_{21} \cos 2\Delta_{21} + \tan^2 \theta_{12}}{\sin^2 2\Delta_{21}}\right)$. The last term of the above formula is the so-called ‘‘solar term’’ that governs the reactor antineutrino oscillation at O(100) km. For the L/E range covered by Daya Bay, $4s_{12}^2 \sin^2 \Delta_{21} \ll 1$. Thus, Eq. 9 can be approximated as

$$P(\bar{\nu}_e \rightarrow \bar{\nu}_e) \approx 1 - 4s_{13}^2 [1 - \cos(2\Delta_{32} \pm \phi)] = 1 - \sin^2 2\theta_{13} \sin^2(\Delta_{32} \pm \phi/2) \quad (10)$$

By comparing Eq. 10 with Eq. 8, we obtain the expression relating Δm_{ee}^2 to Δm_{32}^2 (or Δm_{31}^2):

$$|\Delta m_{ee}^2| = |\Delta m_{32}^2| \pm \Delta m_{21}^2 \cos \phi \mp |\Delta m_{21}^2| \sin \phi \tan \phi/2 \quad (11)$$

where $\Delta m_{ee}^2 = \phi \times \frac{4E}{1.267L}$.

Numerical Evaluation

By definition, Δm_{ee}^2 is a function of L/E. Using the current values of $\Delta m_{21}^2 = 7.50 \times 10^{-5} \text{ eV}^2$ and $\sin^2 2\theta_{12} = 0.857$ [?], Fig. 6 [Figure 6: see original paper] shows the value of $\Delta m_{ee}^2/2$ as a function of energy for $L = 1.6 \text{ km}$. We find that $\Delta m_{ee}^2/2 \approx 5.17 \times 10^{-5} \text{ eV}^2$ is essentially a constant in our L/E region, and numerically identical to $\cos^2 \theta_{12} \Delta m_{21}^2$. Thus, this definition of Δm_{ee}^2 is similar to the definition introduced in Ref. [?]:

$$|\Delta m_{ee}^2| = \cos^2 \theta_{12} |\Delta m_{31}^2| + \sin^2 \theta_{12} |\Delta m_{32}^2| \approx |\Delta m_{32}^2| + \cos^2 \theta_{12} \Delta m_{21}^2 \quad (12)$$

Figure 7 [Figure 7: see original paper] compares the approximated formula with $\phi/2 = 5.17 \times 10^{-5} \text{ eV}^2$,

$$P_{ee} \approx 1 - \sin^2 2\theta_{13} \sin^2 (\Delta_{32} + 5.17 \times 10^{-5} \text{ eV}^2) \quad (13)$$

to the three-flavor formula, Eq. 7. In this comparison, $\Delta m_{32}^2 = 2.44 \times 10^{-3} \text{ eV}^2$, $L = 1.6 \text{ km}$, $\sin^2 2\theta_{13} = 0.09$, and normal mass hierarchy are the inputs. The agreement between the two, better than 10^{-4} , is excellent and exceeds the achievable experimental precision.

This study demonstrates that, once we obtain the value of $|\Delta m_{ee}^2|$ using Eq. 8, we can reliably deduce the values of $|\Delta m_{32}^2|$ and $|\Delta m_{31}^2|$ using Eqs. 11 and 12 with $\phi/2 \approx \cos^2 \theta_{12} \Delta m_{21}^2$. Using the current values of θ_{12} and Δm_{21}^2 , $\Delta m_{\phi}^2/2 \approx 5.17 \times 10^{-5} \text{ eV}^2$, and $(|\Delta m_{31}^2| - |\Delta m_{32}^2|) \approx 2.33 \times 10^{-5} \text{ eV}^2$.

It is important to point out that the exact solution of $\sin^2(\Delta_{31}) = \cos^2 \theta_{12} \sin^2(\Delta_{32} + \Delta_{21}) + \sin^2 \theta_{12} \sin^2(\Delta_{32})$ was never used to extract the value of Δm_{ee}^2 in Daya Bay.

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv – Machine translation. Verify with original.