

A holographic study on vector condensate induced by a magnetic field Postprint

Authors: Cai, Rong-Gen, He, Song, Li, Li, Li, Li-Fang

Date: 2016-12-22T00:00:00+00:00

Abstract

We study a holographic model with vector condensate by coupling the anti-de Sitter gravity to an Abelian gauge field and a charged vector field in $(3 + 1)$ dimensional spacetime. In this model there exists a non-minimal coupling of the vector field to the gauge field. We find that there is a critical temperature below which the charged vector condenses via a second order phase transition. The DC conductivity becomes infinite and the AC conductivity develops a gap in the condensed phase. We study the effect of a background magnetic field on the system. It is found that the background magnetic field can induce the condensate of the vector field even in the case without chemical potential/charge density. In the case with non-vanishing charge density, the transition temperature raises with the applied magnetic field, and the condensate of the charged vector operator forms a vortex lattice structure in the spatial directions perpendicular to the magnetic field.

Full Text

Preamble

A Holographic Study on Vector Condensate Induced by a Magnetic Field

Rong-Gen Cai^{1*}, Song He^{1†}, Li Li^{1‡}, Li-Fang Li²

¹State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China.

²State Key Laboratory of Space Weather, Center for Space Science and Applied Research, Chinese Academy of Sciences, Beijing 100190, China.

December 3, 2013

Abstract

We study a holographic model with vector condensate by coupling the anti-de Sitter gravity to an Abelian gauge field and a charged vector field in $(3 + 1)$ dimensional spacetime. In this model there exists a non-minimal coupling of the vector field to the gauge field. We find that there is a critical temperature below which the charged vector condenses via a second order phase transition. The DC conductivity becomes infinite and the AC conductivity develops a gap in the condensed phase. We study the effect of a background magnetic field on the system. It is found that the background magnetic field can induce the condensate of the vector field even in the case without chemical potential/charge density. In the case with non-vanishing charge density, the transition temperature raises with the applied magnetic field, and the condensate of the charged vector operator forms a vortex lattice structure in the spatial directions perpendicular to the magnetic field.

Introduction

The gauge/gravity duality [1, 2, 3] has proven to be a useful and complementary framework for studying strongly coupled systems through an appropriate gravity theory living in a higher dimensional spacetime. There are two complementary approaches. In the top-down approach, one can obtain low energy effective theories in the bulk as consistent truncations of string/M theory by dimensional reduction on compactified manifolds. The advantage of this approach is that one knows the origin of the effective gravity theory in the bulk and the details of the dual field theory. One well-known example in this category is the duality between IIB superstring theory on $\text{AdS}_5 \times S^5$ and $N = 4$ supersymmetric Yang-Mills theory on the AdS_5 boundary. On the other hand, in the so-called bottom-up approach, the gravity theory in the bulk is usually constructed based on physical considerations according to the AdS/CFT dictionary. The models constructed in this way are simple and universal, much like the Landau-Ginzburg theory describing superconductivity. The disadvantage is that the details of the dual field theory are not very clear.

A well-known example in the second category is the holographic superconductor model [4, 5]. More specifically, to build a holographic superconductor model, one should first introduce a $U(1)$ gauge field in the bulk, which corresponds to a global $U(1)$ symmetry on the boundary side. To break this $U(1)$ symmetry spontaneously, one needs a charged operator condensing at low temperature. Therefore, one includes a charged scalar field in the bulk which is dual to the boundary scalar operator. For simplicity, the gauge field and the charged scalar can be minimally coupled. This holographic toy model admits black holes with scalar hair at low temperatures (superconducting phase), but without scalar hair at high temperatures (normal phase). In this way, one mimics the superconductor/conductor phase transition on the field theory side. Holographic superconductor models constructed in the top-down approach can also be found, for example, in refs. [6, 7, 8, 9].

It is well known that superconductivity involves the formation of a quantum condensate state by pairing conduction electrons. The pair of electrons, called the Cooper pair, can be in a state of either spin singlet with total spin $s = 0$ or spin triplet with $s = 1$. The order parameter in a superconductor is expressed in terms of the gap function. Due to the anti-commuting properties of the electron wave function, the antisymmetric spin-singlet state is associated with a symmetric orbital wave function with orbital angular momentum $l = 0$ (s-wave), 2 (d-wave), etc., while the symmetric spin-triplet state is accompanied by an antisymmetric orbital wave function with orbital angular momentum $l = 1$ (p-wave), 3 (f-wave), etc. Since the condensed field in the holographic setup [4, 5] is a scalar field dual to a scalar operator in the field theory side, it is therefore a holographic s-wave model.

In the holographic setup, a d-wave order parameter is dual to a charged spin-two field propagating in the bulk [10, 11]. To mimic p-wave superconductivity, one can consider a vector order parameter which is dual to a vector field on the gravity side. A holographic p-wave model [12] was constructed by adding an $SU(2)$ Yang-Mills field into the bulk. A $U(1)$ subgroup of the Yang-Mills field is regarded as the gauge group of electromagnetism. A gauge boson generated by another $SU(2)$ generator, charged under this $U(1)$ by the nonlinear coupling of the non-Abelian field, is dual to the vector order parameter. An alternative holographic realization of p-wave superconductivity emerges from the condensation of a 2-form field in the bulk [13]. A holographic chiral $p + ip$ superconductor was discussed by generalizing the $SU(2)$ model, introducing a Maxwell field and a Chern-Simons term [14]. Note that in principle we can also build a holographic p-wave model by introducing a complex vector field charged under a $U(1)$ gauge field in the bulk with possible couplings.

On the other hand, motivated by the possibility of creating very strong magnetic fields, for example at RHIC and LHC, interest in investigating the properties of QCD matter in strong magnetic fields has been growing recently. Some new phenomena have been revealed, such as the chiral magnetic effect and a split between the deconfinement phase transition and chiral symmetry restoration phase transition; see ref. [15] for a review. An interesting new phenomenon is the possibility that the QCD vacuum undergoes a phase transition to a new phase with charged ρ -meson condensation in a sufficiently strong magnetic field [16, 17]. This exotic phase is a kind of anisotropic superconducting phase [18]. The author in ref. [16] adopted an effective quantum electrodynamics action (DSGS model [19]) to discuss the condensate of ρ -meson. A similar study based on the Nambu-Jona-Lasinio model [20] can be found in ref. [17]. The uniform magnetic field background is encoded in an Abelian gauge field. As a vector boson, the possible condensate of the ρ -meson in a uniform magnetic field can be realized in the holographic framework by introducing a charged vector field on the gravity side in the presence of a magnetic field. Based on the Sakai-Sugimoto model, the holographic ρ -meson was studied in the confinement phase at zero temperature [21], where it was shown that the effective mass of the ρ -meson at strong magnetic field becomes tachyonic. The influence of the magnetic field

on the chiral transition temperature and deconfinement transition temperature was discussed in ref. [22].

In the SU(2) model [12], a similar instability triggered by a non-Abelian magnetic field has been found in refs. [23, 24, 25], which is reminiscent of the observation that a non-Abelian magnetic field induces W-boson condensate exhibiting vortex lattices in flat spacetime [26, 27, 28].

In this paper we construct a toy model by introducing a complex vector field charged under an Abelian gauge field in the bulk, which is dual to a strongly coupled system involving a charged vector operator with a global U(1) symmetry. In this model there exists a non-minimal coupling between the vector field and the Abelian gauge field. In this bottom-up approach, we do not know the details of the dual field theory clearly. However, our setup meets the minimal requirement to construct a holographic p-wave superconductor model. This model also has potential to discuss the ρ -meson condensate in the presence of a magnetic field.

We first study the model in the case without a background magnetic field, i.e., turning on only the temporal component of the gauge field, which is the usual way of analyzing holographic superconductors at a first step [4, 10, 11, 12]. We find a critical temperature below which a vector operator acquires a vacuum expectation value, breaking the Abelian symmetry spontaneously. Furthermore, the condensate of this vector operator picks out a special spatial direction, thus the rotational symmetry is also broken in the condensed phase. The system undergoes a second-order phase transition with the critical exponent one-half, which coincides with the result from mean field theory. This condensed phase presents characteristics known from superconductivity, such as infinite DC conductivity and a gap in the optical conductivity.

An instability induced by a background non-Abelian magnetic field has been reported, where the non-Abelian current operators obtain vacuum expectation values resulting in a vortex lattice structure [24, 25]. We are interested in how the applied Abelian magnetic field influences the instability of our model with Abelian gauge symmetry. For this, we turn on a uniform magnetic field B in the bulk, which immerses the condensed phase into an external magnetic field.

We find that the increase of the magnetic field induces instability of the black hole background, which gives rise to a family of condensates induced by the applied magnetic field in the dual strongly coupled system. This magnetic field-induced instability can happen even in the case with vanishing chemical potential or charge density. For finite magnetic field, there is a tower of “droplet” solutions [29] in the sense that they are localized in a finite region. Further, the condensate shrinks in size as one increases the magnetic field, which shares similarity with the Meissner effect. But it is not the exact case with conventional superconductivity, where superconductivity is suppressed by magnetic field in real materials. Of course, it is interesting to notice the fact that it has been reported recently that an applied magnetic field may induce superconductivity

[30, 31]. In addition, the emergence of charged vector operator condensate triggered by the applied magnetic field is consistent with the appearance of electromagnetically superconducting phase in a strong magnetic field studied from both field theory methods [16, 17] and holographic setups [21, 22]. We also manage to construct the vortex lattice solution in the condensed phase near the phase transition, which is very reminiscent of the Abrikosov lattice in common type-II superconductors.

The plan of this paper is as follows. In section 2, we introduce the holographic model and deduce the general equations of motion of the system. In section 3, we turn on a uniform magnetic field and give details about how to recover the Landau levels in this holographic model. Section 4 is devoted to discussing the phase diagram with/without chemical potential. The vortex lattice solutions are constructed in section 5. The conclusion and further discussions are included in section 6. We study, in Appendix A, the condensate of the vector field and the phase transition in the case without magnetic field by calculating the conductivity to ensure the condensed phase is a superconducting state.

2 The Holographic Model

Let us introduce a charged vector field into the (3+1) dimensional Einstein-Maxwell theory with a negative cosmological constant. The full action reads

$$S = \frac{1}{2\kappa^2} \int d^4x \sqrt{-g} \left[R + \frac{6}{L^2} - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{1}{2} \rho_{\mu\nu}^\dagger \rho^{\mu\nu} - m^2 \rho_\mu^\dagger \rho^\mu + iq\gamma \rho_\mu \rho_\nu^\dagger F^{\mu\nu} \right],$$

where the covariant derivative $D_\mu = \nabla_\mu - iqA_\mu$ with q the charge of vector field ρ_μ . The strength of the U(1) field A_μ is $F_{\mu\nu} = \nabla_\mu A_\nu - \nabla_\nu A_\mu$. The tensor $\rho_{\mu\nu}$ in (1) is defined by $\rho_{\mu\nu} = D_\mu \rho_\nu - D_\nu \rho_\mu$. The last interacting term describes the non-minimal coupling of the charged vector field ρ_μ to the U(1) gauge field A_μ . The parameter γ characterizes the magnetic moment of the vector field ρ_μ and is assumed to be non-negative. The form of the action is reminiscent of the DSGS model which describes the quantum electrodynamics of the ρ -meson proposed by Djukanovic, Schindler, Gegelia, and Scherer in ref. [19]. Compared to the action of the DSGS model, it is easy to find that the action in (1) is just a simple generalization of the DSGS model to anti-de Sitter space. Note that the part of neutral ρ -meson in ref. [19] is neglected here since it is not relevant to our goal in this paper.

Varying the action (1) with respect to A_μ yields the equation of motion for the gauge field

$$\nabla_\nu F^{\nu\mu} = iq(\rho_\nu \rho^{\dagger\nu\mu} - \rho_\nu^\dagger \rho^{\nu\mu}) + iq\gamma \nabla_\nu (\rho^\nu \rho_\mu^\dagger - \rho^{\dagger\nu} \rho_\mu),$$

while a variation of the action (1) with respect to ρ_ν^\dagger gives the equation of motion for the charged vector field

$$D_\nu \rho^{\nu\mu} - m^2 \rho^\mu + iq\gamma \rho_\nu F^{\nu\mu} = 0.$$

If one takes the limit $q \rightarrow \infty$ while keeping $q\rho_\mu$ and qA_μ fixed, the back reaction of the matter sources to the background can be ignored. This is the probe limit we will adopt in this paper. The background is taken to be a (3 + 1) dimensional Schwarzschild-AdS black hole

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2(dx^2 + dy^2),$$

with $f(r) = r^2(1 - r_h^3/r^3)$ and r_h the horizon radius. The Hawking temperature for this black hole is $T = f'(r_h)/4\pi = 3r_h/4\pi$, which sets the temperature of the boundary field theory.

In the probe limit, the matter fields ρ_μ and A_μ can be treated as perturbations on the Schwarzschild-AdS black hole background.

3 Adding a Constant Magnetic Field

In Appendix A, we discussed the condensate of the vector field ρ_μ induced by the bulk electric field by turning on only the scalar potential A_t associated with the Maxwell field. The normal phase corresponds to the black hole solution with a vanishing vector field ρ_μ . As one lowers the temperature, the normal phase becomes unstable to developing a non-trivial configuration of the vector field ρ_μ . This gives a non-zero vacuum expectation value of the dual vector operator, which breaks the U(1) gauge symmetry and the rotational symmetry in the x - y plane. The calculation of the optical conductivity reveals that there is a delta function at the origin for the real part of the conductivity, which means the condensed phase is indeed superconducting. For details, see Appendix A.

We now turn on a magnetic field to study how the applied magnetic field influences the system. A consistent ansatz is as follows

$$\rho_\nu dx^\nu = [\epsilon \rho_x(r, x) e^{ipy} + \mathcal{O}(\epsilon^3)] dx + [\epsilon \rho_y(r, x) e^{ipy} e^{i\theta} + \mathcal{O}(\epsilon^2)] dy,$$

$$A_\nu dx^\nu = [\phi(r) + \mathcal{O}(\epsilon^2)] dt + [Bx + \mathcal{O}(\epsilon^2)] dy,$$

where $\rho_x(r, x)$, $\rho_y(r, x)$ and $\phi(r)$ are all real functions, p is a real constant, the constant θ is the phase of ρ_y , and $B > 0$ is the constant magnetic field perpendicular to the x - y plane. We have defined the deviation parameter ϵ from the critical point at which the condensate begins to appear.

The zeroth order of (2) gives the equation of motion for ϕ :

$$\phi''(r) + \frac{f'(r)}{f(r)}\phi'(r) = 0.$$

The asymptotic value of ϕ gives the chemical potential $\mu = A_t(r \rightarrow \infty)$ of the dual field theory. The boundary condition at the horizon is given by requiring that $A_\mu A^\mu$ is finite there. Thus we can obtain a unique solution $\phi(r) = \mu(1 - r_h/r)$.

The equations of motion for ρ_x and ρ_y can be deduced from (3) at order $\mathcal{O}(\epsilon)$. We further separate the variables as $\rho_x(r, x) = \phi_x(r)X(x)$ and $\rho_y(r, x) = \phi_y(r)Y(x)$. We find that to satisfy the equations of motion of the model with the given ansatz, θ can only be chosen as $\theta_+ = \pi/2 + 2n\pi$ or $\theta_- = -\pi/2 + 2n\pi$ with n an arbitrary integer. The equations of motion for $\phi_x(r)$, $\phi_y(r)$, $X(x)$ and $Y(x)$ are divided into the following equations:

$$\phi'_x X + \phi'_y Y + \frac{2f}{r}\phi_x X + \frac{2f}{r}\phi_y Y = 0,$$

$$\phi''_x X + \phi''_y Y + \left(\frac{f'}{f} + \frac{2}{r}\right)(\phi'_x X + \phi'_y Y) + \frac{1}{f} \left[\frac{\omega^2}{f} - m^2 - \frac{(qBx - p)^2}{r^2} \right] (\phi_x X + \phi_y Y) = 0,$$

$$\ddot{X} + \left[E - \frac{(qBx - p)^2}{r^2} \right] X = 0,$$

$$\ddot{Y} + \left[E - \frac{(qBx - p)^2}{r^2} \mp c(1 + \gamma)qB \right] Y = 0,$$

where the prime denotes the derivative with respect to r and the dot denotes the derivative with respect to x . Here and below the upper signs correspond to the θ_+ case and the lower to the θ_- case. In order to satisfy (8), one should impose

$$\phi_y = c\phi_x, \quad c(qBx - p)Y = 0,$$

where c is a real constant. This constraint is automatically satisfied by (9). We can see that only two of the four functions are independent. Substituting (12) into the remaining equations, we can find the following three equations:

$$\phi''_x + \left(\frac{f'}{f} + \frac{2}{r}\right)\phi'_x + \frac{1}{f} \left[\frac{\omega^2}{f} - m^2 - E \right] \phi_x = 0,$$

$$c(1 + \gamma)qBY + (qBx - p)^{2Y} = EY,$$

$$(1 + \gamma)qBX + (qBx - p)^{2X} = EX.$$

One can get the value of E for arbitrary constant c by solving the eigenvalue problem (14) and (15) with the constraint given in (12). There may exist the possibility that one cannot obtain non-trivial solutions for some special values of c . Here we consider a simple case with $c^2 = 1$, in which the equations of motion for $X(x)$ and $Y(x)$ can be solved exactly. Since $c = 1/c$, the c in the denominator in (15) is equivalent to the case in the numerator. Subtracting (14) from (15) and defining a new function as $\psi(x) = X(x) - Y(x)$, one gets the equation

$$\ddot{\psi}(x) + [E - (qBx - p)^2 \mp c(1 + \gamma)qB] \psi(x) = 0.$$

We further introduce a new variable $\xi = \sqrt{|qB|}(x - p/qB)$, then the above equation becomes

$$\frac{d^2}{d\xi^2} \psi(\xi) + (\eta - \xi^2) \psi(\xi) = 0.$$

The regular and bounded solution of (18) is given by Hermite function H_n as

$$\psi_n(\xi) = N_n e^{-\xi^2/2} H_n(\xi),$$

with the corresponding eigenvalue $\eta_n = 2n + 1$. N_n is a normalization constant and n is a non-negative integer. Thus we obtain the solution of equation (17) as

$$\psi_n(x) = N_n e^{-|qB|(x-p/qB)^2},$$

with the corresponding eigenvalue

$$E_n = (2n + 1)|qB| \mp c(1 + \gamma)qB.$$

Combining (12), (16) and (20), one can obtain the exact configurations for $X(x)$ and $Y(x)$, which read

$$X_n(x; p) = e^{\mp \frac{1}{2}|qB|(x-p/qB)^2} \left[X(0) e^{\pm cqB \int_0^x (x' - p/qB) dx'} \mp \frac{cqB}{|qB| \mp cqB} N_n \times \int_0^x e^{\pm |qB|(x' - p/qB)^2} dx' \right],$$

$$Y_n(x; p) = X_n(x; p) - N_n e^{-\frac{1}{2}|qB|(x-p/qB)^2},$$

where $X(0)$ is a constant denoting the value of $X_n(x; p)$ at the origin $x = 0$. The solutions of ϕ_x and ϕ_y corresponding to the eigenvalue E_n , denoted by ϕ_{xn} and ϕ_{yn} , can be obtained by solving the equation of motion (13) with E_n given in (21). So far, we have recovered the Landau levels. As one can see in Appendix A, we have a second-order phase transition from the normal phase with $\rho_\mu = 0$ to the condensed phase with $\rho_\mu \neq 0$. Therefore we should encounter a marginally stable mode at the transition point. These solutions obtained from (13) just correspond to the marginally stable states.

We can see from (13) that the effective mass of ρ_x is

$$m_{\text{eff}}^2 = m^2 + (2n + 1)|qB| \mp c(1 + \gamma)qB,$$

which is clearly shifted by the magnetic field B . Depending on concrete parameters and Landau level, the appearance of magnetic field can increase or decrease the effective mass, thus will hinder or enhance the transition from the normal phase to the condensed phase.

In what follows, we consider the case with the lowest Landau level with $n = 0$, which reads

$$\psi_0(x; p) = N_0 e^{-|qB|(x-p/qB)^2}.$$

4 Phase Diagram

We are interested in how the applied magnetic field influences the transition temperature from the normal phase to the condensed phase. As we can see from (24), the effective mass of the charged vector field ρ_μ in the lowest energy state, i.e., in the lowest Landau level $n = 0$, depends on the magnetic field B and the non-minimal coupling parameter γ as

$$m_{\text{eff}}^2 = m^2 + |qB| \mp c(1 + \gamma)qB.$$

It is clear that the increase of the magnetic field B decreases the effective mass and thus tends to raise the transition temperature.

4.1 Phase Diagram at Vanishing Charge Density

It is well known that the increase of the electric field decreases the effective mass of charged scalar or vector fields, inducing the transition from the normal phase to the condensed phase. We first turn off the electric field, which corresponds to the case with vanishing charge density $\rho = 0$. We introduce a new coordinate $z = r_h/r$. The equation of motion (13) can be rewritten as

$$\phi_x''(z) - \frac{3}{z}\phi_x'(z) + \left[\frac{9}{z^4(1-z^3)} - \frac{m^2}{z^2(1-z^3)} - \frac{E_n}{r_h^2(1-z^3)} \right] \phi_x(z) = 0,$$

with $\zeta = |qB|/T^2$.

To solve such a second-order equation by the shooting method, we impose the regular condition at the horizon $z = 1$ as well as the source-free condition at the boundary $z = 0$. More specifically, we set $\phi_x(1) = 1$ in our numerical calculation due to the linearity of (27). For a given m^2 , only for certain values of $\zeta = |qB|/T^2$ can the boundary conditions be satisfied.

Figure 1 [Figure 1: see original paper] presents the three marginally stable curves of $\phi_x(z)$ for $m^2 = 3/4$. The three lowest-lying modes are in the sequence $\zeta_0 < \zeta_1 < \zeta_2$. The red line corresponding to the minimal value of ζ has no intersecting points with the horizontal axis at non-vanishing z . Such a mode with $\zeta_0 \simeq 72.84$ is considered as a mode of node $n = 0$. Furthermore, the green line to $\zeta_1 \simeq 339.24$ and blue line to $\zeta_2 \simeq 780.10$ are regarded as modes with nodes $n = 1$ and $n = 2$, respectively. Since the radial oscillations in the z -direction of $\phi_x(z)$ will cost more energy, the latter two curves are therefore thought to be unstable. Thus the lowest value ζ_0 just gives the critical magnetic field above which the normal state is unstable to developing vector hair. Figure 2 [Figure 2: see original paper] shows the critical magnetic field in terms of ζ_0 for various squared masses of the vector field. It can be seen clearly that ζ_0 increases as we increase the squared mass.

It should be pointed out that if we turn off the magnetic field, the normal state will not become unstable to developing hair with vanishing charge density. The interesting result here is that only the magnetic field itself can trigger the phase transition. This result has an analogy to the QCD vacuum instability induced by a strong magnetic field to spontaneously develop the ρ -meson condensate. It is clear that the last term in (1) describing a non-minimal coupling of the vector field ρ_μ to the gauge field A_μ plays a crucial role in the instability. Note that similar coupling can be found in many formalisms used to describe the coupling of magnetic moment to the background magnetic field for charged particles of spin 1, i.e., vector particles [19, 32].

4.2 Phase Diagram at Finite Charge Density

Let us now consider the system with fixed charge density ρ . Here we do not limit ourselves to the lowest energy state. The equation of motion (13) can be written in terms of ρ and T as

$$\phi_x''(z) - \frac{3}{z}\phi_x'(z) + \left[\frac{9}{z^4(1-z^3)} - \frac{m^2}{z^2(1-z^3)} - \frac{E_n}{\rho^{2/3}T^{4/3}} \frac{(1+z+z^2)^2}{z^4(1-z^3)} \right] \phi_x(z) = 0,$$

where $\lambda = \rho^{2/3}/T^{4/3}$ and E_n is given in (21). The lowest energy state corresponds to $E_n = E_{LLL}$. For numerical convenience and to match the behavior at the boundary, we further define $\phi_x(z) = (1-z)^\Delta F(z)$. Then we can obtain

$$F''(z) + \left[\frac{2\Delta}{z-1} - \frac{3}{z} \right] F'(z) + \left[\frac{\Delta(\Delta-1)}{(z-1)^2} + \frac{3\Delta}{z(z-1)} + \frac{9}{z^4(1-z^3)} - \frac{m^2}{z^2(1-z^3)} - \frac{E_n}{\rho^{2/3}T^{4/3}} \frac{(1+z+z^2)^2}{z^4(1-z^3)} \right] F(z) =$$

This equation depends on two physical parameters T and ρ . To solve such a second-order equation, we impose the regular condition at the horizon $z = 1$ as well as the source-free condition $F(0) = 0$ at the boundary $z = 0$. It has a non-trivial solution only when there is a relation between these two parameters, which just gives the transition temperature as a function of the magnetic field.

The (T, B) phase diagram with the lowest Landau level is drawn in figure 3 [Figure 3: see original paper]. To determine which side of the phase transition line is the condensed phase, we can consider equation (26). It suggests that the magnetic field decreases the effective mass. So if we increase the magnetic field at a fixed temperature, the normal state will become unstable for sufficiently large magnetic field. Figure 3 looks very similar to figure 9 in ref. [22] where the chiral transition temperature rises with magnetic field, indicating chiral magnetic catalysis.

Furthermore, to compare with the lowest Landau level case, we present an example with positive $E_n = |qB|$ in (28) in figure 4 [Figure 4: see original paper]. One can see clearly in this case that the transition temperature lowers with the increase of the applied magnetic field. This is the well-known property of ordinary superconductors, which was first discussed in a holographic setup in ref. [33].

5 Vortex Lattice Solution

Let us now construct the vortex lattice solution. It is enough to consider the $n = 0$ solution only, i.e., $\psi_0(x; p) = N_0 e^{-|qB|(x-p/qB)^2}$. Due to the fact that the eigenvalue E_n is independent of p , a linear superposition of the solutions $e^{ipy} \phi_{xn}(r) X_n(x; p)$ and $e^{ipy} \phi_{yn}(r) Y_n(x; p)$ with different p is also a solution of the model at $\mathcal{O}(\epsilon)$. We introduce two functions

$$\rho_x(r, x, y) = \phi_{x0}(r) \sum_{\ell} c_{\ell} e^{ip_{\ell}y} X_0^L(x; p_{\ell}),$$

$$\rho_y(r, x, y) = c e^{i\theta_{\pm}} \phi_{x0}(r) \sum_{\ell} c_{\ell} e^{ip_{\ell}y} Y_0^L(x; p_{\ell}),$$

which satisfy the full equations of motion. Following ref. [33], we can obtain the vortex lattice solution

$$\rho(r, x, y) \equiv \rho_x(r, x, y) + i e^{i\theta_{\pm}} \rho_y(r, x, y) = \phi_{x0}(r) \sum_{\ell} c_{\ell} e^{ip_{\ell}y} \psi_0(x; p_{\ell}).$$

$\rho(r, x, y)$ has a pseudo-periodicity

$$\rho(r, x + a_1, y + a_2) = e^{i\alpha_1 x + i\alpha_2 y} \rho(r, x, y),$$

as well as a zero at

$$x_{m,n} = (m + \frac{1}{2})b_1 + (n + \frac{1}{2})b_2,$$

with two vectors $b_1 = a_1 \partial_x$ and $b_2 = 2\pi \partial_y$. m and n are two integers.

Since the expectation value of the operator \hat{J}^μ dual to ρ_μ is given by the coefficient at boundary $r \rightarrow \infty$, the quantity $J_i \equiv \langle \hat{J}_i \rangle$ indeed exhibits the vortex structure with the cores of vortices located at $x_{m,n}$. In particular, the triangular lattice with three adjoining vortices forming an equilateral triangle can be obtained by choosing the following parameters:

$$a_1 = \sqrt{\frac{2\pi}{|qB|}}, \quad a_2 = \sqrt{\frac{2\pi}{|qB|}} e^{i\pi/3}.$$

It should be stressed that it is the special combinations J_\pm which exhibit the vortex lattice structure. In particular, for $q > 0$, it is J_- that corresponds to the lowest Landau level, while for $q < 0$, it is J_+ . The form of operator presenting vortex lattice structure is the same as the one in field theory studies without gravity [16]. This also provides evidence for the correctness of choosing $c^2 = 1$ in section 3.

Figure 5 [Figure 5: see original paper] shows the configuration of the norm of condensate J_i in the x - y plane for the triangular lattice. Obviously, to obtain the true ground state, we should calculate the free energy of the solutions with different lattice structures from the action to find which configuration minimizes the free energy.

It turns out that the linear analysis presented here is not sufficient to determine the most stable solution. We should include higher-order contributions just as done in refs. [24, 33]. The calculation is much more complicated and is not very relevant to the purpose of this paper. We leave it for further study.

6 Conclusion and Discussion

In this paper we studied a holographic model with a complex vector field charged under a U(1) gauge field in a (3 + 1) dimensional AdS black hole background, aiming to shed some light on real strongly coupled systems which are gravity duals. In this model, there is a non-minimal coupling of the vector field to the U(1) gauge field, which describes the interaction between the magnetic dipole moment of the vector field and the background magnetic field. This model

includes the minimal ingredients to build a holographic p-wave superconductor model. We found a critical temperature at which the system undergoes a second-order phase transition. The critical exponent of this transition is one-half, which coincides with the case in Landau-Ginzburg theory. In the condensed phase, a vector operator acquires a vacuum expectation value, breaking the Abelian symmetry as well as rotational symmetry spontaneously. Our calculation showed that this condensed phase exhibits infinite DC conductivity and a gap in the frequency-dependent conductivity, which is quite similar to properties of ordinary superconductivity.

We paid more attention to the response of this system to an applied magnetic field. We obtained the Landau level, from which we can find the contribution to the effective mass of the vector field by the magnetic field (see (24)). Due to the non-minimal coupling given in the last term of (1), the applied magnetic field can reduce the effective mass of the vector field, thus inducing instability of the black hole background even when the chemical potential/charge density is absent. That is, for the case with vanishing chemical potential or charge density, the black hole background becomes unstable when the magnetic field is beyond a certain critical value. For the case with non-vanishing chemical potential, the phase boundary is determined by a relation between the transition temperature and the magnetic field, which is presented in figure 3. The transition temperature increases with the applied magnetic field. The response of this system to the magnetic field is quite different from the behavior of ordinary superconductors where the magnetic field makes the transition more difficult, as drawn in figure 4. But our result is quite similar to the case of QCD vacuum instability induced by strong magnetic field to spontaneously develop the ρ -meson condensate [16, 17]. Although so, it was shown that in our model, the condensate of the vector operator forms a vortex lattice structure in the spatial directions perpendicular to the magnetic field. Of course, the non-minimal coupling term in the action plays a crucial role in both cases. Therefore, in some sense, our model is a holographic setup of the study of ρ -meson condensate in refs. [16, 17].

In ordinary superconductors, an external magnetic field suppresses superconductivity via diamagnetic and Pauli pair-breaking effects. However, it has also been proposed that magnetic field-induced superconductivity can be realized in type-II superconductors [34, 35], in which the Abrikosov flux lattice may enter a quantum limit of low Landau level dominance with spin-triplet pairing. Possible experimental evidence for strong magnetic field-induced superconductivity can be found, for example, in refs. [30, 31]. It was also shown in Gross-Neveu type models that applied magnetic fields might induce superconductivity in planar systems with 4-fermion interaction [36].

We mention here that similar studies can also be generalized to other gravitational backgrounds, such as the AdS soliton background which has been adopted to mimic superconductor/insulator phase transitions [37]. The study of magnetic field effects in the superconductor/insulator case can be found in ref. [38].

In a forthcoming paper [39], we generalize the present study to the case with the AdS soliton as the background. It is found that the magnetic field can induce instabilities of the AdS soliton background. Comparing our model with the SU(2) model with a constant non-Abelian magnetic field [23, 25], we find that our complex vector field model is in some sense a generalization of the SU(2) model to the case with a general mass squared m^2 and magnetic moment characterized by γ . In the setup of the present paper, the SU(2) model corresponds to our model with $m^2 = 0$ and $\gamma = 1$.

In this paper, we restricted ourselves to the probe approximation, neglecting the effect of matter fields on the background geometry. This can indeed reveal some significant properties of the model, but something might be lost in this approximation, see ref. [40] as an example. It is therefore helpful to understand the full properties of the model by considering the back-reaction of matter fields on the background geometry. In a recent paper [41], going beyond the probe approximation, we found a rich phase structure in this model without magnetic field. Depending on the mass squared m^2 and charge q of the vector field, not only second-order but also first-order and zeroth-order phase transitions can appear. Interestingly, there also exists a so-called “retrograde condensation” in which the hairy solution exists only for temperatures above a critical value and is thermodynamically subdominant. Particularly, the zeroth-order transition and retrograde condensation can be observed for the case with small m^2 . Indeed, this model has much richer phase behavior than the SU(2) model, thus it can be used to mimic much richer phenomena in dual strongly coupled systems.

In our model, it is the magnetic field itself that can induce the condensate. A natural question arises regarding the Meissner effect known as the phenomenon where superconductors expel weak external magnetic fields. This effect is due to the large superconducting currents induced in the superconductor by the external magnetic field, which generates a back-reacting magnetic field screening the external magnetic field. In the holographic SU(2) p-wave model, it has been pointed out [25] that the vortex currents flow in the opposite direction to those in conventional superconductors, thus can enhance the applied magnetic field in the regions between the vortices. It seems that the model presented here is similar to the SU(2) p-wave model. It is interesting to ask whether a similar phenomenon also appears in our model. To answer this question, we need to consider the contribution of higher-order terms in (5), which is also required to find the true vortex flux lattice. We leave this for further study.

In this paper we only calculated the conductivity in the case without magnetic field in one spatial direction. Although it is enough to see the superconductivity feature of the condensed phase, in order to study the model in a more realistic manner, it is desirable to calculate the conductivity in another direction and further to study the transport properties of the lattice state. As a phenomenological approach, this toy model would have potential application to mimic strongly coupled systems with a vector-like order parameter. It should be applicable in a wide variety of condensed matter systems, heavy ion physics

and beyond. We hope to report further progress in the future.

Acknowledgements

This work was supported in part by the National Natural Science Foundation of China (No.10821504, No.11035008, No.11205226, No.11305235, and No.11375247), and in part by the Ministry of Science and Technology of China under Grant No.2010CB833004.

Appendix A: Condensate of Vector Field and Superconducting Phase Transition

As a toy model with a charged vector field in the bulk dual to a vector operator in the field theory, there exists the possibility that the condensate of this vector field can serve as an order parameter to mimic a holographic p-wave superconductor phase transition, like the s-wave case [4, 5]. More precisely, we hope that this system has stable black hole solutions with vector hair at low temperatures, but without vector hair at high temperatures. If this is true, in the condensed phase the condensate of the dual vector operator will break not only the U(1) symmetry but also the rotational symmetry since the condensate of the vector field picks out one special direction. This situation is very similar to the one in the holographic p-wave superconductor model with SU(2) gauge field in ref. [12]. In this sense, our model can also be regarded as a holographic p-wave model.

A.1 Condensate of Vector Field

We adopt the following ansatz

$$\rho_\nu dx^\nu = \rho_x(r)dx + \rho_y(r)dy, \quad A_\nu dx^\nu = \phi(r)dt.$$

One can use the U(1) gauge symmetry to set ρ_x to be real. Then one finds that the r component of (2) implies that the phase of ρ_y must be constant. Without loss of generality, we take ρ_y to be real. Then, the independent equations of motion in terms of the above ansatz are deduced as follows:

$$\rho_x'' + \left(\frac{f'}{f} + \frac{2}{r} \right) \rho_x' + \frac{q^2 \phi^2}{f^2} \rho_x = 0,$$

$$\rho_y'' + \left(\frac{f'}{f} + \frac{2}{r} \right) \rho_y' + \frac{q^2 \phi^2}{f^2} \rho_y = 0,$$

$$\phi'' + \frac{2}{r} \phi' - \frac{2(\rho_x^2 + \rho_y^2)}{r^2 f} \phi = 0,$$

where the prime denotes the derivative with respect to r .

In order to find solutions for all three functions, one must impose suitable boundary conditions at the AdS boundary $r \rightarrow \infty$ and at the horizon $r = r_h$. In addition to $f(r_h) = 0$, one must require $\phi(r_h) = 0$ in order for $g_{\mu\nu}A^\mu A^\nu$ to be finite at the horizon. In order to match the asymptotical AdS boundary, the general falloff of the matter fields near the boundary $r \rightarrow \infty$ should behave as

$$\rho_x = \frac{\rho_{x-}}{r^{\Delta_-}} + \frac{\rho_{x+}}{r^{\Delta_+}} + \dots, \quad \rho_y = \frac{\rho_{y-}}{r^{\Delta_-}} + \frac{\rho_{y+}}{r^{\Delta_+}} + \dots, \quad \phi = \mu - \frac{\rho}{r} + \dots,$$

where $\Delta_\pm = 1 \pm \sqrt{1 + 4m^2}/2$. We impose $\rho_{x-} = 0$ and $\rho_{y-} = 0$, since we want the U(1) symmetry to be broken spontaneously. According to the AdS/CFT dictionary, up to a normalization, the coefficients μ , ρ , ρ_{x+} and ρ_{y+} are interpreted as chemical potential, charge density, and the x and y components of the vacuum expectation value of the vector operator \hat{J}^μ in the dual field theory, respectively.

There is a useful scaling symmetry in the equations of motion (37):

$$(t, x, y) \rightarrow \lambda(t, x, y), \quad (\phi, \rho_x, \rho_y) \rightarrow \lambda(\phi, \rho_x, \rho_y),$$

where λ is an arbitrary positive constant. Under this symmetry, the relevant quantities transform as

$$T \rightarrow \lambda T, \quad \mu \rightarrow \lambda \mu, \quad (\rho_{x+}, \rho_{y+}) \rightarrow \lambda^{\Delta_+ + 1}(\rho_{x+}, \rho_{y+}).$$

The condensate as a function of temperature is presented in figure 6 [Figure 6: see original paper]. We choose $q = 1$ and $m^2 = 3/4$. The condensate begins to appear at $T_c \simeq 0.102\sqrt{\rho}$. We assume the condensate picks out the x direction as special, so we can consistently set $\rho_y = 0$. It is clear that as we lower the temperature, the normal phase with vanishing charged vector becomes unstable to developing vector hair which breaks the U(1) symmetry as well as rotational symmetry in the dual field theory. Fitting the curve near the critical temperature T_c , we find that for small condensate there is a critical behavior with critical exponent 1/2, which precisely meets the result given by mean field theory and is typically a second-order phase transition. In the case with $m^2 = 3/4$ we obtain $\langle \hat{J}_x \rangle \simeq 339T_c(1 - T/T_c)^{1/2}$ as $T \rightarrow T_c$.

Thus we have obtained two black hole solutions in the system. When $T > T_c$, we have the black hole solution without the vector field, while we have the black hole solution with non-trivial vector field as $T < T_c$. This behavior is in complete agreement with the holographic superconducting phase transition in the literature. Therefore we expect that the black hole solution with non-trivial vector field can describe a superconducting phase. To prove this, it is helpful to calculate the optical conductivity. Before doing this, we should first ensure that

the black hole solution with non-trivial vector field is more thermodynamically stable than the one without the vector field as $T < T_c$.

A.2 Free Energy

In order to determine which solution is thermodynamically favored, we should calculate the free energy of the system for both black hole solutions. We will work in the canonical ensemble in this paper, where the charge density is fixed. In gauge/gravity duality, the Helmholtz free energy F of the boundary thermal state is identified with temperature times the on-shell bulk action with Euclidean signature. Since we work in the probe approximation, we can ignore the gravity part. Given that the system is stationary, the Euclidean action is related to the Minkowski case by a total minus. Employing the equations of motion (2) and (3), we have

$$\frac{1}{2\kappa^2} S_{\text{Euclidean}} = \int d^4x \sqrt{-g} \left[\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} \rho_{\mu\nu}^\dagger \rho^{\mu\nu} + m^2 \rho_\mu^\dagger \rho^\mu - iq\gamma \rho_\mu \rho_\nu^\dagger F^{\mu\nu} \right] + S_{\text{ct}},$$

where S_{ct} denotes the surface counter term for removing divergence. Substituting the asymptotical expansion (39) into (43) and introducing a counter term $S_{\text{ct}} = \int d^3x \sqrt{h} h^{\mu\nu} \rho_\mu^\dagger \rho_\nu$, we find the free energy F as

$$F = \int d^3x \sqrt{-g} \left[\frac{1}{2} A_\nu \nabla_\mu F^{\mu\nu} + \frac{1}{2} (\rho_\nu^\dagger D_\mu \rho^{\mu\nu} + \text{h.c.}) \right] + S_{\text{ct}}.$$

Regarding ρ_{x-} and ρ_{y-} as sources, the Helmholtz free energy tells us that the sub-leading terms ρ_{x+} and ρ_{y+} are the expectation values of the dual operator in the field theory side. The difference of Helmholtz free energy between the condensed phase and the normal phase as a function of temperature is presented in figure 7 [Figure 7: see original paper]. It is clear that below the critical temperature T_c , the state with non-vanishing vector “hair” is indeed thermodynamically favored compared to the normal phase. The phase transition is second order, which can be seen, for example, from the derivative of the free energy with respect to temperature.

A.3 Conductivity

We now calculate the conductivity in the dual field theory side as a function of frequency ω . We need to turn on fluctuations of the matter contents in the bulk. We assume perturbations have a time dependence of the form $e^{-i\omega t}$ with zero spatial momentum. It turns out that one can consistently turn on the perturbations of A_y only. We can obtain the equation of motion for A_y by linearizing the equations of motion (2), which reads

$$A_y'' + \left(\frac{f'}{f} + \frac{2}{r} \right) A_y' + \left(\frac{\omega^2}{f^2} - \frac{2q^2 \rho_x^2}{r^2 f} \right) A_y = 0.$$

Since the conductivity is related to the retarded two-point function of the U(1) current, we impose the ingoing boundary condition near the horizon. The gauge field A_y near the boundary $r \rightarrow \infty$ falls off as

$$A_y = A^{(0)} + \frac{A^{(1)}}{r} + \dots.$$

According to the AdS/CFT dictionary, one can obtain the conductivity as

$$\sigma_{yy}(\omega) = \frac{i}{\omega} \frac{A^{(1)}}{A^{(0)}}.$$

The AC conductivity as a function of frequency is presented in figure 8 [Figure 8: see original paper]. We can see clearly that the optical conductivity along the y direction in this model behaves qualitatively similar to the case in the p-wave model with SU(2) gauge symmetry [12]. In particular, from the Kramers-Kronig relation, one can conclude that the real part of the conductivity has a Dirac delta function at $\omega = 0$ since the imaginary part has a pole, i.e., $\text{Im}[\sigma_{yy}(\omega)] \sim 1/\omega$. Furthermore, it is clear that the optical conductivity develops a gap at some special frequency ω_g known as the gap frequency. As suggested in ref. [42], it can be identified with the one at the minimum of the imaginary part of the AC conductivity. $\text{Re}[\sigma_{yy}]$ is very small in the infrared and rises quickly at ω_g . There also exists a small “bump” slightly above ω_g which is reminiscent of the behavior due to fermionic pairing [12]. For our chosen parameters, we have $\omega_g \simeq 3.5T_c$, compared to the corresponding BCS value $\omega_g \simeq 8T_c$. The result shown here is consistent with the fact that our holographic model describes a system at strong coupling.

References

- [1] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” *Adv. Theor. Math. Phys.* 2, 231 (1998) [*Int. J. Theor. Phys.* 38, 1113 (1999)] [arXiv:hep-th/9711200].
- [2] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” *Phys. Lett. B* 428, 105 (1998) [arXiv:hep-th/9802109].
- [3] E. Witten, “Anti-de Sitter space and holography,” *Adv. Theor. Math. Phys.* 2, 253 (1998) [arXiv:hep-th/9802150].
- [4] S. A. Hartnoll, C. P. Herzog and G. T. Horowitz, “Building a Holographic Superconductor,” *Phys. Rev. Lett.* 101, 031601 (2008) [arXiv:0803.3295 [hep-th]].

- [5] S. A. Hartnoll, C. P. Herzog and G. T. Horowitz, “Holographic Superconductors,” JHEP 0812, 015 (2008) [arXiv:0810.1563 [hep-th]].
- [6] S. S. Gubser, C. P. Herzog, S. S. Pufu and T. Tesileanu, “Superconductors from Superstrings,” Phys. Rev. Lett. 103, 141601 (2009) [arXiv:0907.3510 [hep-th]].
- [7] J. P. Gauntlett, J. Sonner and T. Wiseman, “Holographic superconductivity in M-Theory,” Phys. Rev. Lett. 103, 151601 (2009) [arXiv:0907.3796 [hep-th]].
- [8] M. Ammon, J. Erdmenger, M. Kaminski and P. Kerner, “Superconductivity from gauge/gravity duality with flavor,” Phys. Lett. B 680, 516 (2009) [arXiv:0810.2316 [hep-th]].
- [9] N. Bobev, A. Kundu, K. Pilch and N. P. Warner, “Minimal Holographic Superconductors from Maximal Supergravity,” JHEP 1203, 064 (2012) [arXiv:1110.3454 [hep-th]].
- [10] J.-W. Chen, Y.-J. Kao, D. Maity, W.-Y. Wen and C.-P. Yeh, “Towards A Holographic Model of D-Wave Superconductors,” Phys. Rev. D 81, 106008 (2010) [arXiv:1003.2991 [hep-th]].
- [11] F. Benini, C. P. Herzog, R. Rahman and A. Yarom, “Gauge gravity duality for d-wave superconductors: prospects and challenges,” JHEP 1011, 137 (2010) [arXiv:1007.1981 [hep-th]].
- [12] S. S. Gubser and S. S. Pufu, “The Gravity dual of a p-wave superconductor,” JHEP 0811, 033 (2008) [arXiv:0805.2960 [hep-th]].
- [13] F. Aprile, D. Rodriguez-Gomez and J. G. Russo, “p-wave Holographic Superconductors and five-dimensional gauged Supergravity,” JHEP 1101, 056 (2011) [arXiv:1011.2172 [hep-th]].
- [14] L. A. Pando Zayas and D. Reichmann, “A Holographic Chiral $p + ip$ Superconductor,” Phys. Rev. D 85, 106012 (2012) [arXiv:1108.4022 [hep-th]].
- [15] D. E. Kharzeev, K. Landsteiner, A. Schmitt and H.-U. Yee, “ ‘Strongly interacting matter in magnetic fields’ : an overview,” Lect. Notes Phys. 871, 1 (2013) [arXiv:1211.6245 [hep-ph]].
- [16] M. N. Chernodub, “Superconductivity of QCD vacuum in strong magnetic field,” Phys. Rev. D 82, 085011 (2010) [arXiv:1008.1055 [hep-ph]].
- [17] M. N. Chernodub, “Spontaneous electromagnetic superconductivity of vacuum in strong magnetic field: evidence from the Nambu-Jona-Lasinio model,” Phys. Rev. Lett. 106, 142003 (2011) [arXiv:1101.0117 [hep-ph]].
- [18] M. N. Chernodub, J. Van Doorselaere and H. Verschelde, “Electromagnetically superconducting phase of vacuum in strong magnetic field: structure of superconductor and superfluid vortex lattices in the ground state,” Phys. Rev. D 85, 045002 (2012) [arXiv:1111.4401 [hep-ph]].

- [19] D. Djukanovic, M. R. Schindler, J. Gegelia and S. Scherer, “Quantum electrodynamics for vector mesons,” *Phys. Rev. Lett.* 95, 012001 (2005) [hep-ph/0505180].
- [20] Y. Nambu and G. Jona-Lasinio, “Dynamical Model of Elementary Particles Based on an Analogy with Superconductivity. 1.,” *Phys. Rev.* 122, 345 (1961).
- [21] N. Callebaut, D. Dudal and H. Verschelde, “Holographic rho mesons in an external magnetic field,” *JHEP* 1303, 033 (2013) [arXiv:1105.2217 [hep-th]].
- [22] N. Callebaut and D. Dudal, “On the transition temperature(s) of magnetized two-flavour holographic QCD,” arXiv:1303.5674 [hep-th].
- [23] M. Ammon, J. Erdmenger, P. Kerner and M. Strydom, “Black Hole Instability Induced by a Magnetic Field,” *Phys. Lett. B* 706, 94 (2011) [arXiv:1106.4551 [hep-th]].
- [24] Y.-Y. Bu, J. Erdmenger, J. P. Shock and M. Strydom, “Magnetic field induced lattice ground states from holography,” *JHEP* 1303, 165 (2013) [arXiv:1210.6669 [hep-th]].
- [25] K. Wong, “A non-abelian vortex lattice in strongly-coupled systems,” arXiv:1307.7839 [hep-th].
- [26] N. K. Nielsen and P. Olesen, “An Unstable Yang-Mills Field Mode,” *Nucl. Phys. B* 144, 376 (1978).
- [27] J. Ambjorn and P. Olesen, “Antiscreening Of Large Magnetic Fields By Vector Bosons,” *Phys. Lett. B* 214, 565 (1988).
- [28] J. Ambjorn and P. Olesen, “A Magnetic Condensate Solution Of The Classical Electroweak Theory,” *Phys. Lett. B* 218, 67 (1989) [Erratum-ibid. B 220, 659 (1989)].
- [29] T. Albash and C. V. Johnson, “A Holographic Superconductor in an External Magnetic Field,” *JHEP* 0809, 121 (2008) [arXiv:0804.3466 [hep-th]].
- [30] F. Levy, I. Sheikin, B. Grenier, A. Huxley, “Magnetic Field-induced Superconductivity in the Ferromagnet URhGe,” *Science* 309, 1343 (2005).
- [31] S. Uji, H. Shinagawa, T. Terashima, T. Yakabe, Y. Terai, M. Tokumoto, A. Kobayashi, H. Tanaka, H. Kobayashi, “Magnetic-field-induced superconductivity in a two-dimensional organic conductor,” *Nature* 410, 908 (2010).
- [32] J. A. Young and S. A. Bludman, “Electromagnetic Properties of a Charged Vector Meson,” *Phys. Rev.* 131, 2326 (1963).
- [33] K. Maeda, M. Natsuume and T. Okamura, “Vortex lattice for a holographic superconductor,” *Phys. Rev. D* 81, 026002 (2010) [arXiv:0910.4475 [hep-th]].
- [34] Rasolt. Mark, “Superconductivity in high magnetic fields,” *Phys. Rev. Lett.* 58, 1482 (1987).

- [35] M. Rasolt and Z. Tesanovic, “Theoretical aspects of superconductivity in very high magnetic fields,” *Rev. Mod. Phys.* 64, 709 (1992).
- [36] K. G. Klimenko, R. N. Zhokhov and V. C. Zhukovsky, “Superconductivity phenomenon induced by external in-plane magnetic field in (2+1)-dimensional Gross-Neveu type model,” *Mod. Phys. Lett. A* 28, 1350096 (2013) [arXiv:1211.0148 [hep-th]].
- [37] T. Nishioka, S. Ryu and T. Takayanagi, “Holographic Superconductor/Insulator Transition at Zero Temperature,” *JHEP* 1003, 131 (2010) [arXiv:0911.0962 [hep-th]].
- [38] R.-G. Cai, L. Li, H.-Q. Zhang and Y.-L. Zhang, “Magnetic Field Effect on the Phase Transition in AdS Soliton Spacetime,” *Phys. Rev. D* 84, 126008 (2011) [arXiv:1109.5885 [hep-th]].
- [39] R.-G. Cai, L. Li, L.-F. Li and Y. Wu, “Vector Condensate and AdS Soliton Instability Induced by a Magnetic Field,” arXiv:1311.7578 [hep-th].
- [40] R.-G. Cai, L. Li, L.-F. Li and Y.-Q. Wang, “Competition and Coexistence of Order Parameters in Holographic Multi-Band Superconductors,” *JHEP* 1309, 074 (2013) [arXiv:1307.2768 [hep-th]].
- [41] R.-G. Cai, L. Li and L.-F. Li, “A Holographic P-wave Superconductor Model,” arXiv:1309.4877 [hep-th].
- [42] G. T. Horowitz and M. M. Roberts, “Holographic Superconductors with Various Condensates,” *Phys. Rev. D* 78, 126008 (2008) [arXiv:0810.1077 [hep-th]].

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv – Machine translation. Verify with original.