

## The morphology of *Chiappeavis magnapremaxillo* (Pengornithidae: Enantiornithes) and a comparison of aerodynamic function in Early Cretaceous avian tail fans (Postprint)

**Authors:** Jingmai K. O' CONNOR, ZHENG Xiao-Ting, HU Han, WANG Xiao-Li, ZHOU Zhong-He, Jingmai K. O' CONNOR

**Date:** 2017-08-10T00:00:00+00:00

### Abstract

*Chiappeavis* is the first enantiornithine discovered to preserve a rectricial fan, showing that the rectricial bulb structure was already developed in relatively primitive enantiornithines. We provide a detailed description of the skeletal morphological features of the holotype specimen of *Chiappeavis magnapremaxillo*. The palatal morphology of *Chiappeavis* is similar to that of *Archaeopteryx*, but distinct from the Late Cretaceous enantiornithine *Gobipteryx*. Even with rectricial bulbs, the pygostyle morphology of pengornithids indicates that this structure was poorly developed. We estimate the lift generated by the rectricial fan in *Chiappeavis* and compare it with other Early Cretaceous birds. The results show that the aerodynamic lift produced by the rectricial fan of *Chiappeavis* is less than that of sympatric ornithuromorphs, which may explain the widespread absence of aerodynamically functional tail morphologies in Enantiornithes.

### Full Text

## The Morphology of *Chiappeavis magnapremaxillo* (Pengornithidae: Enantiornithes) and a Comparison of Aerodynamic Function in Early Cretaceous Avian Tail Fans

Jingmai K. O' Connor<sup>1</sup>, Zheng Xiao-Ting<sup>2,3</sup>, Hu Han<sup>1</sup>, Wang Xiao-Li<sup>2</sup>, Zhou Zhong-He<sup>1</sup>

<sup>1</sup>Key Laboratory of Vertebrate Evolution and Human Origins, Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences, Beijing 100044, China

<sup>2</sup>Institute of Geology and Paleontology, Linyi University, Linyi, Shandong 276000, China

<sup>3</sup>Shandong Tianyu Museum of Nature, Pingyi, Shandong 273300, China

## Abstract

We provide a complete description of the skeletal anatomy of the holotype of *Chiappeavis magnapremaxillo*, the first enantiornithine to preserve a rectricial fan, suggesting that rectricial bulbs were possibly present in basal members of this clade. Notably, *Chiappeavis* preserves a primitive palatal morphology in which the vomers reach the premaxillae, similar to *Archaeopteryx* but unlike the condition in the Late Cretaceous enantiornithine *Gobipteryx*. If rectricial bulbs were present, pengornithid pygostyle morphology suggests they were minimally developed. We estimate the lift generated by the tail fan preserved in this specimen and compare it to the tail fans preserved in other Early Cretaceous birds. Aerodynamic models indicate the tail of *Chiappeavis* produced less lift than that of sympatric ornithuromorphs. This information provides a possible explanation for the absence of widespread aerodynamic tail morphologies in the Enantiornithes.

**Key words:** Mesozoic, Jehol Biota, Aves, rectrix

**Citation:** O' Connor JK, Zheng XT, Hu H et al., 2016. The morphology of *Chiappeavis magnapremaxillo* (Pengornithidae: Enantiornithes) and a comparison of aerodynamic function in Early Cretaceous avian tail fans. *Vertebrata Palasiatica*, 55(1): 41-58

---

## 1 Introduction

Enantiornithes is the most diverse recognized group of Mesozoic birds, considered the first major avian radiation. Although specimens have been collected from continental and marine sediments on all continents except Antarctica, like other bird fossils, these remains are typically isolated and fragmentary (O' Connor et al., 2011). The major exception is the Early Cretaceous deposits in northeastern China that have produced the 130.7–120 Ma Jehol Biota (Pan et al., 2013), preserving the second oldest recognized fossil avifauna, surpassed only by the Late Jurassic Solnhofen Limestones in Germany that produce *Archaeopteryx* (Wellnhofer, 2008). Despite its age, the Jehol avifauna accounts for approximately half of the entire diversity of Mesozoic birds, including a huge diversity of enantiornithines (Wang M et al., 2014; Wang X et al., 2014; Zhou and Zhang, 2006). As new species continue to be discovered, several distinct clades have been recognized (e.g., the Bohaiornithidae, Longipterygidae). The most temporally successful of these enantiornithine lineages is the Pengornithidae. The first specimen, the holotype of *Pengornis houi*, was described by Zhou et al. in 2008, and already this group is one of the most diverse enantiornithine clades in the Jehol avifauna. Currently, there are five specimens representing

four genera (*Pengornis*, *Parapengornis*, *Eopengornis*, and *Chiappeavis*) (Hu et al., 2015; O' Connor et al., 2016; Wang X et al., 2014; Zhou et al., 2008). The holotype of *Pengornis houi* is the largest known Early Cretaceous enantiornithine. *Eopengornis* is from the 130.7 Ma Propteryx-horizon of the Huajiyang Formation, and all other taxa are from the 120 Ma Jiufotang Formation. Pengornithids are the most recognizable enantiornithines, with their characteristic numerous small, low-crowned teeth, hooked scapular acromion, bilaterally formed sternum without intermediate trabeculae, elongate femur, unreduced fibula, metatarsal V, and elongate metatarsal I and hallux. These characters are mostly primitive features, strongly suggesting that pengornithids are basal within the Enantiornithes, consistent with recent phylogenetic analyses and the fact that *Eopengornis* is among the oldest known enantiornithines (Wang X et al., 2014).

The most recently described pengornithid, *Chiappeavis magnapremaxillo*, preserves the first clear evidence that some members of the Enantiornithes possessed rectricial fans (Fig. 1 [Figure 1: see original paper]), similar to those present in the basal pygostylian *Sapeornis* and members of the Ornithuromorpha (Clarke et al., 2006; Zheng et al., 2013), the clade that includes all living birds. The unique shape of the pengornithid pygostyle, being relatively more similar to that of ornithuromorphs and *Sapeornis* than other enantiornithines, suggests that the rectricial fan evolves together with the rectricial bulbs necessary to control them, which in turn shapes the pygostyle (O' Connor et al., 2016). The holotype and only known specimen of *Chiappeavis magnapremaxillo* STM 29-11 was only briefly described with regards to its skeletal morphology. Here we provide a complete description and further explore the unique morphology of the pengornithid pygostyle. We reconstruct the tail fans of several well-preserved Jehol ornithuromorphs and compare the aerodynamic lift generated by different tail shapes and discuss the significance of this information.

---

## 2 Description

This description provides only new anatomical information regarding STM 29-11 (Fig. 2 [Figure 2: see original paper]). The enlarged premaxillary corpus and elongate nasal processes that distinguish *Chiappeavis* from other pengornithids have already been described in detail (O' Connor et al., 2016). The maxillary process of the premaxilla is robust, sharply tapered, and roughly equal to the length of the corpus (Fig. 3 [Figure 3: see original paper]). The premaxillae form a medial wedged articulation with the nasals, although the extent of this articulation is unclear due to poor preservation of the nasals. A fragment of the left nasal is preserved articulating with the frontal; it appears narrower than the nasal in *Pengornis* IVPP V 15336 (Zhou et al., 2008). Although the maxilla forms most of the facial margin in *Pengornis* V 15336 (Zhou et al., 2008) and all other known pengornithids (Hu et al., 2014, 2015; Wang X et al., 2014), only possible fragments are preserved in *Chiappeavis* STM 29-11. Part

of the scleral ring is preserved in the orbit. Other fragments may represent pieces of the pterygoids. Ventrally, a thin, distally upturned rod-like element is identified as the jugal. A robust fragment of bone is preserved dorsal and parallel to the jugal; this bone may be a piece of the jugal process of the maxilla, but it appears unusually robust. Part of the palate is visible through the left external nares (Fig. 3). Both vomers are preserved, presumably in dorsal view. As in *Gobipteryx*, they appear to be fused along their rostral halves, caudally diverging at a  $15^\circ$  angle (nearly parallel in *Gobipteryx*). This suggests that, like *Gobipteryx*, the choana was rostrally located compared to neornithines (Chiappe et al., 2001). However, unlike in *Gobipteryx* and neognathous birds, as preserved the vomers appear to retain the plesiomorphic tetrapod condition, reaching and presumably articulating with the premaxillae, as in paleognathous birds and potentially *Archaeopteryx* (Witmer and Martin, 1987). A fragment of bone preserved between the left vomer and the maxillary process of the premaxilla may be a piece of the maxillary contribution to the palate. The caudal half of the skull is poorly preserved, crushed, and heavily abraded, revealing no anatomical details. The tip of the left dentary is blunt; barely visible, numerous small teeth are preserved in separate alveoli. The postdentary bones are poorly preserved.

Approximately seven cervical vertebrae are poorly preserved in articulation with the skull (Fig. 2). The cranial articular surfaces appear to be slightly concave. Proximally, the cervicals are preserved in ventral view; distally, the series appears in caudal view. The third preserved vertebra reveals small carotid processes. The costal processes are short and sharply tapered. The proximal portion of the thoracic series is obscured by overlap with the thoracic girdle. There are five visible dorsal vertebrae; the first two are disarticulated, but the distal three are in articulation with the synsacrum. The dorsal vertebrae show the typical enantiornithine condition, with the parapophyses located one-third from the proximal end and the lateral surface deeply excavated by a groove (Chiappe and Walker, 2002). The neural spines of the synsacrum are fused into a continuous spinous crest, which narrows and decreases in height distally, as in V 15336.

The first four transverse processes are quadrangular, short, and wide; the last three are elongate and caudolaterally oriented, and the fifth is intermediate in morphology (Figs. 1, 2). Five free caudal vertebrae are preserved, which is fewer than reported in other pengornithids (Hu et al., 2014, 2015); the cranial articular surface is exposed in the first caudal vertebra, revealing a weakly concave surface. The transverse processes are long, approximately equal to the centrum width. The neural canal is smaller than the size of the exposed caudal articular surfaces. The unfused haemal arches are rectangular with blunt distal margins.

As noted in a more detailed study of the pygostyle (Wang and O' Connor, in press), the pengornithid pygostyle bears all the same characteristic features of other enantiornithines, although this clade differs in how these morphologies are expressed. The pygostyle of *Chiappeavis* has a pair of ventrolateral processes

restricted to the proximal ventral half of the pygostyle, whereas in other enantiornithines these processes extend for most of the pygostyle length (Chiappe and Walker, 2002). The ventrolateral processes project further ventrally and are limited to the proximal third of the pygostyle in *Pengornis* V 15336 (Fig. 4B [Figure 4: see original paper]). Visible on the right, a small cranial fork is present and continuous with a dorsolateral process that appears to extend the entire length of the pygostyle (Fig. 4A). The dorsal surface between these dorsolateral processes is broadly concave, as in other pengornithids, whereas this concavity is narrow and deeper in other enantiornithines, even when preserved dorsoventrally crushed (Wang and O' Connor, in press).

The left ramus of the furcula is preserved in dorsal view; the rami are slightly bowed medially, similar to *Pengornis* V 15336 (Fig. 5 [Figure 5: see original paper]). The dorsolateral excavation is limited to the proximal 2/3 of the ramus; the dorsal surface of the omal third is flat. The omal tip tapers bluntly, as in *Pengornis* V 15336, and is pitted and striated, suggesting incomplete ossification of the articular surfaces. A hypocleidium was present, as in other pengornithids. As in *Pengornis*, the process measures approximately half the length of the furcular rami, whereas the hypocleidium appears proportionately longer in *Parapengornis* IVPP V 18687 (Fig. 5). Both coracoids are in articulation with the coracoidal sulci of the sternum in dorsal view; the dorsal lips of the sulci cover the sternal margin of the coracoids, indicating that the sulci are deep. The acrocoracoid, glenoid, and scapular articular surface are weakly aligned, as in other enantiornithines (Chiappe and Walker, 2002). The poorly developed glenoid and scapular cotyla are separated by a deep pit, which may be an artifact of crushing and obscuring the presence of an acrocoracoidal tubercle. Just distal to the scapular articular surface, the supracoracoid nerve foramen pierces the neck of the coracoid, separated from the medial margin by a complete bony bar; it does not appear to open into a medially located groove as it does in many other enantiornithines (Chiappe and Walker, 2002). The corpus makes up the distal half of the coracoid. The distal quarter of the lateral margin is convex, expanding the width of the sternal margin. The convex distal portion of the lateral margin appears distinctly thinner and flatter than the rest of the coracoidal corpus. This morphology of the lateral margin of the coracoid is also observed in other pengornithids (*Eopengornis*, *Parapengornis* V 18687) as well as bohaiornithid enantiornithines (Wang M et al., 2014). The corpus is weakly excavated by a shallow dorsal fossa, also observed in some ornithuromorph birds (e.g., *Yixianornis*), inferred to be the attachment of the m. supracoracoideus (Clarke et al., 2006).

The left scapula in lateral view overlaps the right in costal view, and where the two shafts overlap, the left is missing a piece of the shaft; both distal ends are unclear. The scapular acromion process is slightly longer than the glenoid facet and hooked, as in other pengornithid enantiornithines (Wang X et al., 2014). The cranial margin of the process is wide and flat. The scapular glenoid facet is large, concave, slightly tapered distally, and forms a labrum where it contacts the scapular blade (Fig. 6 [Figure 6: see original paper]). The body of

the scapula is relatively wide and short, as in other pengornithids. The costal surface is smooth, lacking the groove present in more derived enantiornithines (e.g., *Elsornis*, *Neuquenornis*) (Chiappe and Walker, 2002).

The straight coracoidal sulci meet at a  $120^\circ$  angle so that the rostral margin of the sternum is weakly vaulted (Fig. 6). The cranio-lateral corners are weakly developed into slight dorsolateral projections, but no distinct cranio-lateral process is present. The entire lateral margin of the sternum, including the lateral trabecula, is weakly concave. The dorsal surface of each trabecula is keeled, giving this process a triangular cross-section; distally, the apex moves from dorso-laterally located to centered on the dorsal surface, and the distal third is flat. The distal ends of the trabeculae are weakly expanded—this area of bone is also heavily pitted and striated, indicating ossification was incomplete. The lateral trabeculae extend distally beyond the caudal margin of the xiphial region, as in V 18632 (level with caudal margin in *Eopengornis* STM 24-11). The lateral margins of the median trabeculae are concave as the sternal plates narrow caudally, whereas they are straight in *Eopengornis* and *Parapengornis*.

Compared to other known pengornithids, the xiphial region is more elongate and narrow in STM 29-11, forming an incipient xiphoid process and defining approximately a  $40^\circ$  angle ( $75^\circ$  in *Eopengornis* and  $70^\circ$  in *Parapengornis*). The xiphial region bears a short, straight caudal margin, absent in *Eopengornis* and V 18632, in which the xiphial region defines a blunt V-shaped margin (Hu et al., 2014; Wang X et al., 2014). Given the lack of maturity in all specimens preserving sternal material, these apparent differences in the morphology of the distal ends of the lateral trabeculae and proportions of the xiphial region may change with the discovery of adult material. However, features like the concavity of the lateral margins in *Chiappeavis* STM 29-11 are unlikely to change at this stage in maturity.

Well-defined costal facets are not visible, but three robust sternal ribs are visible articulating on the right costal margin of the sternum. The third rib is more robust than the others and has an uneven caudal margin that may be pathological in origin (Fig. 6). The left sternal ribs are displaced over the dorsal surface of the sternum—they are short, robust, and weakly curved. Several disarticulated thoracic ribs are preserved; a few are located cranial to the sternum, and a few are associated with this element. Compared to the cranially located ribs, the ribs preserved near and overlying the sternum are shorter, more robust, and more weakly curved, and probably articulated with the sternal ribs.

The left humerus is preserved in cranial view, while the right is in caudal view. Proximally in caudal view, the ventral tubercle is small, separated by a wide, shallow capital incision, and a pneumatic fossa is absent. In cranial view, a small bicipital crest is present, weakly projecting cranially. The width of the deltopectoral crest is less than the width of the shaft and extends along the proximal third of the humerus. The shaft weakly increases in width distally from its narrowest point located midshaft, also observed in other pengornithids. The distal caudal surface is poorly preserved, revealing no anatomical details. The distal

condyles are small and located primarily on the cranial surface, as in other birds. The circular dorsal condyle is larger than the oval ventral condyle, although the two are approximately equal in diameter. The long axis of the ventral condyle is transversely oriented. The round ventral epicondyle is smaller than the ventral condyle and located on the craniodistal surface of the ventrodistal margin. The brachial fossa is not developed.

Both antebrachia are complete; the left is in caudal view, and the right is cranially exposed. As in other basal birds, the ulnae are bowed along their proximal halves. The dorsal cotyla is flat, and the ventral cotyla appears slightly concave. The radii are straight and more than half the thickness of the ulna; crushing prevents identification of a longitudinal groove like that present on the interosseous surface of some enantiornithines (Chiappe and Walker, 2002). The ulnare is U-shaped; one ramus is bluntly tapered, while the other is short and more robust, similar to that of *Pengornis* V 15336. The radiale is quadrangular, but it is unclear which surface is exposed. The semilunate carpal appears only partially fused to the metacarpals.

The right alular metacarpal is preserved unfused to the major metacarpal. It is narrow and approximately 1/5 the length of the carpometacarpus. The proximal end is rounded, and the distal end appears ginglymous. As in V 15336, the proximal end of the minor metacarpal wraps onto the ventral surface of the major metacarpal, and the ventral surface of the proximal third of the minor metacarpal forms a ridge rather than a distinct pisiform process, a morphology common in Early Cretaceous enantiornithines (O' Connor, 2009). The first phalanx of the major digit is incomplete, but like other enantiornithines, it maintains the plesiomorphic theropod condition and lacks any caudal expansion like that present in ornithuromorph birds. The penultimate phalanx is not preserved; the ungual phalanx is small and weakly curved. The first phalanx of the minor digit is wedge-shaped with a flat cranial margin and convex caudal margin, tapering to a blunt distal margin.

Both ilia are preserved in articulation with the synsacrum (Figs. 1, 2). The right ilium is preserved in medial view, while the left appears in dorsal view. In lateral view, the dorsal margin is weakly convex. The ventral margins of the preacetabular and postacetabular alae are straight. The pubic pedicel of the ischium is wider than the iliac pedicel. The preacetabular wing is longer and dorsoventrally taller than the postacetabular wing, and the postacetabular wing is bluntly tapered, as in V 18632 and most enantiornithines. The right ischium, preserved in medial view, is long, delicate, and gently tapered; the dorsal margin is concave, and the ventral margin is convex (Figs. 1, 2). The pubes have a thick oval cross-section with a craniodorsal-caudoventrally oriented long axis. The distal third is heavily pitted and striated.

Both femora are preserved, although the right is overlain by the pubes and ischia, and the proximal end is not exposed on either side (Figs. 1, 2). As described, a small tibiofibular crest is present distally (O'Connor et al., 2016). The right tibia is in caudal view, exposing the popliteal tuberosity and lateral articular facet on

the proximocaudal surface, also developed in *Pengornis* and *Eopengornis* (Wang X et al., 2014). The flexor fossa is weak or absent. The left tibia is exposed in craniolateral view; the proximal tarsals are fused to each other, forming the condyles. The medial condyle is wider than the lateral condyle. They are separated by a wide, shallow intercondylar incisure. The lateral margin of the lateral condyle and the medial margin of the medial condyle are both straight, and the opposing surfaces are strongly convex, as in *Pengornis* V 15336. The lateral condyle is excavated by a deep concavity on the lateral surface, as in some enantiornithines (e.g., *Qiliania*) (Ji et al., 2011). The ascending process is triangular, tapering proximally, and taller than the height of the condyles. Only the right fibula is preserved, and it is incomplete, so it cannot be determined if it reached the distal condyles as in other pengornithids (Wang X et al., 2014).

No distal tarsals are preserved, as in other pengornithids. The metatarsals are unfused and slightly disarticulated; the exposed surfaces are abraded, preserving very little information. Metatarsal IV is thinner than metatarsal III, which is in turn thinner than metatarsal II. The digit of metatarsal II is robust with a large ungual phalanx; the ungual phalanx in digit III is long but not as robust as those of digits I and II. The digit IV ungual is the smallest. No metatarsal V is preserved, although it was likely present, as in other pengornithids (Wang X et al., 2014).

**Ontogenetic status.** STM 29-11 is clearly immature based on the incomplete ossification of the periosteal surface of the bone, evident from the striated pits that sparsely cover the surface of most elements. As would be expected, fusion is incomplete in the sternum and proximal carpometacarpus, and the compound bones of the hindlimb have yet to fully co-ossify. The compound bones of the axial skeleton, the pygostyle and synsacrum, are fully fused in all subadult enantiornithine specimens, supporting the inference that the difference in the number of sacral vertebrae is a true distinction between *Chiappeavis* and *Pengornis*. The age of the specimen has not been explored through histology because there are no breaks in the long bones of STM 29-11 to allow non-destructive sampling of this particularly nice specimen. However, the sternum is fully fused in the subadult holotype of *Eopengornis martini* STM 24-1 and V 18632. This indicates that STM 29-11 is at a relatively earlier ontogenetic stage and strongly suggests it would have increased in size with maturity. The sternum in *Parapengornis* V 18687 is also medially unfused, despite its much smaller size (Table 1), and histology confirms that this specimen is immature, with no remodeling present to indicate the individual was mature or nearly so. *Parapengornis* sp. V 18632 has a fully fused sternum but is roughly 15% smaller than *Parapengornis* V 18687, indicating these specimens may represent different species (see phylogenetic analysis). As preserved, STM 29-11 is 20% smaller than the holotype of *Pengornis houi* V 15336, although given the relative immaturity of the former specimen, we suggest the terminal body size in *Pengornis* and *Chiappeavis* would have been similar.

### 3 Phylogenetic Analysis

We investigated the phylogenetic placement of *Chiappeavis* using a modified version of the O' Connor and Zhou (2013) dataset that includes the revised character 220 used by Wang X et al. (2014). The dataset includes five pengornithids: *Eopengornis*, *Parapengornis* IVPP V 18687 and IVPP V 18632, *Pengornis*, and *Chiappeavis* (Hu et al., 2014, 2015; O' Connor et al., 2016; Wang X et al., 2014; Zhou et al., 2008). The matrix includes a total of 63 taxa, 24 of which are referable to the Enantiornithes. Using TNT software (Goloboff et al., 2008), we conducted a heuristic search using tree-bisection reconnection (TBR), retaining the single shortest tree out of every 1,000 replications. This produced 824 most parsimonious trees with a length of 897 steps. A second round of TBR produced more than 10,000 trees of the same length. In the strict consensus, all nodes collapsed except Aves itself. Investigation of the MPTs revealed that in 78% of the trees, Pengornithidae was resolved. The clade formed by *Jeholornis* + all more derived birds collapses due to the basal position of the fragmentary taxon *Chaoyangia* resolved by a small percentage of MPTs. In order to reduce the effects of homoplasy (Goloboff et al., 2008), which strongly characterizes early avian evolution (Brusatte et al., 2015), we reran the analysis with the same parameters using implied weighting ( $k = 1.0, 2.0, \text{ and } 3.0$ ) (Goloboff, 1993). With a  $k$  value of 1.0, the analysis produced 24 trees (TBR = 129.47). The strict consensus is well resolved, with pengornithids forming successive outgroups to all other enantiornithines. With higher  $k$  values (2.0, 3.0), Pengornithidae is resolved as a clade that is sister taxon to all other enantiornithines (Fig. 7 [Figure 7: see original paper]).

---

### 4 Discussion

**Pengornithid diversity.** Five pengornithid specimens are now recognized, representing at least four distinct genera (*Chiappeavis*, *Eopengornis*, *Parapengornis*, and *Pengornis*). Pengornithids are unusual birds, differing from other enantiornithines in the morphology of the sternum (e.g., ossifying from a pair of medially fused bilateral plates and having only a single pair of caudal trabeculae), pygostyle (shorter, wider, with proximally restricted ventrolateral processes, often having a midline invagination on the caudal margin), and ischium (slender without a dorsal process). In addition, they possess features more typical of long bony-tailed birds or basal pygostylians, such as the presence of a metatarsal V and an elongate fibula. This marked increase in homoplasy caused by the inclusion of five pengornithids has thus resulted in a collapse in weakly resolved relationships from previous analyses, making a strong case for the use of implied weighting. Without the use of implied weighting, Pengornithidae is only resolved in 78% of the MPTs, but in 98% of all MPTs, V 18632 and *Eopengornis* form a clade, thus not supporting previous inferences that V 18632 should be referred to *Parapengornis* (Hu et al., 2015). This is further supported by differences in body size between V 18632 and *Parapengornis* V

18687 (see Ontogenetic status). Differences in size, morphology (Fig. 5), and stratigraphic level also suggest V 18632 is not referable to *Eopengornis martini*. Because the specimen is a subadult, Hu et al. (2014) originally refrained from naming V 18632 (*Pengornis* sp.), later referring it to *Parapengornis* (Hu et al., 2015). Here we regard this specimen as Pengornithidae indeterminate. Notably, *Eopengornis* and *Parapengornis* share a common tail morphology consisting of a pair of elongate, fully pennaceous rachis-dominated feathers but are not found to be closely related; instead, *Eopengornis* is found to be more closely related to *Chiappeavis*, despite their disparate tail morphologies. This may suggest that this analysis does not accurately portray pengornithid relationships or that pengornithid plumage was as evolutionarily labile as in extant avian families (Gluckman, 2014). In favor of the former, the pygostyle of *Chiappeavis* is most similar to that of *Pengornis* with regards to proportions, whereas that of *Parapengornis* and *Eopengornis* are slightly shorter, suggesting that *Pengornis* may have had a rectricial fan. This subtle interclade diversity is not captured by the abstract morphologies encapsulated by this and other current character matrices used for the phylogenetic analysis of Mesozoic birds.

Large body size (Table 1) and stratigraphic level (Jiufotang) suggest that *Chiappeavis* is more closely related to *Pengornis* than to *Parapengornis* (smaller) or *Eopengornis* (smaller, Huajiying), which is further supported by the morphology of the furcula (furcular rami curved in *Chiappeavis* and *Pengornis*, whereas they are straight in *Parapengornis*) (Fig. 5). The sternum in *Chiappeavis* is proportionately more elongate (length greater than width) than in *Parapengornis* and *Eopengornis* (length and width subequal). It further differs from smaller pengornithids in the morphology of the xiphial region of the sternum: the lateral margins of median trabeculae are concave in the new taxon, whereas they are straight in *Eopengornis* and *Parapengornis*. Ontogenetic and preservational differences obfuscate comparison with *Pengornis* V 15336. Postcranially, *Pengornis* V 15336 and *Chiappeavis* STM 29-11 are similar, but STM 29-11 has a greater number of sacral vertebrae despite its immaturity (although the cranial margin of the synsacrum is obscured, clearly only seven sacrals are present in *Pengornis* V 15336), and the proximal articular surface of the tibia is inclined. The increased number of sacral vertebrae and expanded premaxillae suggest *Chiappeavis* is more advanced than other pengornithids.

**Tail-fan performance in Early Cretaceous birds.** Even in the Jehol Biota, at the earliest known stage of pygostylian evolution with data limited by fossilization, there exist observable differences in the shape and relative size of the rectricial fan between clades and individual taxa. Compared to all Jehol ornithuromorphs preserving tail fans (*Yixianornis*, *Yanornis*, *Hongshanornis*, *Schizoura*, *Piscivoravis*) (Chiappe et al., 2014; Clarke et al., 2006) (Fig. 8 [Figure 8: see original paper]), the tail is proportionately shorter relative to body length in *Chiappeavis* STM 29-11 (Table 2). These taxa also differ in the degree of gradation (measured as the difference in length between the longest and shortest rectrices). Subtle differences in tail shape and length can be used to understand the selective pressures responsible for producing each phenotype.

Lift is determined by the maximum continuous width; therefore, any elongation beyond this point is considered the product of sexual selection, being not optimized for flight (Thomas and Balmford, 1995). In one test area, 80% of all bird species were found to have tail displays of some kind (Fitzpatrick, 1997). Although two tails of equal width have the same lift, longer tails generate greater moments for turning; the trade-off is that longer tails require greater muscle force and incur more drag (Thomas and Balmford, 1995). Thus, the huge diversity of avian tail morphologies represents the product of these and the numerous other selective forces that exist in the varied ecologies and lifestyles occupied by birds. The graded morphology in *Sapeornis* and even the round morphology in ornithuromorphs indicate that, although capable of generating lift, these tails were not optimized for aerodynamic function, being also shaped by sexual/signal selection. The difference between the longest and shortest rectrices is highest in *Sapeornis* and *Chiappeavis*, suggesting this influence was strongest in these taxa. Shorter tails tend to be adapted for higher flight speeds and greater lift-to-drag ratios, whereas longer tails increase maneuverability and are found in woodland birds where the cluttered environment selects for increased maneuverability and high-speed flight is infrequent (Thomas, 1997; Thomas and Balmford, 1995). Thus, the presence of long round-tails in most ornithuromorphs indicates these taxa were well adapted for the forested Jehol paleoenvironment (Zhou, 2004).

Since longer tails require greater muscle force, the proportionately shorter tail in *Chiappeavis* supports inferences based on pygostyle morphology that rectricial bulbs, if present, were poorly developed in enantiornithines. Some extant birds have a pygostyle similar to that of enantiornithines, in which the dorsal surface is expanded (having a dorsal platform and thus no dorsally blade-like pygostyle lamina) to provide additional surface area for the attachment of enlarged caudal levator muscles, yet these taxa retain rectricial bulbs (Wang and O' Connor, in press). Thus, the absence of a pygostyle lamina does not exclude enantiornithines from having this structure.

In order to more directly quantify aerodynamic differences in tail shape, we reconstructed the rectricial fan for several taxa based on the specimens most clearly preserving this feature (Table 2) and estimated their lift (Thomas, 1993). We recognize that the preserved width is not necessarily the optimal or maximal spread of the tail in flight, but like living birds, we assume a range of widths were possible at least in the Ornithuromorpha. We assume equal post-mortem compression of the rectricial bulbs (or comparable tail musculature in *Chiappeavis* and *Sapeornis*), resulting in a comparable degree of tail spread between Jehol specimens. Specimens preserved in lateral view were considered unsuitable for analysis. For Jehol birds, these measurements represent conservative estimates of the tail's lift capabilities. The tail fan of *Chiappeavis* is estimated to have the least efficient tail shape, whereas the tails in most Jehol ornithuromorphs have almost double the lifting power for their given body mass (Table 2). These measurements support our predictions based on the shape of the pygostyle and associated rectricial fan. The limited function of the enantiornithine tail fan, quantified here for the first time, provides support for the

hypothesis that this is the cause of the restricted distribution of this feature in the Enantiornithes. Paired with limited musculature (as inferred from pygostyle morphology) and the potential absence of the ability to control the spread of the tail fan, the enantiornithine aerodynamic rectricial fan may not have provided a great enough reproductive advantage to be retained through natural selection. This is at odds with derived skeletal features present in *Chiappeavis* (enlarged premaxilla, longer synsacrum), which may alternatively suggest that a tail fan was independently evolved in the *Chiappeavis* lineage.

---

## References

- Brusatte SL, O' Connor JK, Jarvis ED, 2015. The origin and diversification of birds. *Curr Biol*, 25: R888-R898
- Chiappe LM, Walker CA, 2002. Skeletal morphology and systematics of the Cretaceous Euenantiornithes (Ornithothoraces: Enantiornithes). In: Chiappe LM, Witmer LM eds. *Mesozoic Birds: Above the Heads of Dinosaurs*. Berkeley: University of California Press. 240-267
- Chiappe LM, Norell M, Clark J, 2001. A new skull of *Gobipteryx minuta* (Aves: Enantiornithes) from the Cretaceous of the Gobi Desert. *Am Mus Novit*, 3346: 1-15
- Chiappe LM, Zhao B, O' Connor JK et al., 2014. A new specimen of the Early Cretaceous bird *Hongshanornis longicresta*: insights into the aerodynamics and diet of a basal ornithuromorph. *PeerJ*, 2(e234): 1-28
- Clarke JA, Zhou ZH, Zhang FC, 2006. Insight into the evolution of avian flight from a new clade of Early Cretaceous ornithurines from China and the morphology of *Yixianornis grabaui*. *J Anat*, 208(3): 287-308
- Fitzpatrick S, 1997. Patterns of morphometric variation in birds' tails: length, shape and variability. *Biol J Linn Soc*, 62: 145-162
- Gatesy SM, Dial KP, 1996. From frond to fan: *Archaeopteryx* and the evolution of short-tailed birds. *Evolution*, 50(5): 2037-2048
- Gluckman TL, 2014. Pathways to elaboration of sexual dimorphism in bird plumage patterns. *Biol J Linn Soc*, 111: 262-273
- Goloboff PA, 1993. Estimating character weights during tree search. *Cladistics*, 9: 83-91
- Goloboff PA, Farris JS, Nixon KC, 2008. TNT, a free program for phylogenetic analysis. *Cladistics*, 24: 774-786
- Hu H, Zhou ZH, O' Connor JK, 2014. A subadult specimen of *Pengornis* and character evolution in Enantiornithes. *Vert Palasiat*, 52(1): 77-97
- Hu H, O' Connor JK, Zhou ZH, 2015. A new species of Pengornithidae (Aves: Enantiornithes) from the Lower Cretaceous of China suggests a specialized scansorial habitat previously unknown in early birds. *PLoS ONE*, 10(6): e0126791
- Ji SA, Atterholt JA, O' Connor JK et al., 2011. A new, three-dimensionally preserved enantiornithian (Aves: Ornithothoraces) from Gansu Province, north-western China. *Zool J Linn Soc*, 162: 201-219

- Liu D, Zhou ZH, Zhang YG, 2012. Mass estimate and evolutionary trend in Chinese Mesozoic fossil birds. *Vert Palasiat*, 50(1): 39-52
- O' Connor JK, 2009. A systematic review of Enantiornithes (Aves: Ornithothoraces). Ph.D thesis. Los Angeles: University of Southern California. 1-600
- O' Connor JK, Zhou ZH, 2013. A redescription of *Chaoyangia beishanensis* (Aves) and a comprehensive phylogeny of Mesozoic birds. *J Syst Palaeont*, 11(7): 889-906
- O' Connor JK, Chiappe LM, Bell A, 2011. Pre-modern birds: avian divergences in the Mesozoic. In: Dyke GD, Kaiser G eds. *Living Dinosaurs: the Evolutionary History of Birds*. New Jersey: J. Wiley & Sons. 39-114
- O' Connor JK, Wang XL, Sullivan C et al., 2013. The unique caudal plumage of *Jeholornis* and complex tail evolution in early birds. *Proc Natl Acad Sci USA*, 110(43): 17404-17408
- O' Connor JK, Wang XL, Zheng XT et al., 2016. An enantiornithine with a fan-shaped tail, and the evolution of the rectricial complex in early birds. *Curr Biol*, 26: 114-119
- Pan YH, Sha JG, Zhou ZH et al., 2013. The Jehol Biota: definition and distribution of exceptionally preserved relicts of a continental Early Cretaceous ecosystem. *Cretaceous Res*, 44: 30-38
- Thomas ALR, 1993. On the aerodynamics of birds' tails. *Philos Trans R Soc Lond B Biol Sci*, 340: 361-380
- Thomas ALR, 1997. On the tails of birds. *Bioscience*, 47(4): 215-225
- Thomas ALR, Balmford A, 1995. How natural selection shapes bird's tails. *Am Nat*, 146(6): 848-868
- Wang M, O' Connor JK, Zelenkov NZ et al., 2014. A new diverse enantiornithine family (Bohaiornithidae fam. nov.) from the Lower Cretaceous of China with information from two new species. *Vert Palasiat*, 52(1): 31-76
- Wang W, O' Connor JK, in press. Morphological co-evolution of the pygostyle and tail feathers in Mesozoic birds. *Vert Palasiat*
- Wang XL, O' Connor JK, Zheng XT et al., 2014. Insights into the evolution of rachis-dominated tail feathers from a new basal enantiornithine (Aves: Ornithothoraces). *Biol J Linn Soc*, 113(3): 805-819
- Wellnhofer P, 2008. *Archaeopteryx. Der Urvogel von Solnhofen*. München: Friedrich Pfeil. 1-256
- Witmer LM, Martin LD, 1987. The primitive features of the avian palate, with special reference to Mesozoic birds. In: Mourer-Chauviré C ed. *L' Evolution des Oiseaux d' apres le Temoignage des Fossiles*. Lyon: Département des Sciences de la Terre, Université Claude-Bernard. 21-40
- Zheng XT, Zhou ZH, Wang XL et al., 2013. Hind wings in basal birds and the evolution of leg feathers. *Science*, 339: 1309-1312
- Zhou ZH, 2004. Vertebrate radiations of the Jehol Biota and their environmental background. *Chin Sci Bull*, 49(8): 754-756
- Zhou ZH, Zhang FC, 2006. Mesozoic birds of China—a synoptic review. *Vert Palasiat*, 44(1): 74-98
- Zhou ZH, Clarke J, Zhang F, 2008. Insight into diversity, body size and morphological evolution from the largest Early Cretaceous enantiornithine bird. *J*

## Chinese Abstract

### Morphological Description of *Chiappeavis magnapremaxillo* (Pengornithidae: Enantiornithes) and Comparison of Aerodynamic Function of Tail Feathers in Early Cretaceous Birds

*Chiappeavis* is the first enantiornithine discovered to preserve a fan-shaped tail, indicating that the rectricial bulb structure was already developed in relatively primitive enantiornithines. The skeletal morphology of the holotype specimen of *Chiappeavis magnapremaxillo* is described in detail. The palatal morphology of *Chiappeavis* is similar to that of *Archaeopteryx* but differs from the Late Cretaceous enantiornithine *Gobipteryx*. Even if rectricial bulbs were present, the pygostyle morphology of pengornithids suggests that this structure was poorly developed. We estimated the lift generated by the fan-shaped tail feathers in *Chiappeavis* and compared it with other Early Cretaceous birds. The results show that the fan-shaped tail feathers of *Chiappeavis* generated less aerodynamic lift than those of sympatric ornithuromorphs, which may explain the widespread absence of aerodynamically functional tail morphologies in Enantiornithes.

**Keywords:** Mesozoic, Jehol Biota, Aves, rectrix

*Note: Figure translations are in progress. See original paper for figures.*

*Source: ChinaXiv – Machine translation. Verify with original.*