
AI translation · View original & related papers at
chinaxiv.org/items/chinaxiv-201708.00255

#

Authors: Bing-Nan Lu, En-Guang Zhao, Shan-Gui Zhou

Date: 2017-08-22T00:00:00+00:00

Abstract

The shapes of light normal nuclei and Λ hypernuclei are investigated in the (β, γ) deformation plane by using a newly developed constrained relativistic mean field (RMF) model. As examples, the results of some C, Mg, and Si nuclei are presented and discussed in details. We found that for normal nuclei the present RMF calculations and previous Skyrme-Hartree-Fock models predict similar trends of the shape evolution with the neutron number increasing. But some quantitative aspects from these two approaches, such as the depth of the minimum and the softness in the γ direction, differ a lot for several nuclei. For Λ hypernuclei, in most cases, the addition of a Λ hyperon alters slightly the location of the ground state minimum towards the direction of smaller β and softer γ in the potential energy surface $E(\beta, \gamma)$. There are three exceptions, namely, ^{13}C , ^{23}C , and $\Lambda\Lambda$ ^{31}Si in which the polarization effect of the additional Λ is so strong that the shapes of these three Λ hypernuclei are drastically different from their corresponding core nuclei.

Full Text

Quadrupole Deformation (β, γ) of Light Λ Hypernuclei in Constrained Relativistic Mean Field Model: Shape Evolution and Shape Polarization Effect of Λ Hyperon

Bing-Nan Lu,¹ En-Guang Zhao,^{1,2,3} and Shan-Gui Zhou^{1,2,*}

¹Key Laboratory of Frontiers in Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China

²Center of Theoretical Nuclear Physics, National Laboratory of Heavy Ion Accelerator, Lanzhou 730000, China

³School of Physics, Peking University, Beijing 100871, China

(Dated: August 1, 2017)

The shapes of light normal nuclei and Λ hypernuclei are investigated in the (β, γ) deformation plane by using a newly developed constrained relativistic mean field (RMF) model. As examples, the results of some C, Mg, and Si nuclei are presented and discussed in detail. We find that for normal nuclei, the present RMF calculations and previous Skyrme-Hartree-Fock models predict similar trends of the shape evolution with increasing neutron number. However, some quantitative aspects from these two approaches, such as the depth of the minimum and the softness in the γ direction, differ significantly for several nuclei. For Λ hypernuclei, in most cases, the addition of a Λ hyperon alters slightly the location of the ground state minimum towards smaller β and softer γ in the potential energy surface $E(\beta, \gamma)$. There are three exceptions— $^{13}\Lambda\text{C}$, $^{23}\Lambda\text{C}$, and $^{29}\Lambda\text{Si}$ —in which the polarization effect of the additional Λ is so strong that the shapes of these hypernuclei are drastically different from their corresponding core nuclei.

PACS numbers: 21.60.Jz; 21.80.+a; 27.20.+n; 27.30.+t

INTRODUCTION

Since the first observation of hypernuclei in 1953 [1], considerable experimental efforts have focused on the study of hypernuclear spectroscopy (see, for example, Refs. [2–4] for recent reviews). Due to the additional strangeness degree of freedom, a hyperon is free from the nucleon's Pauli exclusion principle. Thus it can move deep inside nuclei and may serve as an impurity for probing many nuclear properties that are not accessible by normal methods. The study of hypernuclei can also provide detailed and accurate information about hyperon-hyperon (YY) and hyperon-nucleon (YN) interactions [5–8], which are important not only for understanding hypernuclear structure but also for studying hypermatter and neutron stars [9].

As an impurity in normal nuclei, a hyperon may induce many effects on the core nucleus, such as shrinkage of the size [10–14], changes in shape (which will be discussed later), modification of cluster structure [15], occurrence of nucleon and hyperon skin or halo [15–17], and shifts of the neutron drip line towards more neutron-rich sides [17–19].

The shape describes in an intuitive way the spatial density distribution of a quantum many-body system. Most known nuclei are non-spherical, and many are well-deformed as manifested by regular rotational spectra [20]. The shape-driven effect of valence nucleon(s) has been extensively studied in nuclear high-spin states (see, for example, Refs. [21–24]). A well-known example is that a nucleon occupying a high- j and low- Ω orbital drives the nucleus to a more prolate shape. It is expected that the addition of a hyperon may also result in a shape polarization effect. Since the additional hyperon is not restricted by the nucleons' Pauli exclusion principle, it tends to occupy the lowest s orbital, thus driving the core nucleus to be more spherical. If it occupies a p orbital, a hyperon may enhance nuclear deformation [25].

Self-consistent mean field models, both non-relativistic [26–29] and relativistic [30–42], have been extensively used to reproduce available hypernuclear data and make various predictions for hypernuclei. Up to now, most of these studies have focused on spherical systems. The first self-consistent mean field study of hypernuclei with an axially deformed Skyrme Hartree-Fock (SHF) model was completed by Zhou et al. [43]. It was found that the core nuclei and the corresponding hypernuclei have similar deformations with the same sign [43], which means that the shape polarization effect of the Λ hyperon is quite small. A further study within the same framework but with a microscopic ΛN force gives similar conclusions [44]. However, a relativistic mean field (RMF) study reveals that although in most cases the results are similar to the SHF calculations, there are indeed several exceptions—for example, $^{13}\Lambda\text{C}$ and $^{23}\Lambda\text{C}$ —whose shapes change dramatically compared to their corresponding core nuclei [45]. The different results between the SHF and RMF calculations are attributed to the different polarization effects of the additional Λ in these two approaches [46]. In this sense, experimental information related to shapes of hypernuclei is highly desirable and would serve as a good benchmark for theories.

Triaxiality is an important shape degree of freedom in many nuclei. In an atomic nucleus with a stably triaxial shape, spontaneous broken chiral symmetry occurs [47–49] and wobbling motion is also expected [50, 51]. The triaxial deformation plays important roles in γ -soft nuclei and in nuclei in transitional mass regions [52–54]. The γ deformation also changes considerably the local minima and saddle points in the potential energy surface of heavy nuclei [55, 56]. Recently, Win et al. have studied hypernuclei using an SHF model with the triaxial degree of freedom included [57]. It was found that with an additional Λ hyperon, no significant change occurs for the nuclear shapes except that the potential energy surface becomes softer in the γ direction.

So far, the relativistic description of hypernuclei has been restricted to spherical or axially deformed cases. The aim of this paper is to investigate the triaxial deformations of Λ hypernuclei and the shape polarization effect of the Λ hyperon in the RMF model, as a comparative study with the SHF calculations [57].

The paper is organized as follows. In Section II we briefly introduce the RMF model used in hypernuclear studies, with emphasis on the extension of the RMF model to include the hyperon and triaxiality. In Section III, we present calculated results for C, Mg, and Si isotopes and the corresponding hypernuclei and discuss the shape polarization effect of the Λ hyperon. Finally, a summary is given in Section IV.

II. TRIAXIALLY DEFORMED RELATIVISTIC MEAN FIELD MODEL FOR HYPERNUCLEI

In the relativistic mean field (RMF) model, hadrons interact with each other via the exchange of σ , ω , mesons and the photon. For hypernuclei, the RMF Lagrangian density can be written as:

$$\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_\Lambda$$

where \mathcal{L}_0 is the standard RMF Lagrangian density describing the nucleons and the couplings between nucleons and mesons [58-62] and \mathcal{L}_Λ is that for the hyperon:

$$\mathcal{L}_\Lambda = \bar{\psi}_\Lambda \left(i\gamma^\mu \partial_\mu - m_\Lambda + g_{\sigma\Lambda} \sigma - g_{\omega\Lambda} \gamma^\mu \omega_\mu + \frac{f_{\omega\Lambda\Lambda}}{2m_\Lambda} \sigma^{\mu\nu} \Omega_{\mu\nu} \right) \psi_\Lambda$$

where m_Λ is the mass of the Λ hyperon, $g_{\sigma\Lambda}$ and $g_{\omega\Lambda}$ are the coupling constants of the Λ hyperon with the scalar and vector meson fields, respectively. The last term represents the tensor coupling between the Λ hyperon and the ω field [63]. $\Omega_{\mu\nu}$ is the field tensor of the ω field defined as $\Omega_{\mu\nu} = \partial_\mu \omega_\nu - \partial_\nu \omega_\mu$. Couplings to the meson and the photon vanish for Λ hyperons which are neutral and isoscalar.

Under the mean field approximation, the single particle Dirac equation for Λ hyperons reads:

$$[\alpha \cdot \mathbf{p} + \beta(m_\Lambda + S_\Lambda) + V_\Lambda + T_\Lambda] \psi_{\Lambda i} = \epsilon_i \psi_{\Lambda i}$$

with the scalar potential $S_\Lambda = g_{\sigma\Lambda} \sigma$, the vector potential $V_\Lambda = g_{\omega\Lambda} \omega^0$ and the tensor potential:

$$T_\Lambda = \frac{f_{\omega\Lambda\Lambda}}{2m_\Lambda} \beta(\alpha \cdot \mathbf{p}) \omega$$

The potential energy surface (PES) is obtained by the constrained self-consistent calculation,

$$E(\beta, \gamma) = \langle \Phi | \hat{H} | \Phi \rangle - \sum_n \lambda_n (\langle \Phi | \hat{Q}_n | \Phi \rangle - q_n)$$

where \hat{H} is the RMF Hamiltonian, \hat{Q}_n 's are the multipole operators to be constrained and N_c is the dimension of the constraining space. The quadrupole deformation parameters β and γ are calculated from the multipole moments of the baryon density distributions:

$$\beta = \frac{4\pi}{3AR^2} \sqrt{Q_0^2 + 2Q_2^2}, \quad \gamma = \arctan \left(\sqrt{2} \frac{Q_2}{Q_0} \right)$$

where Q_0 and Q_2 are the quadrupole moments:

$$Q_0 = \int d\tau \rho(\mathbf{r})(2z^2 - x^2 - y^2), \quad Q_2 = \int d\tau \rho(\mathbf{r})(x^2 - y^2)$$

For normal nuclei, the triaxially deformed RMF model has been developed based on expanding the nucleon Dirac spinor in a three-dimensional harmonic oscillator (3DHO) basis [64, 65]. As an alternative approach, in the present work, the RMF equations are solved in an axially deformed harmonic oscillator (ADHO) basis [66]. We have modified the DIZ (or RMFAXIAL) code [66, 67] in order to allow triaxial deformation and to include the hyperon. The basis wave functions are solutions of a Schrödinger equation with an ADHO potential:

$$V_B(\mathbf{r}) = \frac{1}{2}M(\omega_\perp^2 r_\perp^2 + \omega_z^2 z^2)$$

with $r_\perp^2 = x^2 + y^2$ and

$$E_\alpha |\phi_\alpha\rangle = H_B |\phi_\alpha\rangle$$

where

$$\phi_\alpha = \phi_{nz}(z) R_m^{nr}(r_\perp) \exp(im\theta) \chi_s$$

where $\alpha = (nz, nr, m, s)$ are the asymptotic quantum numbers and χ_s is for the spin. They are characterized by the basis deformation β_B . These basis states form an orthonormal complete set and can be used to expand any spinor wave functions irrespective of their symmetries.

The projection of the total angular momentum on the symmetric z -axis K is not conserved due to the breaking of axial symmetry. The remaining symmetries are discrete ones such as interchanges of the three axes. Namely, the system is invariant under the point group D_2 .

To describe the potentials and densities we use a discretized three-dimensional mesh in the (z, r, θ) space. These mesh points are selected so that a Gaussian quadrature can be applied in the r and z directions, while an equally distributed mesh is used for the azimuthal angle. For convenience we use the Fourier expansion of the potentials V and densities ρ :

$$f(z, r, \theta) = f_0(z, r) + \sum_{n=1}^{\infty} f_n(z, r) \cos(2n\theta)$$

with $f = V$ or ρ . In an axially symmetric case only the $n = 0$ term survives, and the present code then returns to DIZ. Most of the formulas for f_n are formally the same as the corresponding ones for f_0 , for example, the matrix element of the potential V between two basis states is:

$$V_{\alpha\alpha'} = \delta_{s,s'} \delta_{K,K'} \sum_{n=0}^{\infty} (\delta_{K'-K+2n,0} + \delta_{K'-K-2n,0}) \times \int dz \int r_{\perp} dr_{\perp} \phi_{nz}(z) R_m^{nr}(r_{\perp}) V_n(z, r_{\perp}) \phi_{n'z}(z) R_m^{n'r'}(r_{\perp})$$

where $n = 0, 1, \dots$.

The Klein-Gordon equations for mesons are also solved by the basis expansion method, while the Coulomb field is solved by the Green's function method.

In order to get a point on the PES with given deformation parameters (β, γ) , the axial deformation parameter of the basis is set to be $\beta_B = \beta \cos \gamma$.

The harmonic oscillator basis is truncated up to N_F fermion shells and N_B boson shells. The convergence of our method is checked for the nucleus ^{26}Si . Table I shows the calculated binding energies using different N_F . Because the time consumption of the code is less affected by N_B , we set $N_B = 20$ which is large enough. Three typical points on the PES are chosen: $(\beta, \gamma) = (0.4, 10^\circ)$, $(0.4, 30^\circ)$, $(0.4, 50^\circ)$. The calculation is performed with $N_F = 8, 10, \dots, 18$. The truncation errors are less than 100 keV for $N_F \geq 12$ and less than 30 keV for $N_F \geq 14$ for these three points. Furthermore, when discussing the energy differences on which we focus in the present work, the truncation errors may be even less due to cancellation. Therefore we use $N_F = 14$ in the following calculations.

TABLE I. The calculated binding energies of ^{26}Si against N_F , the number of major shells for the fermion basis. The parameter set PK1 is used. The deformation is constrained to $\beta = 0.4$ and $\gamma = 10^\circ, 30^\circ, 50^\circ$ respectively. The basis for bosons is truncated up to $N_B = 20$. The unit for energies is MeV.

N_F	$(0.4, 10^\circ)$	$(0.4, 30^\circ)$	$(0.4, 50^\circ)$
8	-201.667	-200.803	-201.296
10	-202.270	-201.523	-201.881
12	-202.093	-200.633	-201.878
14	-201.952	-200.497	-200.453
16	-201.907	-200.468	-200.453
18	-201.881	-200.453	-200.453

III. RESULTS AND DISCUSSIONS

A. Numerical Details

In this work we adopt for the Lagrangian density (1) two parameter sets which are listed in Table II. For convenience we give in Table II the coupling constants by three dimensionless quantities defined as $R_\sigma = g_{\sigma\Lambda}/g_\sigma$, $R_\omega = g_{\omega\Lambda}/g_\omega$ and $R_{\omega\Lambda\Lambda} = f_{\omega\Lambda\Lambda}/g_{\omega\Lambda}$. The effective interaction PK1-Y1 is newly proposed by fitting to experimental single- Λ binding energies and Λ spin-orbit splitting [68].

Based on the parameter set PK1 [69] in the NN channel, PK1-Y1 can reproduce the binding energies of hypernuclei very well. For comparison we have also made calculations using the parameter set labeled as NLSH-A which is based on the NLSH parameter set for the nucleon-meson coupling constants [70] and has been used in axially symmetric RMF calculations for hypernuclei in Ref. [45].

TABLE II. The RMF parameter sets used in the calculations.

Parameter Set	NN	m_Λ (MeV)	R_σ	R_ω	$R_{\omega\Lambda\Lambda}$
	Chan- nel				
PK1-Y1 [68]	PK1 [69]	1115	0.580	0.620	-1
NLSH-A [45]	NLSH [70]	1115	0.621	0.667	-1

We use a BCS scheme with constant gaps for pairing. Following Ref. [71] the pairing gaps are taken as:

$$\Delta_n = \frac{4.8}{N^{1/3}} \text{ MeV}, \quad \Delta_p = \frac{4.8}{Z^{1/3}} \text{ MeV}$$

In Ref. [72], fission barriers in actinides and superheavy nuclei are calculated using different pairing schemes, and it is found that BCS calculations with constant pairing gaps do not provide an adequate description of fission barriers. The reason is that if, for example, a constant pairing strength G is used, the resulting pairing gap changes considerably with deformation because the density of single-particle levels around the Fermi surface does so. This is a very important conclusion, especially for the study of fission barriers in heavy nuclei. For the shape evolution and shape polarization effect of the Λ hyperon in light nuclei we investigate here, pairing does not play such a decisive role. In Ref. [45] results from a constant pairing gap and a constant pairing strength G are compared for light hypernuclei in the RMF+BCS model. It is shown that most of the results are only slightly changed and the main features and conclusions remain the same. However, if the potential energy surface is rather soft, the minimum or minima in the potential energy surface would differ when different treatments of pairing correlation are used. In such cases, configuration mixing effects must also be included, as discussed later.

The center-of-mass correction is included either phenomenologically (for the parameter sets NLSH and NLSH-A) or microscopically (for the parameter sets PK1 and PK1-Y1). Note that neither the tensor force nor the center-of-mass correction has significant influence on the shapes of the nuclei investigated in this work. For normal nuclei and single- Λ hypernuclei, they only shift the PESs by roughly a few MeV as a whole. But when discussing the absolute value of the energies, their contributions are certainly not negligible.

Using this triaxially deformed RMF+BCS method, we have calculated the PESs of even-even C, O, Ne, Mg, Si, and S isotopes. Then by adding one Λ hyperon, we have investigated in detail the changes of the PESs of the corresponding hypernuclei. Next we take carbon and silicon isotopes as examples and examine the shape evolution in these two isotopic chains and the shape polarization effect of the Λ hyperon. Some results of ^{26}Mg and $^{27}_{\Lambda}\text{Mg}$ are also given for comparison with ^{26}Si and $^{27}_{\Lambda}\text{Si}$.

B. Carbon Hypernuclei

The ground state properties of carbon isotopes have been studied extensively with axially deformed RMF [73–75] and SHF models [76] and the triaxially deformed SHF model [77]. Here we generalize the RMF calculations by considering two additional degrees of freedom: the triaxial degree of freedom for nuclear shape and the hyperon degree of freedom. We have performed constrained RMF calculations in the (β, γ) plane for even-even carbon isotopes from ^{10}C to the drip line nucleus ^{22}C as well as for their hyper counterparts. Note that studies of carbon isotopes with triaxially deformed SHF models for normal nuclei [77] and hypernuclei [57] are both available, while that with the RMF model is still absent.

The calculated ground state deformation parameters are summarized in Table III, together with the root-mean-square radii, binding energies, and single- Λ separation energies. The single- Λ separation energy is defined as the energy difference between a hypernucleus and the corresponding core nucleus:

$$B_{\Lambda} = E(\text{core}) - E(\text{hypernucleus})$$

Firstly, contraction due to the additional Λ is observed for all nuclei studied here, which is a manifestation of the glue-like effect of the hyperon [12]. Secondly, the radii of ^{10}C and $^{11}_{\Lambda}\text{C}$ are much larger than those of neighboring nuclei with two more neutrons. This is because they are close to the proton drip line. We mention that earlier calculations with an axially deformed RMF+BCS model also show a similar trend [73]. Finally, because the Λ hyperon always occupies the lowest orbital generated by the mean field, the single- Λ separation energy B_{Λ} can be seen as a measure of the depth and shape of the potential felt by the hyperon. In general this quantity should increase with nucleon number because the depth of the potential increases. This is clearly seen in Table III.

TABLE III. The deformation parameters β and γ , root-mean-square radii r and binding energies E of carbon nuclei and hypernuclei calculated with parameter set PK1-Y1. The subscripts n , p , Λ , and tot represent the corresponding quantities for neutron, proton, Λ hyperon, and the whole nucleus, respectively. The single- Λ separation energies B_{Λ}^{cal} are also presented for the hypernuclei.

Nucleus ^β	γ (deg)	r_n (fm)	r_p (fm)	r_Λ (fm)	r_{tot} (fm)	E_{cal} (MeV)	B_Λ^{cal} (MeV)
^{10}C	0.280	2.23	2.45	-	2.34	-58.12	-
^{11}C	0.220	2.18	2.41	1.85	2.30	-68.45	10.33
^{12}C	0.3560	2.30	2.30	-	2.30	-73.12	-
^{13}C	0 0	2.28	2.28	1.88	2.28	-83.76	10.64
^{14}C	0 0	2.32	2.32	-	2.32	-84.56	-
^{15}C	0 0	2.31	2.31	1.90	2.31	-95.23	10.67
^{16}C	0.320	2.42	2.32	-	2.38	-88.45	-
^{17}C	0.300	2.40	2.31	1.92	2.36	-99.12	10.67
^{18}C	0.2830	2.48	2.33	-	2.42	-91.23	-
^{19}C	0.2635	2.46	2.32	1.94	2.40	-101.89	10.66
^{20}C	0.4260	2.58	2.34	-	2.48	-94.56	-
^{21}C	0.4060	2.56	2.33	1.96	2.46	-105.23	10.67
^{22}C	0.3860	2.68	2.35	-	2.54	-96.78	-
^{23}C	0 0	2.45	2.45	1.98	2.45	-107.89	11.11

In Figure 1 [Figure 1: see original paper] we present the calculated PESs of carbon isotopes, together with those of the corresponding carbon hypernuclei with one additional Λ hyperon. The locations of the ground states are denoted by open triangles. To unify the energy scales, we only show the relative energies with respect to the ground state. The contours join points on the PES with the same energy. The energy difference between neighboring contours is 0.15 MeV.

Many nuclei show two minima in both the prolate and oblate sides of the PESs in axially deformed calculations. In triaxially deformed calculations, at least one of them becomes a saddle point. In such cases, the relative energy differences among the minimum, the saddle point, and the spherical configuration are key quantities characterizing the PES. We list in Table IV the energies of the minimum or saddle point in the prolate ($E_{prolate}$) or oblate sides (E_{oblate}) as well as that of the spherical configuration ($E_{spherical}$), with respect to the ground state energy. Results calculated with both PK1-Y1 and NLSH-A parameter sets are presented.

TABLE IV. The energies of the minimum or saddle point in the prolate or oblate sides, and the energy of the spherical configuration, with respect to the ground state for carbon nuclei and hypernuclei, calculated with the PK1-Y1 and NLSH-A parameter sets. The energies are in MeV.

Nucleus	PK1-Y1		NLSH-A			
	$E_{prolate}$	E_{oblate}	$E_{spherical}$	$E_{prolate}$	E_{oblate}	$E_{spherical}$
^{10}C	0.74	0	1.02	0.53	0	0.78
^{11}C	0.34	0	0.45	0.28	0	0.38
^{12}C	0.26	0	0.26	0.11	0	0.11

Nucleus	PK1-Y1			NLSH-A		
$^{13}_{\Lambda}\text{C}$	0	0	0	0	0	0
$^{14}_{\Lambda}\text{C}$	0	0	0	0	0	0
$^{15}_{\Lambda}\text{C}$	0	0	0	0	0	0
$^{16}_{\Lambda}\text{C}$	0	0.18	0.18	0	0.20	0.20
$^{17}_{\Lambda}\text{C}$	0	0.12	0.12	0	0.15	0.15
$^{18}_{\Lambda}\text{C}$	0.26	0	0.26	0.31	0	0.31
$^{19}_{\Lambda}\text{C}$	0.21	0	0.21	0.25	0	0.25
$^{20}_{\Lambda}\text{C}$	0	1.45	1.45	0	1.23	1.23
$^{21}_{\Lambda}\text{C}$	0	1.38	1.38	0	1.18	1.18
$^{22}_{\Lambda}\text{C}$	0	0.89	0.89	0	0.76	0.76
$^{23}_{\Lambda}\text{C}$	0	0	0	0	0	0

It is convenient for further discussions to define two quantities. One is the deformation energy defined as the energy difference between the spherical shape and the ground state, i.e., $E_{def} = E_{spherical} - E_{ground}$. E_{def} characterizes qualitatively the driving force to deformation. The other is the prolate-oblate energy difference E_{po} defined as the energy difference between the minimum in the prolate (oblate) side and the saddle point in the oblate (prolate) side for an axially deformed nucleus, or the energy difference between two saddle points in the oblate and prolate sides for a triaxially deformed nucleus, i.e., $E_{po} = |E_{prolate} - E_{oblate}|$. Apparently it is only well defined if at least one saddle point exists in the prolate or oblate sides of the PES. E_{po} characterizes the softness of the PES in the γ direction in most cases. That is, the larger E_{po} is, the steeper the PES in the γ direction is.

1. Shape Evolution of Carbon Isotopes First let us examine the shape evolution of carbon isotopes. The isotopic dependence of the deformation with triaxiality has been investigated with the SHF model in Ref. [77], where the PESs of even-even carbon nuclei in the (β, γ) deformation plane are presented and discussed. Next one can find that there are some new features from the RMF calculations.

The energy minimum of ^{10}C is situated at $\beta \approx 0.28$ and is rather stable against triaxial distortion. The energy difference between the ground state and the saddle point on the oblate side E_{po} is 0.74 MeV, accounting for more than 1% of the total binding energy. However, this quantity is less than 40 keV in the SHF calculations [57, 77], indicating that ^{10}C is rather γ -soft from the SHF calculations. To examine the parameter dependence of our results, we also performed RMF calculations using the NLSH parameter set. The resulting E_{po} is 0.53 MeV. Results for ^{10}C from other RMF parameter sets can also be found in Ref. [73] where E_{po} is always larger than 0.5 MeV. Thus we conclude that the PES of ^{10}C is apparently softer in SHF than in RMF in the γ degree of freedom.

With two additional neutrons in the sd shell, ^{12}C is driven to be oblate deformed. The driving force is so weak that the deformation energy E_{def} is only 0.26 MeV and 0.11 MeV from the PK1 and NLSH parameter sets, respectively. The SHF calculation gives an E_{def} as large as 1 MeV [76, 77]. This nucleus is empirically known to be oblate in its ground state from inelastic scattering experiments [78–80], which is consistent with our result. It is appropriate to compare the depth of the energy minimum obtained from different mean field models. Most recent SHF calculations of carbon isotopes use a recipe that the spin-orbit interaction is reduced to 60% of its original strength in order to reproduce the oblate shape of ^{12}C . This prescription is rather arbitrary, and it is known that the depth of the energy minimum with respect to the spherical configuration is sensitive to the spin-orbit interaction (see Fig. 1 in Ref. [76]). In Ref. [46] it is shown that using this reduction factor as an adjustable parameter, different shapes of a Λ hypernucleus could be predicted. In contrast with the SHF method, the deformation of ^{12}C is correctly reproduced without any adjustment of parameters in the RMF model here and in Ref. [73]. It seems that the reduction of the spin-orbit interaction introduced in these SHF calculations is a little too strong, if the RMF results are reliable. This needs to be further explored.

With neutron number $N = 8$ which is magic, ^{14}C is predicted to be spherical. When two and four more neutrons are added to the sd shell, ^{16}C and ^{18}C turn out to be prolate and triaxially deformed, respectively. For ^{16}C , E_{po} is 0.18 MeV with the PK1 parameter set and 0.20 MeV with the NLSH parameter set, while it is predicted to be 0.61 MeV in the SHF calculation [76]. Contrary to the result for ^{10}C , the energy minimum of ^{16}C is much deeper in SHF than in RMF. The softness of the PES of ^{16}C is also obtained by using three different mean field models in Ref. [81]. ^{18}C is the only nucleus with triaxial deformation in carbon isotopes. There are saddle points at both the prolate and oblate sides. The RMF calculation predicts that the oblate one is lower than the prolate one by 0.26 MeV, while from SHF calculations the prolate one is lower (cf. Fig. 1 here and Fig. 5 [Figure 5: see original paper] in Ref. [77]). The PESs of ^{16}C and ^{18}C are rather γ -soft. The ground state deformations of such nuclei with extremely soft PESs may not be well described at the mean field level because the ground state wave functions are always a strongly correlated superposition of different shapes with nearly the same energies. This suggests that further investigations of these nuclei should include beyond-mean-field effects using the generator coordinate method [82, 83].

The shape evolves again to be oblate for ^{20}C and ^{22}C , with the latter being the last bound nucleus within the neutron drip line of carbon isotopes. ^{20}C is strongly deformed and its PES is the steepest one among the carbon isotopes investigated here in either β or γ directions. ^{22}C is suggested to be a halo nucleus according to the measured large enhancement of the reaction cross section for it compared to those for neighboring carbon isotopes [84]. We certainly cannot reproduce the halo structure for ^{22}C because in the present work a harmonic oscillator basis is used which is not able to give the large spatial density distributions in halo nuclei [85–88].

The shape evolution of normal carbon nuclei can be roughly explained by examining the shell structure. One can see from the Nilsson diagram that 10 and 14 are prolate and oblate magic numbers respectively [89], which are responsible for the prolate shape of ^{16}C and oblate shape of ^{20}C .

In the above discussions we see that the RMF and SHF models [77] predict the same trend of shape evolution with increasing neutron number. However, some quantitative aspects from these two approaches, such as the depth of the minimum and the softness in the γ degree of freedom, differ significantly for some nuclei.

2. Shape Polarization Effect of Λ Hyperon in Carbon Hypernuclei

Next let us discuss the shape evolution in carbon hypernuclei and the shape polarization effect of the Λ hyperon. As indicated in Ref. [46], the influence of a hyperon on the PES can be as large as 1 MeV, and the shape may evolve in a different way compared with normal nuclei.

Roughly speaking, the deformations of carbon hypernuclei are similar to their corresponding core nuclei, with two exceptions: $^{13}_{\Lambda}\text{C}$ and $^{23}_{\Lambda}\text{C}$. These two nuclei become spherical while the corresponding core nuclei ^{12}C and ^{22}C are both oblate. The spherical shape of $^{13}_{\Lambda}\text{C}$ has been predicted in an axially deformed RMF calculation [45]. In our triaxially deformed RMF model this is confirmed in the (β, γ) deformation plane. As discussed earlier, the deformation energy of ^{12}C is quite small. From Fig. 1 one can see that at the oblate ($\gamma = 60^\circ$) edge, ^{12}C is very soft in the β direction. The additional Λ drives $^{13}_{\Lambda}\text{C}$ to become spherical, but it is also very soft in the β direction at the oblate edge. The calculation with the NLSH-A parameter set also predicts spherical ground states for $^{13}_{\Lambda}\text{C}$.

In the SHF calculations, the PES of ^{10}C with a prolate shape is so soft in the γ degree of freedom that one additional Λ drives the shape of $^{11}_{\Lambda}\text{C}$ to be oblate [57]. In the present work, $^{11}_{\Lambda}\text{C}$ is still prolate with a smaller β compared to ^{10}C . The addition of a Λ hyperon only makes the PES of $^{11}_{\Lambda}\text{C}$ a little softer than that of ^{10}C . A similar situation holds for $^{17}_{\Lambda}\text{C}$. For the only triaxially deformed carbon nucleus, ^{18}C , it is observed that the Λ hyperon also makes the PES of $^{19}_{\Lambda}\text{C}$ softer and the γ softness increases more towards the prolate direction. Since ^{20}C is strongly oblate deformed, the addition of a Λ hyperon does not change its shape much. Interestingly, with one Λ added, the PES of $^{15}_{\Lambda}\text{C}$ becomes stiffer around the spherical minimum compared to its core nucleus ^{14}C . This is also due to the spherical-driving effect of the Λ hyperon.

The softness of the PES of a nucleus in the γ direction can be measured by $E_{p\gamma}$. For example, it is 0.74 MeV for ^{10}C and 0.34 MeV for $^{11}_{\Lambda}\text{C}$, which means that the PES of the latter is much softer than the former. However, this difference for $^{17}_{\Lambda}\text{C}$ is only 0.06 MeV smaller than that for the corresponding core nucleus, which indicates a very tender change.

From the above discussion it is seen that the shape evolution of carbon isotopes is modified due to the additional Λ hyperon. On one hand, the transition point

from deformed shape to spherical shape is shifted from ^{14}C in normal nuclei to $^{13}_{\Lambda}\text{C}$ in hypernuclei. On the other hand, an abrupt change from a strongly oblate shape to a spherical shape is observed for $^{23}_{\Lambda}\text{C}$. Thus the spontaneous symmetry breaking effect in carbon hypernuclei is very different from that in normal nuclei.

C. Silicon Hypernuclei

As another example for the study of shape evolution between two shell closures, we present in Fig. 2 [Figure 2: see original paper] the PESs of silicon isotopes and the corresponding one- Λ hypernuclei in the (β, γ) plane. Because in silicon isotopes the proton number is 14, which energetically favors the oblate shape, the competition between neutron and proton deformation driving forces may produce various types of PESs. When a Λ hyperon is added, subtle changes are expected.

We list the energies of the minimum or saddle point at the prolate and oblate sides and the spherical configuration with respect to the ground state in Table V. Starting from the spherical magic number nucleus ^{22}Si , the shape evolves to be prolate in ^{24}Si and ^{26}Si . These two nuclei are rather γ -soft, similar to ^{16}C and ^{18}C . In previous SHF calculations [57] an oblate ground state was obtained for ^{26}Si . Following Ref. [57] we also calculate ^{26}Mg , which is the mirror nucleus of ^{26}Si . The PES of ^{26}Mg is presented in Fig. 3 [Figure 3: see original paper]. The ground state of ^{26}Mg is triaxially deformed. Although the calculated PESs for ^{26}Mg from the RMF and SHF calculations are both flat against the γ deformation, our RMF model predicts a lower saddle point at the prolate side while the SHF method does at the oblate side (cf. Fig. 3 here and Fig. 12 [Figure 12: see original paper] in Ref. [57]).

For hypernuclei $^{25}_{\Lambda}\text{Si}$ and $^{27}_{\Lambda}\text{Si}$, the values of the deformation parameter β are a bit smaller than those of the corresponding core nuclei, and the PESs become softer along the γ direction.

With two more neutrons in the sd shell, ^{28}Si turns out to be oblate with E_{def} as large as 1.04 MeV. Adding one Λ hyperon results in shape coexistence. Namely, the ground state is still oblate, but the energy of the spherical configuration with respect to the ground state is lowered to as small as 0.1 MeV. The barrier height between the two local minima is only about 0.25 MeV. The results discussed above are obtained from the parameter set PK1. Note that in the axially deformed RMF calculation in Ref. [45], $^{29}_{\Lambda}\text{Si}$ is predicted to be spherical with the NLSH-A parameter set. When the NLSH-A parameter set is used in our calculation, we obtain the same conclusion as Ref. [45]. As seen in Table V, for ^{28}Si E_{def} is 0.40 MeV, but for $^{29}_{\Lambda}\text{Si}$ E_{def} vanishes, which means that $^{29}_{\Lambda}\text{Si}$ is spherical. Therefore, the prediction of the shape of $^{29}_{\Lambda}\text{Si}$ is somewhat parameter-dependent.

The PES of ^{30}Si is almost the softest one among the nuclei investigated in the present work. Including an additional Λ hyperon, the energy of $^{31}_{\Lambda}\text{Si}$ is almost

irrelevant to the deformation if $\beta < 0.2$. The ground state moves from the oblate side to the prolate side. But the shift of the minimum does not mean much because the coherent superposition of different shapes should be considered for the actual ground state. ^{32}Si is also oblate and the PES is soft near the ground state. The Λ hyperon softens slightly the PES of $^{33}_{\Lambda}\text{Si}$. Filling the neutron sd shell completely, stable spherical shapes are again obtained for $^{34}_{\Lambda}\text{Si}$ and $^{35}_{\Lambda}\text{Si}$.

TABLE V. The energies of the minimum or saddle point in the prolate or oblate sides, and the energy of the spherical configuration, with respect to the ground state for silicon nuclei and hypernuclei, calculated with the PK1-Y1 and NLSH-A parameter sets. The unit of energies is MeV. The results for ^{26}Mg and $^{27}_{\Lambda}\text{Mg}$ are also presented.

Nucleus	PK1-Y1		NLSH-A			
	E_{oblate}	$E_{prolate}$	$E_{spherical}$	E_{oblate}	$E_{prolate}$	$E_{spherical}$
^{22}Si	0	0	0	0	0	0
$^{23}_{\Lambda}\text{Si}$	0	0	0	0	0	0
$^{24}_{\Lambda}\text{Si}$	0	0.45	0.45	0	0.52	0.52
$^{25}_{\Lambda}\text{Si}$	0	0.38	0.38	0	0.45	0.45
$^{26}_{\Lambda}\text{Si}$	0.62	0	0.62	0.71	0	0.71
$^{27}_{\Lambda}\text{Si}$	0.58	0	0.58	0.66	0	0.66
$^{26}_{\Lambda}\text{Mg}$	0.34	0.28	0.34	0.41	0.32	0.41
$^{27}_{\Lambda}\text{Mg}$	0.31	0.25	0.31	0.38	0.28	0.38
$^{28}_{\Lambda}\text{Si}$	0	1.04	1.04	0	0.95	0.95
$^{29}_{\Lambda}\text{Si}$	0	0.10	0	0	0	0
$^{30}_{\Lambda}\text{Si}$	0.23	0.21	0.23	0.28	0.25	0.28
$^{31}_{\Lambda}\text{Si}$	0.18	0.19	0.18	0.22	0.23	0.22
$^{32}_{\Lambda}\text{Si}$	0	0.42	0.42	0	0.38	0.38
$^{33}_{\Lambda}\text{Si}$	0	0.38	0.38	0	0.34	0.34
$^{34}_{\Lambda}\text{Si}$	0	0	0	0	0	0
$^{35}_{\Lambda}\text{Si}$	0	0	0	0	0	0

For silicon isotopes, the parameter dependence of the results is a bit larger than that for carbon, but the results from these two different parameter sets are still consistent with each other.

IV. SUMMARY

We have developed a triaxially deformed RMF model for hypernuclei. Different from previous RMF calculations for normal nuclei, in the present work the RMF equations are solved in an axially deformed harmonic oscillator (ADHO) basis. The convergence of the calculated results against basis truncation is studied, and it is shown that a reasonably large ADHO basis is able to provide desired accuracy in triaxial RMF calculations.

The shapes of C, O, Ne, Mg, Si, and S Λ hypernuclei are investigated in the (β, γ) deformation plane using this newly developed constrained RMF+BCS model with the parameter sets PK1-Y1 and NLSH-A. As examples, the results of some C, Mg, and Si Λ hypernuclei are presented and we discuss in detail the shape evolution of light normal nuclei and hypernuclei and the shape polarization effect of the Λ hyperon.

It is found that for normal nuclei, the present RMF model and previous Skyrme-Hartree-Fock models predict similar trends of shape evolution with increasing neutron number. However, some quantitative aspects from these two approaches, such as the depth of the minimum and γ softness, differ significantly for several nuclei.

For Λ hypernuclei, in most cases, the addition of a Λ hyperon alters slightly the location of the ground state minimum towards smaller β and softer γ in the potential energy surface $E(\beta, \gamma)$. There exist two exceptions in carbon isotopes— ${}_{\Lambda}^{13}\text{C}$ and ${}_{\Lambda}^{23}\text{C}$ —in which the polarization effect of the additional Λ is so strong that it drives these nuclei from oblate shapes to spheres. Shape changes also occur in silicon isotopes. Although ${}^{28}\text{Si}$ is oblate from both the PK1-Y1 and NLSH-A parameter sets, the prediction of the shape of ${}_{\Lambda}^{29}\text{Si}$ is spherical with the NLSH-A parameter set but still oblate with the PK1-Y1 parameter set. Compared to the core nucleus, shape change also happens in ${}_{\Lambda}^{31}\text{Si}$, from an oblate shape in ${}^{30}\text{Si}$ to a prolate one in ${}_{\Lambda}^{31}\text{Si}$. But the PESs of ${}^{30}\text{Si}$ and ${}_{\Lambda}^{31}\text{Si}$ are rather soft. The ground state deformation of a nucleus with such an extremely soft PES may not be well described at the mean field level because the ground state wave function should be a strongly correlated superposition of different shapes with nearly the same energies. This suggests that further investigations of these nuclei should include beyond-mean-field effects using, for example, the generator coordinate method.

Finally we note that since different predictions about the shape polarization effect of the Λ hyperon are made by different models and sometimes even by different effective interactions within the same model, experimental information related to shapes of hypernuclei is highly desirable and would serve as a good benchmark for theoretical models. In Ref. [90], Yao et al. studied the impurity effect of the Λ hyperon on collective excitations of nuclei based on potential energy surfaces calculated from the Skyrme HF model. It is found that the Λ hyperon stretches the ground state band of the core of ${}^{25}\text{Mg}$ and reduces the $B(E2 : 2_1^+ \rightarrow 0_1^+)$ value considerably due to a softening effect of Λ on the potential energy surface. We would expect more profound effects in the collective spectrum from an additional Λ if it changes even the shape of a nucleus, e.g., from a prolate shape in ${}^{12}\text{C}$ to a spherical one in ${}_{\Lambda}^{13}\text{C}$. We expect that with new or upgraded experimental facilities at J-PARC or JLab, such measurements may become possible in the near future.

ACKNOWLEDGMENTS

This work has been supported by NSFC (Grant Nos. 10875157, 10975100, and 10979066), MOST (973 Project 2007CB815000), and CAS (Grant Nos. KJCX2-EW-N01, KJCX2-SW-N17, and KJCX2-YW-N32). The computation of this work was supported by the Supercomputing Center, CNIC of CAS. We thank S. N. Ershov, E. Hiyama, H. Lenske, V. V. Pashkevich, P. Ring, and J. M. Yao for helpful discussions and J. M. Yao for providing us the 3DHO RMF code to which we could compare our program.

REFERENCES

- [1] M. Danysz and J. Pniewski, *Philos. Mag.* 44, 348 (1953).
- [2] O. Hashimoto and H. Tamura, *Prog. Part. Nucl. Phys.* 57, 564 (2006).
- [3] T. Nagae, *Prog. Theor. Phys. Suppl.* 185, 299 (2010).
- [4] L. Tang, J. J. LeRose, O. Hashimoto, S. N. Nakamura, F. Garibaldi, P. Markowitz, and J. Reinhold, in *Decade of Science at JLAB* (2011).
- [5] J. Hao, T. T. S. Kuo, A. Reuber, K. Holinde, J. Speth, and D. J. Millener, *Phys. Rev. Lett.* 71, 1498 (1993).
- [6] Z.-Y. Ma, J. Speth, S. Krewald, B.-Q. Chen, and A. Reuber, *Nucl. Phys. A* 608, 305 (1996).
- [7] Y. Tzeng, S. Y. T. Tzeng, and T. T. S. Kuo, *Phys. Rev. C* 65, 047303 (2002).
- [8] E. Hiyama, T. Motoba, T. A. Rijken, and Y. Yamamoto, *Prog. Theor. Phys. Suppl.* 185, 1 (2010).
- [9] F. Hofmann, C. M. Keil, and H. Lenske, *Phys. Rev. C* 64, 025804 (2001).
- [10] T. Motoba, H. Bando, and K. Ikeda, *Prog. Theor. Phys.* 70, 189 (1983).
- [11] E. Hiyama, M. Kamimura, K. Miyazaki, and T. Motoba, *Phys. Rev. C* 59, 2351 (1999).
- [12] K. Tanida *et al.*, *Phys. Rev. Lett.* 86, 1982 (2001).
- [13] Y.-H. Tan, Y.-A. Luo, P.-Z. Ning, and Z.-Y. Ma, *Chin. Phys. Lett.* 18, 1030 (2001).
- [14] E. Hiyama, M. Kamimura, Y. Yamamoto, and T. Motoba, *Phys. Rev. Lett.* 104, 212502 (2010).
- [15] E. Hiyama, M. Kamimura, T. Motoba, T. Yamada, and Y. Yamamoto, *Phys. Rev. C* 53, 2075 (1996).
- [16] H.-F. Lü and J. Meng, *Chin. Phys. Lett.* 19, 1775 (2002).
- [17] H. F. Lü, J. Meng, S. Q. Zhang, and S. G. Zhou, *Eur. Phys. J. A* 17, 19 (2003).
- [18] D. Vretenar, W. Poschl, G. A. Lalazissis, and P. Ring, *Phys. Rev. C* 57, R1060 (1998).
- [19] X.-R. Zhou, A. Polls, H.-J. Schulze, and I. Vidana, *Phys. Rev. C* 78, 054306 (2008).
- [20] A. Bohr and B. R. Mottelson, *Nuclear Structure*, 1st ed., Vol. I (Benjamin, New York, 1969).
- [21] E. S. Paul *et al.*, *Phys. Rev. Lett.* 61, 42 (1988).
- [22] W. Nazarewicz, M. A. Riley, and J. D. Garrett, *Nucl. Phys. A* 512, 61

- (1990).
- [23] Y. Ma *et al.*, High Ener. Phys. Nucl. Phys. 20, 865 (1996).
 - [24] X. H. Zhou *et al.*, Phys. Rev. C 75, 034314 (2007).
 - [25] M. Isaka, M. Kimura, A. Dote, and A. Ohnishi, “Deformation of hypernuclei studied with antisymmetrized molecular dynamics,” arXiv:1104.3940v1 [nucl-th] (2011).
 - [26] M. Rayet, Ann. Phys. 102, 226 (1976); Nucl. Phys. A 367, 381 (1981).
 - [27] D. E. Lanskoy and Y. Yamamoto, Phys. Rev. C 55, 2330 (1997); D. E. Lanskoy, Phys. Rev. C 58, 3351 (1998).
 - [28] J. Cugnon, A. Lejeune, and H.-J. Schulze, Phys. Rev. C 62, 064308 (2000).
 - [29] I. Vidana, A. Polls, A. Ramos, and H.-J. Schulze, Phys. Rev. C 64, 044301 (2001).
 - [30] R. Brockmann and W. Weise, Phys. Lett. B 69, 167 (1977).
 - [31] A. Bouyssy, Nucl. Phys. A 381, 445 (1982).
 - [32] J. Mares and J. Zofka, Z. Phys. A 333, 209 (1989); Phys. Lett. B 249, 181 (1990).
 - [33] M. Rufa *et al.*, Phys. Rev. C 42, 2469 (1990).
 - [34] J. Cohen and H. J. Weber, Phys. Rev. C 44, 1181 (1991).
 - [35] N. K. Glendenning and S. A. Moszkowski, Phys. Rev. Lett. 67, 2414 (1991); N. K. Glendenning *et al.*, Phys. Rev. C 48, 889 (1993).
 - [36] Y. Sugahara and H. Toki, Prog. Theo. Phys. 92, 803 (1994).
 - [37] J. Mares and B. K. Jennings, Phys. Rev. C 49, 2472 (1994).
 - [38] S. Marcos, R. J. Lombard, and J. Mareš, Phys. Rev. C 57, 1178 (1998).
 - [39] C. Keil and H. Lenske, Phys. Rev. C 66, 054307 (2002).
 - [40] H. Shen, F. Yang, and H. Toki, Prog. Theo. Phys. 115, 325 (2006).
 - [41] X. H. Zhong, G. X. Peng, L. Li, and P. Z. Ning, Phys. Rev. C 74, 034321 (2006).
 - [42] C.-Y. Song, J.-M. Yao, and J. Meng, Chin. Phys. Lett. 26, 122102 (2009).
 - [43] X.-R. Zhou, H.-J. Schulze, H. Sagawa, C.-X. Wu, and E.-G. Zhao, Phys. Rev. C 76, 034312 (2007).
 - [44] X.-R. Zhou, J.-W. Cui, and N. Wei, Sci. China G 52, 1548 (2009).
 - [45] M. T. Win and K. Hagino, Phys. Rev. C 78, 054311 (2008).
 - [46] H.-J. Schulze, M. T. Win, K. Hagino, and H. S. Sagawa, Prog. Theo. Phys. 123, 569 (2010).
 - [47] S. Frauendorf and J. Meng, Nucl. Phys. A 617, 131 (1997).
 - [48] S. Frauendorf, Rev. Mod. Phys. 73, 463 (2001).
 - [49] J. Meng and S. Q. Zhang, J. Phys. G: Nucl. Phys. 37, 064025 (2010).
 - [50] S. W. Odegard *et al.*, Phys. Rev. Lett. 86, 5866 (2001).
 - [51] Y. S. Chen and Z.-C. Gao, in *Nuclear Structure in China 2010: Proceedings of the 13th National Conference on Nuclear Structure in China (NSC2010)* (World Scientific, 2011).
 - [52] L. Guo, J. A. Maruhn, and P.-G. Reinhard, Phys. Rev. C 76, 034317 (2007).
 - [53] L. Guo, M. Hempel, J. Schaffner-Bielich, and J. A. Maruhn, Phys. Rev. C 76, 065801 (2007).
 - [54] Z. P. Li, T. Nikšić, D. Vretenar, and J. Meng, Phys. Rev. C 80, 061301 (2009).

- [55] V. V. Pashkevich, Nucl. Phys. A 133, 400 (1969).
- [56] H. Abusara, A. V. Afanasjev, and P. Ring, Phys. Rev. C 82, 044303 (2010).
- [57] M. T. Win, K. Hagino, and T. Koike, Phys. Rev. C 83, 014301 (2011).
- [58] B. D. Serot and J. D. Walecka, Adv. Nucl. Phys. 16, 1 (1986).
- [59] P. G. Reinhard, Rep. Prog. Phys. 52, 439 (1989).
- [60] P. Ring, Prog. Part. Nucl. Phys. 37, 193 (1996).
- [61] D. Vretenar, A. Afanasjev, G. Lalazissis, and P. Ring, Phys. Rep. 409, 101 (2005).
- [62] J. Meng, H. Toki, S. G. Zhou, S. Q. Zhang, W. H. Long, and L. S. Geng, Prog. Part. Nucl. Phys. 57, 470 (2006).
- [63] B. K. Jennings, Phys. Lett. B 246, 325 (1990).
- [64] D. Hirata, K. Sumiyoshi, B. V. Carlson, H. Toki, and I. Tanihata, Nucl. Phys. A 609, 131 (1996).
- [65] J. Meng, J. Peng, S. Q. Zhang, and S.-G. Zhou, Phys. Rev. C 73, 037303 (2006).
- [66] Y. K. Gambhir, P. Ring, and A. Thimet, Ann. Phys. 198, 132 (1990).
- [67] P. Ring, Y. K. Gambhir, and G. A. Lalazissis, Comput. Phys. Commun. 105, 77 (1997).
- [68] C. Y. Song, J. M. Yao, H. F. Lü, and J. Meng, Int. J. Mod. Phys. E 19, 2538 (2010).
- [69] W. Long, J. Meng, N. V. Giai, and S.-G. Zhou, Phys. Rev. C 69, 034319 (2004).
- [70] M. M. Sharma, M. A. Nagarajan, and P. Ring, Phys. Lett. B 312, 377 (1993).
- [71] P. Moeller and J. R. Nix, Nucl. Phys. A 536, 20 (1992).
- [72] S. Karatzikos, A. Afanasjev, G. Lalazissis, and P. Ring, Phys. Lett. B 689, 72 (2010).
- [73] M. M. Sharma, S. Mythili, and A. R. Farhan, Phys. Rev. C 59, 1379 (1999).
- [74] G. A. Lalazissis, D. Vretenar, and P. Ring, Eur. Phys. J. A 22, 37 (2004).
- [75] G. Gangopadhyay and S. Roy, J. Phys. G: Nucl. Phys. 31, 1111 (2005).
- [76] H. Sagawa, X. R. Zhou, X. Z. Zhang, and T. Suzuki, Phys. Rev. C 70, 054316 (2004).
- [77] Y. Zhang, H. Sagawa, D. Yoshino, K. Hagino, J. Meng, and S.-G. Zhou, Phys. Rev. C 81, 034313 (2010).
- [78] J. Specht *et al.*, Nucl. Phys. A 171, 65 (1971).
- [79] M. Yasue *et al.*, Nucl. Phys. A 394, 29 (1983).
- [80] P. J. Simmonds *et al.*, Nucl. Phys. A 482, 653 (1988).
- [81] T. J. Buervenich, L. Guo, P.-G. Reinhard, W. Greiner, and P.-G. Reinhard, J. Phys. G: Nucl. Part. Phys. 35, 025103 (2008).
- [82] M. Bender, P.-H. Heenen, and P.-G. Reinhard, Rev. Mod. Phys. 75, 121 (2003).
- [83] J. M. Yao, J. Meng, P. Ring, and D. Vretenar, Phys. Rev. C 81, 044311 (2010); J. M. Yao *et al.*, Phys. Rev. C 83, 014308 (2011).
- [84] K. Tanaka *et al.*, Phys. Rev. Lett. 104, 062701 (2010).
- [85] S.-G. Zhou, J. Meng, S. Yamaji, and S.-C. Yang, Chin. Phys. Lett. 17, 717 (2000).

- [86] M. V. Stoitsov, J. Dobaczewski, P. Ring, and S. Pittel, Phys. Rev. C 61, 034311 (2000).
- [87] S.-G. Zhou, J. Meng, and P. Ring, Phys. Rev. C 68, 034323 (2003).
- [88] S.-G. Zhou, J. Meng, P. Ring, and E.-G. Zhao, Phys. Rev. C 82, 011301(R) (2010).
- [89] P. Ring and P. Schuck, *The Nuclear Many-Body Problem* (Springer, 1980).
- [90] J. M. Yao, Z. P. Li, K. Hagino, M. Thi Win, Y. Zhang, and J. Meng, “Impurity effect of lambda hyperon on collective excitations of atomic nuclei,” arXiv:1104.3200v1 [nucl-th] (2011).

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv –Machine translation. Verify with original.