

## Spin symmetry in the anti-nucleon spectrum postprint

**Authors:** Shan-Gui Zhou Jie Meng P. Ring

**Date:** 2017-08-22T00:00:00+00:00

### Abstract

We discuss spin and pseudo-spin symmetry in the spectrum of single nucleons and single anti-nucleons in a nucleus. As an example we use relativistic mean field theory to investigate single anti-nucleon spectra. We find a very well developed spin symmetry in single anti-neutron and single anti-proton spectra. The dominant components of the wave functions of the spin doublet are almost identical. This spin symmetry in anti-particle spectra and the pseudo-spin symmetry in particle spectra have the same origin. However it turns out that the spin symmetry in anti-nucleon spectra is much better developed than the pseudo-spin symmetry in normal nuclear single particle spectra.

### Full Text

---

#### Spin Symmetry in the Anti-Nucleon Spectrum

Shan-Gui Zhou<sup>1,2,3,4</sup>, Jie Meng<sup>1,3,4</sup>, and P. Ring<sup>5</sup>

<sup>1</sup>School of Physics, Peking University, Beijing 100871, China

<sup>2</sup>Max-Planck-Institut für Kernphysik, 69029 Heidelberg, Germany

<sup>3</sup>Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100080, China

<sup>4</sup>Center of Theoretical Nuclear Physics, National Laboratory of Heavy Ion Accelerator, Lanzhou 730000, China

<sup>5</sup>Physikdepartment, Technische Universität München, 85748 Garching, Germany

(Dated: August 1, 2017)

We discuss spin and pseudo-spin symmetry in the spectrum of single nucleons and single anti-nucleons in a nucleus. As an example we use relativistic mean

field theory to investigate single anti-nucleon spectra. We find a very well developed spin symmetry in single anti-neutron and single anti-proton spectra. The dominant components of the wave functions of the spin doublet are almost identical. This spin symmetry in anti-particle spectra and the pseudo-spin symmetry in particle spectra have the same origin. However it turns out that the spin symmetry in anti-nucleon spectra is much better developed than the pseudo-spin symmetry in normal nuclear single particle spectra.

PACS numbers: 21.10-k, 14.20-c, 21.10.Hw, 21.10.Pc, 21.60.Jz

Symmetries in single particle spectra of atomic nuclei have been discussed extensively in the literature, particularly the violation of spin-symmetry by the spin-orbit term and approximate pseudo-spin symmetry in nuclear single particle spectra. Atomic nuclei are characterized by a very large spin-orbit splitting, i.e., pairs of single particle states with opposite spin ( $j = \ell \pm \frac{1}{2}$ ) have very different energies. This fact allowed the understanding of magic numbers in nuclei and forms the basis of nuclear shell structure. More than thirty years ago [1, 2], pseudo-spin quantum numbers were introduced with  $\tilde{\ell} = \ell \pm 1$  and  $\tilde{j} = \tilde{\ell} \pm \frac{1}{2}$  for  $j = \ell \pm \frac{1}{2}$ , and it was observed that the splitting between pseudo-spin doublets in nuclear single particle spectra is smaller by an order of magnitude than the normal spin-orbit splitting.

After the observation that relativistic mean field models yield spectra with nearly degenerate pseudo spin-orbit partners [3], Ginocchio showed clearly that the origin of pseudo-spin symmetry in nuclei is given by a relativistic symmetry in the Dirac Hamiltonian ([4, 5] and references given therein). He found that pseudo-spin symmetry becomes exact in the limiting case where the strong scalar and vector potentials have the same magnitude but opposite sign. However, this condition is never fulfilled exactly in real nuclei, because in this limit the average nuclear potential vanishes and nuclei are no longer bound. It has been found that the quality of pseudo-spin symmetry is related to the competition between the centrifugal barrier and the pseudo-spin orbital potential [6].

In relativistic investigations a Dirac Hamiltonian is used. In its spectrum one finds single particle levels with positive energies as well as those with negative energies. The latter are interpreted as anti-particles under charge conjugation. This has led to much effort to explore configurations with anti-particles and their interaction with nuclei. The possibility of producing a new kind of nuclear system by putting one or more anti-baryons inside ordinary nuclei has recently gained renewed interest [7]. For future studies of anti-particles in nuclei it is therefore of great importance to investigate the symmetries of such configurations.

In a relativistic description, nuclei are characterized by two strong potentials: an attractive scalar field  $-S(r)$  and a repulsive vector field  $V(r)$  in the Dirac equation, which for nucleons (labelled by a subscript "N") reads:

$$[\alpha \cdot \mathbf{p} + V_N(r) + \beta(M - S_N(r))]\psi_N(r) = \epsilon_N \psi_N(r),$$

where  $V_N(r) = V(r)$  and  $S_N(r) = S(r)$ . For a spherical system, the Dirac spinor  $\psi_N$  has the form

$$\psi_N(r, s) = iG_{n\kappa}(r)Y_{\ell jm}(\theta, \phi, s)j m(\theta, \phi, s) - \tilde{F}_{n\kappa}(r)Y,$$

where  $Y_{\ell jm}(\theta, \phi)$  are the spin spherical harmonics.  $G_{n\kappa}(r)/r$  and  $\tilde{F}_{n\kappa}(r)/r$  form the radial wave functions for the upper and lower components with  $n$  and  $\tilde{n}$  radial nodes.  $\kappa = \langle 1 + \sigma \cdot \mathbf{l} \rangle = (-1)^{j+\ell+1/2}(j+1/2)$  characterizes the spin orbit operator and the quantum numbers  $\ell$  and  $j$ .  $\tilde{\ell} = \ell - \text{sign}(\kappa)$  is the orbital angular momentum of the lower component. It is therefore well accepted that the pseudo-spin quantum numbers of a particle state with positive energy are nothing but the quantum numbers of the lower component [4, 5].

Charge conjugation leaves the scalar potential  $S_N(r)$  invariant while it changes the sign of the vector potential  $V_N(r)$ . That is, for anti-nucleons (labelled by “A”),  $V_A(r) = -V_N(r) = -V(r)$  and  $S_A(r) = S_N(r) = S(r)$ . Charge conjugation of Eq. (2) gives the Dirac spinor for an anti-nucleon,

$$\psi_A(r, s) = j m(\theta, \phi, s) - \tilde{F}_{n\tilde{\kappa}}(r)Y i G_{n\tilde{\kappa}}(r)Y_{\ell jm}(\theta, \phi, s),$$

with  $\tilde{\kappa} = -\kappa$ .

We are only interested in positive energy states of the Dirac equations. Therefore normal quantum numbers follow the upper component which is dominant. A particle state is labeled by  $\{n\ell\kappa m\}$ , while its pseudo-quantum numbers are  $\{\tilde{n}\tilde{\ell}\tilde{\kappa}m\}$ . Following Ref. [8],  $\tilde{n} = n + 1$  for  $\kappa > 0$  and  $\tilde{n} = n$  for  $\kappa < 0$ . An anti-particle state is labeled by  $\{\tilde{n}\tilde{\ell}\tilde{\kappa}m\}$  and its pseudo-quantum numbers are  $\{n\ell\kappa m\}$ . In analogy to Ref. [8], we deduce the relation  $n = \tilde{n} + 1$  for  $\tilde{\kappa} > 0$  and  $n = \tilde{n}$  for  $\tilde{\kappa} < 0$ .

With  $\kappa(1-\kappa) = \tilde{\ell}(\tilde{\ell}+1)$  and  $\kappa(1+\kappa) = \ell(\ell+1)$  in mind, one derives Schrödinger-like equations for the upper and lower components:

$$2M_+ \frac{d^2}{dr^2} + \frac{\ell(\ell+1)}{2M_-} \frac{d^2}{dr^2} - \frac{\tilde{\ell}(\tilde{\ell}+1)}{r^2} + M - V_- G(r) = +\epsilon_N G(r), -\epsilon_A G(r),$$

$$+M - V_+ F(r) = -\epsilon_N F(r), +\epsilon_A F(r),$$

where  $V_{\pm}(r) = V(r) \pm S(r)$  and  $M_{\pm} = M_{\pm}(\epsilon) = M \pm \epsilon \mp V_{\pm}$  with  $\epsilon = +\epsilon_N$  or  $-\epsilon_A$ . Both equations are fully equivalent to the exact Dirac equation with the full spectrum of particle and anti-particle states, but they carry different

quantum numbers. For particle states the first equation carries spin-quantum numbers and the second carries pseudo-spin quantum numbers; for anti-particle states the opposite is true. In the following discussions we will use either the first or second equations according to the type of quantum numbers (spin or pseudo-spin) we are interested in.

We give the relation between spin or pseudo-spin symmetry and the external fields in Table I.

**TABLE I: Relation between symmetry and external fields.**

Condition	Particle Spectrum	Anti-Particle Spectrum
$dV_+/dr = 0$	Pseudo spin symmetry	Spin symmetry
$dV_-/dr = 0$	Spin symmetry	Pseudo spin symmetry

If  $dV_+/dr = 0$ , we have exact spin symmetry in the particle spectrum and exact pseudo-spin symmetry in the anti-particle spectrum because states with the same  $\ell$  (but different  $\kappa$ ) are degenerate in Eq. (5). Here  $\ell$  is the orbital angular momentum of particle states and pseudo orbital angular momentum of anti-particle states. When  $dV_+/dr \neq 0$ , the symmetries are broken. But if  $dV_+/dr$  is so small that the spin-orbit term (the term  $\sim \kappa$ ) in Eq. (5) is much smaller than the centrifugal term, there will be approximate symmetries. For nuclei far from stability where the nuclear potential is expected to be more diffuse, the spin-orbit splitting in single nucleon spectra will also be smaller as compared to stable nuclei. This quenching of the spin-orbit splitting could be one of the reasons for the change of magic numbers in exotic nuclei.

Similarly, when  $dV_-/dr = 0$  in Eq. (6), there is exact pseudo-spin symmetry in the particle spectra [5]. On the other hand, if we focus on anti-particle states, we have in this case exact spin symmetry because now  $\tilde{\ell}$  is the orbital angular momentum. If  $dV_-/dr \neq 0$  but small, we have approximate pseudo-spin symmetry in particle spectra and approximate spin symmetry in anti-particle spectra. This implies that the spin symmetry in the anti-particle spectrum has the same origin as the pseudo-spin symmetry in particle spectrum as realized in Ref. [5]. However, there is an essential difference in the degree to which the symmetry is broken in both cases: the factor  $1/M_-^2 = 1/(M - \epsilon + V_-)^2$  is much smaller for anti-nucleon states than that for nucleon states. The bound anti-particle energies  $\epsilon_A$  are in the region between  $M - V_+(0) \leq \epsilon_A \leq M$ . For realistic nuclei roughly  $0.3 \text{ GeV} \leq \epsilon_A \leq 1 \text{ GeV}$ . On the other hand the bound particle states are in the region of  $M - |V_-(0)| \leq \epsilon_N \leq M$ , i.e., for realistic nuclei close to  $1 \text{ GeV}$ . We therefore have  $|M_-(\epsilon_A)| > 2|M - S(0)|$  and  $|M_-(\epsilon_N)| < |V_-(0)|$ . Thus the factor in front of the  $\tilde{\kappa}$ -term is for anti-particle states by more than a factor  $(2|M - S(0)|/|V_-(0)|)^2 \approx 400$  smaller than for particle states. Spin-symmetry for anti-particle states is therefore much less broken than pseudo-spin symmetry for particle states.

Since the spin-orbit term in Eq. (6) is so small for anti-nucleon states, we expect in addition that the radial wave functions of the spin-doublets are nearly identical, i.e., the dominant components of spin partners for anti-particle solutions are much more similar than the small components of pseudo-spin partners for particles. Although the present discussion is meant for single particle spectra in atomic nuclei, the idea is very general.

It was first discovered that the equality of the vector and scalar potentials results in spin symmetry in Refs. [9, 10] where the authors suggested applications to meson spectra. However, this symmetry was only recently found to be valid for mesons with one heavy quark [11]. In the present letter, we illustrate for the first time in realistic nuclei nearly exact spin symmetry in the single particle spectra for anti-nucleons. We use for that purpose non-linear relativistic mean field (RMF) theory [12] with the modern parameter set NL3. Relativistic Hartree calculations are carried out in coordinate space for the doubly magic nuclei  $^{16}\text{O}$  and  $^{208}\text{Pb}$ .

For  $^{16}\text{O}$ , pseudo-spin symmetry cannot be studied successfully because there are only very few bound nucleon states. However, as seen in [Figure 1: see original paper], there are many more anti-particle states. We find excellent spin symmetry for them. Since there are too many levels in anti-particle spectra of  $^{208}\text{Pb}$  (around 400 for either anti-neutrons or anti-protons), we will not give a similar figure in this case.

In [Figure 2: see original paper] we present the spin-orbit splitting in anti-neutron spectra of  $^{16}\text{O}$  and  $^{208}\text{Pb}$  versus the average energy of a pair of spin doublets. For  $^{16}\text{O}$ , the spin-orbit splittings are around 0.2-0.5 MeV for p states ( $\ell = 1$ ). With increasing particle number  $A$  the spin symmetry in the anti-particle spectra becomes even more exact. For  $^{208}\text{Pb}$ , the spin-orbit splittings are  $\sim 0.1$  MeV for p states and less than 0.2 MeV even for h states ( $\ell = 5$ ) as seen in the lower panel of [Figure 2: see original paper]. We show in **TABLE II** the pseudo-spin orbit splitting of the neutron spectrum of  $^{208}\text{Pb}$  to compare them with the spin-orbit splitting in anti-nucleon spectra. In most cases, the pseudo-spin orbit splittings for particles are larger than 0.4 MeV and for deeply bound states, it can reach even values around 4 MeV.

In general, the spin-orbit splitting decreases as the state approaches the continuum limit. But for very deeply bound anti-neutron p, d, f and g states in  $^{208}\text{Pb}$ , the spin-orbit splitting is smaller. This might be due to the competition between the centrifugal barrier and the spin-orbit potential in Eq. (6).

In order to investigate this in more detail, we calculated the expectation value of the spin-orbit potential,  $\text{SOP} = -\int dr F(r)^2$ . Since the lower amplitudes of the two spin doublets are nearly equal to each other (cf. [Figure 4: see original paper] and [Figure 5: see original paper]), we expect the difference,  $\Delta\text{SOP}$ , gives the main part of  $\Delta\epsilon$  of a pair of spin doublets. In [Figure 3: see original paper] we present  $\Delta\text{SOP}$  as a function of the average energy for spin doublets in  $^{208}\text{Pb}$ . The variational trend of  $\Delta\text{SOP}$  is roughly in agreement with that of

$\Delta\epsilon$ . Particularly, for deeply bound states,  $\Delta\text{SOP} \sim \Delta\epsilon$ .

Wave functions of pseudo-spin doublets in single nucleon spectra have been studied extensively in the literature [5]. The lower amplitudes of pseudo-spin doublets are found to be close to each other. Since the spin symmetry in the anti-nucleon spectrum is much more exact than the pseudo-spin symmetry in the single nucleon spectrum, we expect that the upper amplitudes of the spin doublets coincide with each other even more.

In [Figure 4: see original paper] and [Figure 5: see original paper], we show radial wave functions  $F(r)$  and  $G(r)$  for several anti-nucleon spin doublets in  $^{16}\text{O}$  and  $^{208}\text{Pb}$ . The dominant components  $F(r)$  are nearly exactly identical for the two spin partners. On the other hand the small components  $G(r)$  of the two spin-partners show dramatic deviations from each other. The relation between the node numbers of the upper and lower amplitudes given in Eq. (4) is seen in [Figure 4: see original paper] and [Figure 5: see original paper].

In summary, we discussed the relation between the (pseudo)-spin symmetry in single (anti)-particle states and the external fields where the (anti)-particle moves. We present the single anti-nucleon spectra in atomic nuclei as examples and find an almost exact spin symmetry. The origin of the spin symmetry in anti-nucleon spectra and the pseudo-spin symmetry in nucleon spectra have the same origin but the former is much more conserved in real nuclei. We performed RMF calculations for some doubly magic nuclei. Even in a very light nucleus,  $^{16}\text{O}$ , the spin symmetry in the anti-nucleon spectrum is very good. The spin splitting increases with the orbital quantum number and decreases as the anti-nucleon state approaches the continuum. An investigation of wave functions shows that the dominant components of the Dirac spinor of the anti-nucleon spin doublets are almost identical.

This work was partly supported by the Major State Basic Research Development Program Under Contract Number G2000077407 and the National Natural Science Foundation of China under Grant No. 10025522, 10047001 and 10221003 and by the Bundesministerium für Bildung und Forschung under the project 06 TM 193.

## References

- [1] A. Arima, M. Harvey, and K. Shimizu, Phys. Lett. 30B, 517 (1969).
- [2] K. Hecht and A. Adler, Nucl. Phys. A137, 129 (1969).
- [3] A. L. Blokhin, C. Bahri, and J. P. Draayer, Phys. Rev. Lett. 74, 4149 (1995).
- [4] J. N. Ginocchio, Phys. Rev. Lett. 78, 436 (1997).
- [5] J. N. Ginocchio, Phys. Rep. 315, 231 (1999).
- [6] J. Meng et al., Phys. Rev. C58, R628 (1998); *ibid.* C59, 154 (1999).
- [7] T. Bürvenich et al., Phys. Lett. B542, 261 (2002).
- [8] A. Leviatan and J. N. Ginocchio, Phys. Lett. B518, 214 (2001).
- [9] G. B. Smith and L. J. Tassie, Ann. Phys. 65, 352 (1971).
- [10] J. S. Bell and H. Rugg, Nucl. Phys. B98, 151 (1975).
- [11] P. R. Page, T. Goldman, and J. N. Ginocchio, Phys. Rev. Lett. 86, 204

(2001).

[12] P. Ring, Prog. Part. Nucl. Phys. 37, 193 (1996).

*Note: Figure translations are in progress. See original paper for figures.*

*Source: ChinaXiv – Machine translation. Verify with original.*