

Evaluating the Adaptation of Five Plant Species to Tropical Coral Island Environments Based on Functional Traits: Postprint

Authors: Luo Qi, Liu Hui, Wu Guilin, He Pengcheng, Hua Lei, Zhu Liwei, Zhang Hui, Liu Nan, Jian Shuguang, Ye Qing

Date: 2018-03-08T00:00:00+00:00

Abstract

Vegetation establishment is a critical step for protecting and improving ecosystems of tropical coral islands. The extremely arid habitats of tropical coral islands represent one of the main limiting factors affecting plant survival and colonization; therefore, selecting suitable plant species is crucial for vegetation establishment on tropical coral islands. By measuring photosynthetic/hydraulic functional traits of five plant species—*Scaevola taccada*, *Ipomoea pescaprae*, *Casuarina equisetifolia*, *Arachis hypogaea*, and *Cocos nucifera*—grown in a nursery in Wenchang City, Hainan Province and on tropical coral islands, we explored plant adaptation to tropical coral island habitats. The results showed that, compared with the control (Wenchang nursery), the maximum photosynthetic rate (A_{max}) of *Scaevola taccada*, *Ipomoea pescaprae*, and *Casuarina equisetifolia* on tropical coral islands increased significantly; except for *Cocos nucifera*, the specific leaf area (SLA) of the other four species decreased, while long-term water use efficiency increased (with significant increases in *Casuarina equisetifolia* and *Arachis hypogaea*); leaf carbon content (LC) of all five species decreased to varying degrees. Additionally, leaf hydraulic conductivity (K_{leaf}) of *Ipomoea pescaprae*, *Casuarina equisetifolia*, and *Cocos nucifera* increased significantly; leaf turgor loss point (Ψ_{tlp}) of *Ipomoea pescaprae* and *Arachis hypogaea* decreased significantly; and stomatal conductance (g_s) of *Ipomoea pescaprae* and *Casuarina equisetifolia* increased significantly. The results indicate that *Scaevola taccada*, *Cocos nucifera*, and *Arachis hypogaea* mainly adapt to the arid environment of tropical coral islands through non-stomatal regulation (increasing K_{leaf} , decreasing Ψ_{tlp} and SLA, etc.); whereas *Ipomoea pescaprae* and *Casuarina equisetifolia* promote carbon assimilation and water utilization through both stomatal regulation (decreasing g_s) and non-stomatal regulation (increasing K_{leaf} , decreasing Ψ_{tlp} and SLA, etc.). In summary, *Scaevola taccada*, *Ipomoea pescaprae*, and *Casuarina equisetifolia* exhibit higher photosyn-

thetic capacity and water use efficiency, can effectively coordinate carbon assimilation and water utilization, and demonstrate good adaptive capacity, making them suitable for vegetation establishment on tropical coral islands.

Full Text

Preamble

ACTA ECOLOGICA SINICA

ChinaXiv Partner Journal

Vol. 38, No. 4, Feb. 2018

DOI: 10.5846/stxb201612152597

Using Functional Traits to Evaluate the Adaptability of Five Plant Species on Tropical Coral Islands

LUO Qi^{1,2,3}, LIU Hui^{1,2,3}, WU Guilin^{1,2,3}, HE Pengcheng^{1,2,3}, HUA Lei^{1,2,3}, ZHU Liwei^{1,2}, ZHANG Hui^{1,2}, LIU Nan^{1,2}, JIAN Shuguang^{1,2}, YE Qing^{1,2,*}

¹Key Laboratory of Vegetation Restoration and Management of Degraded Ecosystems, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China

²Guangdong Provincial Key Laboratory of Applied Botany, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China

³University of Chinese Academy of Sciences, Beijing 100049, China

*Corresponding author. E-mail: qye@scbg.ac.cn

<http://www.ecologica.cn>

Abstract

Vegetation establishment is a critical component for protecting and improving tropical coral island ecosystems. Selecting suitable species is therefore essential for successful vegetation restoration on these islands. In this study, we quantified photosynthetic and hydraulic functional traits of five plant species—*Casuarina equisetifolia*, *Ipomoea pescaprae*, *Scaevola taccada*, *Arachis hypogaea*, and *Cocos nucifera*—grown in a nursery in Wenchang, Hainan Province, and transplanted to a tropical coral island with harsh habitats characterized by high solar radiation and low freshwater availability.

Compared to nursery-grown plants, the maximum photosynthetic rate (A_{max}) of *S. taccada*, *C. equisetifolia*, and *I. pescaprae* on the coral island increased significantly, with particularly notable increases in *C. equisetifolia* and *A. hypogaea*. All five species showed decreased leaf carbon content (LC) on the island. The turgor loss point (Ψ_{tlo}) of *I. pescaprae* and *A. hypogaea* decreased significantly, while leaf hydraulic conductivity (K_f) increased significantly in *I. pescaprae* and *C. equisetifolia*. However, stomatal conductance (g_s) of *C. nucifera* increased significantly.

Our results demonstrate that *S. taccada* and *A. hypogaea* adapted to the island environment primarily through non-stomatal regulation (i.e., increased K_f but decreased Ψ_m and SLA). In contrast, *I. pescaprae* and *C. equisetifolia* employed both stomatal regulation (i.e., decreased g) and non-stomatal regulation (increased K_f but decreased Ψ_m and SLA) to promote carbon assimilation and water use. *C. equisetifolia* exhibited higher photosynthetic capacity and water use efficiency, enabling better coping with harsh environmental conditions. Therefore, *C. equisetifolia*, *I. pescaprae*, and *S. taccada* are recommended as suitable species for vegetation restoration on tropical coral islands.

Keywords: tropical coral island; ecological adaptability; vegetation restoration; photosynthesis; functional traits

Introduction

With rapid global economic development and depletion of terrestrial resources, ocean development has become a strategic priority for many countries. As important components of marine systems, islands possess tremendous resource value, and their rational development is crucial for national sustainable development, attracting high attention from governments and academia. Islands also serve as important tourism resources that satisfy growing recreational demands. However, tropical coral islands present extreme environmental challenges including high temperatures, intense light, and substrates with low water-holding capacity, minimal soil, high salinity, and poor nutrient content. These conditions severely limit plant survival and establishment, making vegetation restoration essential for creating habitable environments and supporting sustainable development.

Species selection is key to successful vegetation restoration on tropical coral islands. Plant adaptability is influenced by multiple factors including soil conditions, species characteristics, and environmental stress. Drought stress represents a primary limiting factor, significantly affecting plant survival rates. During long evolution, plants have developed effective mechanisms to adapt to drought through stomatal regulation (physiological control of transpiration through stomatal structure and environmental feedback) and non-stomatal regulation (including leaf hydraulic conductivity, turgor loss point, and specific leaf area). Leaf hydraulic resistance constitutes a major bottleneck (>30%) in whole-plant water transport, with leaf hydraulic conductivity reflecting water transport rates and efficiency closely linked to photosynthesis. The turgor loss point indicates drought resistance capacity, while specific leaf area (SLA)—the fresh leaf area per unit dry weight—is associated with drought tolerance, with lower SLA indicating stronger turgor maintenance capacity.

Previous research on tropical coral island ecology has focused on environmental assessment and community dynamics, with few studies examining physiological adaptation mechanisms of suitable species. While Li et al. investigated eco-

logical adaptability of four plant species from carbon assimilation perspectives, they did not consider hydraulic traits or differences in drought strategies. This study examines five potential suitable species transplanted from a Wenchang nursery to a tropical coral island, comparing their photosynthetic and hydraulic characteristics to evaluate adaptability to the island's arid environment.

1. Study Site Description

The experimental nursery was located in the suburbs of Wenchang City, Hainan Province (110°45 E, 19°31 N), characterized by a tropical maritime monsoon climate with mean annual temperature of ~24°C and precipitation of ~1,800 mm. The soil type was coastal deposit sandy loam.

The tropical coral island study site had an elevation of approximately 5 m, with tropical maritime monsoon climate, mean annual temperature of ~28°C, and precipitation of ~2,800 mm with uneven distribution. The substrate consisted of coral sand with extreme environmental characteristics including high salinity and poor nutrient content, supporting virtually no vegetation.

2. Experimental Materials and Methods

Based on previous surveys of the Xisha Islands, five potential suitable species were selected: *Scaevola taccada*, *Ipomoea pescaprae*, *Casuarina equisetifolia*, *Arachis hypogaea*, and *Cocos nucifera*. All island plants were transplanted in March 2015. Except for minimal imported soil used during transplantation to improve survival, no additional artificial care was provided. The differences in environmental factors (water conditions, light radiation) between sites were the focus of this comparative study.

Measurements were taken on fully expanded, healthy mature leaves in September 2015.

2.1 Photosynthesis

Maximum photosynthetic rate (A_{max}), transpiration rate (Tr), and stomatal conductance (g) were measured in situ on sunny days between 9:00-11:00 using a LI-6400 portable photosynthesis system (LI-COR, Nebraska, USA). Light intensity was set at 1,500 $\mu\text{mol m}^{-2} \text{s}^{-1}$, CO₂ concentration at 400 $\mu\text{mol/mol}$, and data were recorded after 3-5 minutes of stabilization. Instantaneous water use efficiency was calculated as $WUE = A_{max} / \text{Tr}$.

2.2 Turgor Loss Point

Sun-exposed leaves were collected, immediately placed in water-filled buckets, sealed in black plastic bags to prevent transpiration, and saturated overnight

(16:00–18:00). Saturated fresh weight (SW) and water potential (Ψ) were measured using a pressure chamber (PMS, Corvallis, Oregon, USA). Leaves were then placed on a bench to desiccate, with water potential and weight measured repeatedly until water potential declined insignificantly. After measurement, leaves were oven-dried to determine dry weight (DW). Relative water content was calculated as: $RWC = 100 \times (FW - DW)/(SW - DW)$. The turgor loss point water potential (Ψ_{tlo}) was determined by fitting the water potential-RWC curve.

2.3 Leaf Hydraulic Conductivity

Leaf hydraulic conductivity (K_{lf}) was measured using the method of Franks: $K_{lf} = \Delta f / (10 \times A_{lf} \times (\Psi_{sat} - \Psi)^2)$, where Δf is water flow rate over 10 seconds, A_{lf} is leaf area, Ψ_{sat} is saturated leaf water potential, and Ψ is applied pressure potential (0.3–0.5 MPa).

2.4 Specific Leaf Area

Leaf area (LA) was measured with a leaf area meter (Li-3000A, Li-Cor, Lincoln, NE, USA). Leaves were then oven-dried to determine dry weight (DW). SLA was calculated as: $SLA = LA/DW$.

2.5 Stable Carbon Isotope Content

Dried, ground samples were sent to the South China Botanical Garden Public Laboratory for stable carbon isotope analysis using an IsoPrime100 isotope ratio mass spectrometer (IsoPrime, Manchester, UK). Stable carbon isotope content (^{13}C) is a reliable indicator of long-term water use efficiency, with higher values indicating greater efficiency.

3. Data Analysis

SPSS 18.0 software was used for t-tests comparing traits of the same species between habitats. Origin 16.0 was used for graphical presentation.

3. Results

3.1 Photosynthetic Rate and Leaf Economic Traits

Compared to Wenchang nursery plants, A_{lf} of *S. taccada*, *C. equisetifolia*, and *I. pescaprae* on the coral island increased significantly, while *C. nucifera* showed no significant difference. Transpiration rate of *C. equisetifolia* on the island increased significantly, while other species showed no significant differences. Specific leaf area (SLA) of four species (except *C. nucifera*) on the island decreased

significantly compared to nursery plants. Leaf carbon content (LC) of all five species on the island decreased, with significant reductions in *S. taccada* and *C. nucifera*.

[Figure 1: see original paper] Comparison of A_{net} , Tr, SLA, and LC values of five plant species in Wenchang nursery and island (Mean \pm SE). * indicates significant difference between sites ($P < 0.05$); ** indicates highly significant difference ($P < 0.01$); NS indicates no significant difference.

3.2 Water Use Efficiency

Instantaneous water use efficiency (WUE) of *I. pescaprae* on the coral island was significantly higher than nursery plants, while *C. equisetifolia* and *A. hypogaea* showed significant differences. Stable carbon isotope content (^{13}C) of all species except *C. nucifera* increased on the island, with significant differences in *C. equisetifolia* and *A. hypogaea*.

[Figure 2: see original paper] Comparison of WUE and ^{13}C values of five plant species in Wenchang nursery and island.

3.3 Hydraulic Traits

Stomatal conductance (g_s) of *I. pescaprae* and *C. equisetifolia* on the coral island increased significantly compared to nursery plants, while other species showed no significant differences. Leaf hydraulic conductivity (K_{leaf}) of *I. pescaprae* and *C. nucifera* increased significantly, while *A. hypogaea* decreased significantly. Turgor loss point water potential (Ψ_{tlo}) of *I. pescaprae* and *A. hypogaea* on the island decreased significantly compared to Wenchang.

[Figure 3: see original paper] Comparison of g_s , K_{leaf} , and Ψ_{tlo} values of five plant species in Wenchang nursery and island.

4. Discussion

4.1 Photosynthetic Rate and Leaf Economic Traits

Photosynthetic rate reflects carbon assimilation capacity and is an important indicator of comprehensive environmental adaptability. This study found that A_{net} of transplanted *S. taccada*, *I. pescaprae*, and *C. equisetifolia* was higher than nursery plants, indicating strong adaptability to coral island conditions. The lack of significant difference in *C. nucifera* suggests different adaptive strategies. Previous studies show transplanted seedlings undergo tolerance, recovery, and stable phases with different physiological responses, which may explain discrepancies with earlier research.

Specific leaf area (SLA) is closely related to drought resistance and photosynthetic capacity. Lower SLA indicates greater investment in protective structures

and higher mesophyll cell density to prevent excessive water loss. The strong light and drought conditions on coral islands promoted most species to allocate more resources to leaf construction, reducing SLA to avoid radiation damage and water loss.

Leaf carbon content (LC) represents construction costs and reflects the leaf economics spectrum—the trade-off between resource investment and returns. The worldwide leaf economics spectrum shows that species with low LC and high photosynthetic rates occupy one end, while the opposite traits occupy the other. The significant increase in photosynthetic rate combined with decreased LC in island plants confirms this trade-off, showing greater energy allocation to photosynthetic capacity rather than leaf construction.

4.2 Water Use Efficiency

Water use efficiency (WUE) measures assimilated carbon per unit water consumed. Instantaneous WUE (A / Tr) reflects short-term efficiency, while stable carbon isotope composition (^{13}C) estimates long-term efficiency. Our results show that *I. pescaprae* and *C. equisetifolia* on the island had significantly higher instantaneous WUE, while ^{13}C increased in all species except *C. nucifera*, indicating improved long-term water use efficiency under drought conditions. This aligns with research showing drought increases ^{13}C values.

4.3 Stomatal vs. Non-stomatal Regulation

Stomata regulate the balance between water loss and carbon assimilation. Compared to nursery plants, *I. pescaprae* and *C. equisetifolia* on the island showed increased g but no corresponding increase in transpiration rate, suggesting stomatal structural adjustments that maintain photosynthesis while reducing water loss. This contradicts some studies showing stomatal closure under drought, indicating that island stress may trigger structural modifications for more uniform stomatal distribution.

Leaf hydraulic conductivity (K_f) is positively correlated with photosynthetic rate within certain ranges. *I. pescaprae* and *C. nucifera* significantly increased K_f on the island, enhancing water transport to support photosynthesis. In contrast, *A. hypogaea* showed decreased K_f , suggesting its stress tolerance mechanisms were compromised.

The significantly lower Ψ in *I. pescaprae* and *A. hypogaea* indicates enhanced osmotic adjustment capacity, an important non-stomatal adaptation to drought. Combined with reduced SLA, these mechanisms prevent excessive water loss and improve drought tolerance.

5. Conclusion

This study demonstrates that different species employ distinct strategies to adapt to tropical coral island environments. *Arachis hypogaea* primarily uses non-stomatal regulation (increased K_f , decreased Ψ and SLA). *Casuarina equisetifolia* and *Ipomoea pescaprae* use both stomatal and non-stomatal regulation to promote carbon assimilation and water use. *C. equisetifolia* and *I. pescaprae* exhibit higher photosynthetic capacity and water use efficiency, showing good adaptability and suitability for tropical coral island restoration.

Considering photosynthesis, water transport, and drought resistance, these three species demonstrate strong adaptability to the harsh conditions of tropical coral islands and are recommended for ecosystem restoration. However, plant adaptation is a long-term process requiring continuous monitoring of more species to provide a scientific basis for selecting suitable plants and improving ecosystem services on tropical coral islands.

References

- [1] Study on land resource development suitability of uninhabited islands in Shanghai. *Journal of Shanghai Ocean University*, 2013.
- [2] Risk assessment and management of tropical coral island reef ecosystems. East China Normal University, 2013.
- [3] Vegetation ecological survey of Xisha Islands. *Chinese Agricultural Science Bulletin*, 2012, 21(1): 92-96.
- [4] Screening of drought-resistant plants and research on drought resistance technology in island vegetation restoration. *Journal of Applied Oceanography*, 2016, 35(2): 223-228.
- [5] Research progress on water consumption regulation mechanisms of forest trees. *Chinese Journal of Ecology*, 2006, 25(6): 692-697.
- [6] Sack L, Holbrook N M. Leaf hydraulics. *Annual Review of Plant Biology*, 2006, 57: 361-381.
- [7] Scoffoni C, McKown A D, Rawls M, Sack L. Dynamics of leaf hydraulic conductance with water status: quantification and analysis of species differences under steady state. *Journal of Experimental Botany*, 2012, 63(2): 643-658.
- [8] Brodribb T J, Holbrook N M. Changes in leaf hydraulic conductance during leaf shedding in seasonally dry tropical forest. *New Phytologist*, 2003, 158(2): 295-303.
- [9] Sack L, Frole K. Leaf structural diversity is related to hydraulic capacity in tropical rainforest trees. *Ecology*, 2006, 87(2): 483-491.

- [10] Study on water parameters of artificial forests in western Beijing mountains (I). *Journal of Beijing Forestry University*, 1994, 16(1): 1-12.
- [11] Bucci S J, Goldstein G, Meinzer F C, Scholz F G, Franco A C, Bustamante M. Functional convergence in hydraulic architecture and water relations of tropical savanna trees: from leaf to whole plant. *Tree Physiology*, 2004, 24(8): 891-899.
- [12] Reich P B, Walters M B, Ellsworth D S, Vose J M, Volin J C, Gresham C, Bowman W D. Relationships of leaf dark respiration to leaf nitrogen, specific leaf area and leaf life-span: a test across biomes and functional groups. *Oecologia*, 1998, 114(4): 471-482.
- [13] Ecological research progress on leaf lifespan and related leaf traits. *Chinese Journal of Plant Ecology*, 2004, 28(6): 844-852.
- [14] Response of specific leaf area of main tree species in secondary forests of Northeast China to light intensity. *Chinese Journal of Ecology*, 2009, 28(8): 1437-1442.
- [15] Smith Jr K L, Baldwin R J, Kaufmann R S, Sturz A. Ecosystem studies at Deception Island, Antarctica: an overview. *Deep Sea Research Part II: Topical Studies in Oceanography*, 2003, 50(10/11): 1595-1609.
- [16] Community dynamics during vegetation restoration on Nanao Island, Guangdong. *Chinese Journal of Ecology*, 2006, 25(4): 566-571.
- [17] Discussion on comprehensive island ecological assessment methods. *Biodiversity Science*, 2013, 21(3): 364-374.
- [18] Ecological adaptability of four plant species to tropical coral island environments. *Chinese Journal of Plant Ecology*, 2001, 25(3): 298-305.
- [19] Plant diversity of Xisha Islands, China. *Ecology and Environmental Sciences*, 2016, 25(5): 790-794.
- [20] Tyree M T, Richter H. Alternative methods of analysing water potential isotherms: some cautions and clarifications: I. the impact of non-ideality and of some experimental errors. *Journal of Experimental Botany*, 1981, 32(3): 643-653.
- [21] Schulte P J, Hinckley T M. A comparison of pressure-volume curve data analysis techniques. *Journal of Experimental Botany*, 1985, 36(10): 1590-1602.
- [22] Franks P J. Higher rates of leaf gas exchange are associated with higher leaf hydrodynamic pressure gradients. *Plant, Cell & Environment*, 2006, 29(4): 584-592.
- [23] Gower S T, Kucharik C J, Norman J M. Direct and indirect estimation of leaf area index, APAR, and net primary production of terrestrial ecosystems. *Remote Sensing of Environment*, 1999, 70(1): 29-51.

- [24] Application of stable carbon isotope technology in ecological research. *Chinese Journal of Plant Ecology*, 2002, 26(5): 549-560.
- [25] Lambers H, Chapin III F S, Pons T L. *Plant Physiological Ecology*. 2nd ed. New York: Springer, 2008: 11-99.
- [26] Physiological responses of jujube seedlings to transplanting stress and physiological basis of anti-transpirant mitigation. Henan Agricultural University, 2008.
- [27] Wright I J, Reich P B, Westoby M, et al. The worldwide leaf economics spectrum. *Nature*, 2004, 428(6985): 821-827.
- [28] Leaf functional traits of ferns in different habitats in Dinghushan monsoon evergreen broad-leaved forest. *Journal of Tropical and Subtropical Botany*, 2013, 21(6): 489-495.
- [29] Response and simulation of water use efficiency in different plant leaves to light and CO₂. *Acta Ecologica Sinica*, 2008, 28(2): 525-533.
- [30] Farquhar G D, O' Leary M H, Berry J A. On the relationship between carbon isotope discrimination and the intercellular carbon dioxide concentration in leaves. *Australian Journal of Plant Physiology*, 1982, 9(2): 121-137.
- [31] Saugier B, Ehleringer J R, Hall A E, Farquhar G S. *Stable Isotopes and Plant Carbon-Water Relations*. San Diego: Elsevier, 2012.
- [32] Application of carbon stable isotope technology in plant water stress research. *Chinese Journal of Plant Ecology*, 2004, 24(5): 1027-1033.
- [33] Relationship between leaf carbon stable isotope composition and water use efficiency. *Chinese Journal of Plant Ecology*, 2011, 35(2): 119-124.
- [34] Meinzer F C, Grantz D A. Coordination of stomatal, hydraulic, and canopy boundary layer properties: do stomata balance conductances by measuring transpiration? *Physiologia Plantarum*, 1991, 83(2): 324-329.
- [35] Meinzer F C, Goldstein G, Holbrook N M, Jackson P, Cavelier J. Stomatal and environmental control of transpiration in a lowland tropical forest tree. *Plant, Cell & Environment*, 1993, 16(4): 429-436.
- [36] Hérault A, Lin Y S, Bourne A, Medlyn B E, Ellsworth D S. Optimal stomatal conductance in relation to photosynthesis in climatically contrasting *Eucalyptus* species under drought. *Plant, Cell & Environment*, 2013, 36(2): 262-274.
- [37] Comparative analysis of crop photosynthesis: constraints of stomatal and internal conductance. *Journal of Northeast Agricultural University*, 2011, 42(1): 129-132.
- [38] Effects of elevated CO₂ concentration on stomatal density and distribution in spring wheat in arid regions. *Chinese Journal of Plant Ecology*, 2001, 25(3): 312-316.

- [39] Effects of drought stress on cucumber seedling growth, photosynthetic physiology, and stomatal characteristics. *Journal of China Agricultural University*, 2010, 15(5): 12-18.
- [40] Brodribb T J, Holbrook N M, Zwieniecki M A, Palma B. Leaf hydraulic capacity in ferns, conifers and angiosperms: impacts on photosynthetic maxima. *New Phytologist*, 2005, 165(3): 839-846.
- [41] Brodribb T J, Feild T S, Jordan G J. Leaf maximum photosynthetic rate and venation are linked by hydraulics. *Plant Physiology*, 2007, 144(4): 1890-1898.
- [42] Brodribb T J, Jordan G J. Water supply and demand remain balanced during leaf acclimation of *Nothofagus cunninghamii* trees. *New Phytologist*, 2011, 192(2): 437-448.
- [43] Holloway-Phillips M M, Brodribb T J. Minimum hydraulic safety leads to maximum water-use efficiency in a forage grass. *Plant, Cell & Environment*, 2011, 34(2): 302-313.
- [44] Effects of seasonal drought stress on water and photosynthetic physiology of three evergreen dominant tree species in limestone forests. *Acta Botanica Yunnanica*, 2008, 30(3): 325-332.

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv – Machine translation. Verify with original.