

Lyapunov-type inequalities for a class of nonlinear higher-order differential equations

Authors: Jun Zheng, Haofan Wang, Jun Zheng

Date: 2018-06-04T00:00:00+00:00

Abstract

In this work, we establish several Lyapunov-type inequalities for a class of nonlinear higher order differential equations having a form

$$(\psi(u^{(m)}(x)))' + \sum_{i=0}^n r_i(x) f_i(u^{(i)}(x)) = 0,$$

with anti-periodic boundary conditions, where $m > n \geq 0$ are integers, ψ and $f_i (i = 0, 1, 2, \dots, n)$ satisfy certain structural conditions such that the considered equations have general nonlinearities. The obtained inequalities are extensions and complements of the existing results in the literature.

Full Text

Preamble

Lyapunov-type inequalities for a class of nonlinear higher order differential equations

Jun Zheng¹, Haofan Wang²

¹School of Mathematics, Southwest Jiaotong University, Chengdu, Sichuan, P. R. of China 611756

²School of Information Science & Technology, Southwest Jiaotong University, Chengdu 611756, Sichuan, China

Abstract

In this work, we establish several Lyapunov-type inequalities for a class of nonlinear higher order differential equations having a form $(\psi(u^{(m)}(x)))' + \sum_{i=0}^n r_i(x) f_i(u^{(i)}(x)) = 0$, with anti-periodic boundary conditions, where $m > n \geq 0$ are integers, and $f_i (i = 0, 1, 2, \dots, n)$ satisfy certain structural conditions such that

the considered equations have general nonlinearities. The obtained inequalities are extensions and complements of the existing results in the literature.

Key words: Lyapunov inequality, Higher order differential equation, nonlinear equation.

Introduction

If u is a nontrivial solution of the following second order differential equation $u'' + r(x)u(x) = 0$, $x \in (a, b)$, $u(a) = u(b) = 0$, where r is a continuous and nonnegative function defined in $[a, b]$, then the following inequality $\int_a^b r(x)dx > 2$ holds, which was first proved by Lyapunov [7] and known as “Lyapunov inequality”. The Lyapunov inequality and its various generalizations have proved to be useful tools in oscillation theory, disconjugacy, eigenvalue problems, and numerous other applications for the theories of differential and difference equations, and also in time scales. Since the appearance of Lyapunov’s fundamental paper, there are many improvements and generalizations of (1) in the literature. Especially, the Lyapunov inequality has been generalized extensively to the higher-order linear equations see [1-4,6,8-13,15,14,16,18] and references therein.

But so far, only a few results have been achieved for higher order differential equations that have either general nonlinearities or anti-periodic boundary conditions. For example, [12] and [17] have contributed to certain higher equations with anti-periodic boundary conditions, and to general nonlinear second order equations with Dirichlet boundary conditions respectively. Corresponding author: J. Zheng Email addresses: zhengjun@swjtu.edu.cn (Jun Zheng¹), wanghaofan@my.swjtu.edu.cn (Haofan Wang²).

Preprint submitted to

4 June 2018

In 2012, Wang [12] considered Lyapunov-type inequalities for certain higher order differential equations with anti-periodic boundary conditions. The main result in [12] is as follows:

Theorem A If $u(x)$ is a nonzero solution of the following $(m+1)$ -order half-linear differential equation anti-periodic boundary conditions: $(|u'(x)|^2 u(x))' + r(x)|u(x)|^2 u(x) = 0$, $x \in (a, b)$, $u(a) + u(b) = 0$, $i = 0, 1, 2, \dots, m$, $\int_a^b |r(x)|dx > 2$

In a recent work [17], the authors established Lyapunov-type inequalities for a class of nonlinear second order equations with a general form as $(|u'(x)|^2 + r(x)f(u(x)))' = 0$, $x \in (a, b)$, $u(a) = u(b) = 0$.

The structural conditions are given as below, which allow the functions ψ and f to have a large class of nonlinearities (see Section 1 and 4 in [17], and Remark 1, for instance): (A) $\psi \in C((-\infty, \infty)) \cap C^1((0, \infty))$ with $f \geq 0$ on $(-\infty,$

∞). (B) ϕ is odd on $(-\infty, \infty)$. (C) $f(t) \geq 0$ for all $t \in [0, \infty)$. (D) There exists $k > 0$ such that $|f(t)| \leq k(|t|)$ for all $t \in (-\infty, \infty)$. (E) There exists constants $\alpha, \beta > 0$ such that $\phi(t) \leq \alpha \phi(t) + \beta$, $t > 0$. (F) There exists constants $\alpha, \beta > 0$ such that $\phi(t) \leq \alpha \phi(t) + \beta$, $t > 0$.

The main result in [17] is the following theorem.

Theorem B Suppose that conditions (A)-(D) are satisfied. Let u be a nontrivial solution of (3). (i) If ϕ satisfies (E), then $\int_a^b |r(x)|dx \leq \int_a^b |r(x)|dx + 1 + \alpha \cdot \min\{2, 1 + \alpha\} \cdot \min\{2, 1 + \alpha\}$, (cid:1) , (cid:0) 2 (cid:1) , (cid:0) 2 (cid:1) (cid:9). (cid:1) (cid:9).

Our motivation for this paper comes from the papers of [12] and [17]. We attempt to establish several new Lyapunov-type inequalities for a large class of nonlinear higher differential equations under anti-periodic boundary conditions. The results obtained in this paper are generalizations of [12] and [17].

2 Problem setting and main results

In this paper, we establish Lyapunov inequalities for the following equation $(u^{(m)}(x))' + r(x)f(u^{(n)}(x)) = 0$, $x \in (a, b)$, $u(a) + u(b) = 0$, $j = 0, 1, 2, \dots, m$, where r is a given function defined in $[a, b]$, and f satisfy some of the following structural conditions: (H1) $f \in C((-\infty, \infty)) \cap C^1((0, \infty))$ with $|f| \leq k(|t|)$ on $(-\infty, \infty)$, $i = 0, 1, 2, \dots, n$. (H2) ϕ is odd on $(-\infty, \infty)$. (H3) $f(t) \geq 0$ for all $t \in [0, \infty)$, $i = 0, 1, 2, \dots, n$. (H4) There exists constants $k > 0$ satisfying $k > 0$ such that $|f(t)| \leq k(|t|)$ for all $t \in (-\infty, \infty)$, $i = 0, 1, 2, \dots, n$. (H) There exists constants $\alpha, \beta > 0$ such that $\phi(t) \leq \alpha \phi(t) + \beta$, $t > 0$. (Hf) There exists constants $\alpha, \beta > 0$ such that $\phi(t) \leq \alpha \phi(t) + \beta$, $t > 0$, $i = 0, 1, 2, \dots, n$.

Remark 1 (i) The structural condition (H) (or (Hf)) was firstly presented in [17], which is a slight version given by $\ln d$ in (H) when $\log(b|t|+d)$, $b > 0$, $c, d > \ln d$, $\alpha = a$ in (H). More examples of nonlinear functions satisfying (H) can be found in [17, Lieberman [5]. Observe that $\alpha = p - 1$ when $\phi(t) = |t|^{2p} (p > 1)$, and $\alpha = a$, $\beta = a + 1$ $\phi(t) = |t|^{-1} t \log(b|t| + d)$, $a, b > 0$, $c, d > 1$. Another interesting example of ϕ is $\phi(t) = |t|^{-1} t$, $a > 1$ Section 1 and 4]. In d with $\alpha = a - 1$

- (ii) By (H1) – (H4), $\phi(0) = f(0) = 0$ ($i = 0, 1, 2, \dots, n$) and $\phi(t) \geq 0$ for any $t \geq 0$. Furthermore, if ϕ (or f) satisfies $\phi(t) \geq 0$, which guarantees the increasing monotonicity of $\phi(t)$ (or $f(t)$) in $t \geq 0$. (H) (or (Hf)), then $\phi(t) \geq 0$ (or $f(t) \geq 0$), which guarantees the increasing monotonicity of $\phi(t)$ (or $f(t)$) in $t \geq 0$.

Throughout this paper, we always assume that $m > n \geq 0$ are integers, $r \in L^1(a, b)$ with $r \geq 0$ on (a, b) ($i = 0, 1, 2, \dots, n$), and conditions (H1) – (H4) are satisfied. Moreover, we say u is a solution of (4) if $u \in C^1(a, b) \cap C([a, b])$,

$(u(x))$ is absolutely continuous in x , and u satisfies the equation in (4) almost everywhere in (a, b) .

The main result in this paper is given below.

Theorem 1 Suppose that (4) admits a nontrivial solution. (i) If ψ satisfies (H), then (cid:18) (cid:80) (ii) If f satisfies (Hf) ($i = 0, 1, 2, \dots, n$), then (cid:82) $b - a \int_a^b |dx| \cdot k \cdot \max$ (cid:18) (cid:80) (cid:26)(cid:18) (cid:19) $(m-i)$ (cid:18) (cid:19) $(m-i)$ (cid:27)(cid:19) $2(1+)$ $1+$. (cid:18) (cid:19) $(m-i)$ (cid:82) $b - a \int_a^b |dx| \cdot k \cdot 1+$ $\cdot \max$ (cid:26)(cid:18) (cid:19) $(m-i)$ (cid:27)(cid:19)

Corollary 2 If u is a nontrivial solution of (4) with $(t) = f(t) = |t|^{2t}$, $p > 1$, $i = 0, 1, 2, \dots, n$, then (cid:88) (cid:18) (cid:90) $b - a \int_a^b |dx| \cdot$ (cid:18) $b - a$ (cid:19) (cid:19) $(^{-1})$ (cid:19) Particulary, if $n = 0$, there holds (cid:82) $b - a \int_a^b |dx| 2^{-1} (b-a)^{-1}$, which is the same as (2).

3 Proof of main result

We present first some auxiliary results needed in the main proofs. In this section, we always write $\Psi(t) =$ (cid:82) $t F(t) =$ (cid:82) $t \int_0^t f(s)ds$ ($i = 0, 1, 2, \dots, n$) for $t \in [0, +\infty)$ and

Lemma 3 [17] Assume that ψ satisfies (H1)-(H4) and (H). The following results hold true. (i) $(st) \max\{s, s\}(t)$, $s, t \geq 0$. (ii) Ψ is C^2 -continuous on $(0, +\infty)$, and convex on $[0, +\infty)$. (iii) $t(t) \geq \Psi(t) \geq t(t) \geq 0$, $t \geq 0$.

Remark 2 The function f satisfying (H1)-(H4) and (Hf) and the function F have similar properties as above.

Lemma 4 Let u be a C -continuous function on $[a, b]$ satisfying (4b). The following inequalities hold: $\int_a^b |u(x)|$ (cid:18) $b - a$ (cid:19) $b - a$ (cid:90) $b - a$ (cid:19) $\int_a^b |u(x)|dx$, $i = 0, 1, 2, \dots, m - 1$.

Proof. It suffices to note that $|u(x)| =$ (cid:90) x (cid:12) (cid:12) (cid:12) (cid:12) (cid:12) (cid:18) (cid:90) x (cid:90) $b - a \int_a^b u^{-1}(y)dy +$ (cid:12) $u^{-1}(y)$ (cid:12) (cid:12) (cid:12) $dy +$ (cid:90) $b - a \int_a^b u^{-1}(y)dy$ (cid:12) (cid:12) (cid:12) (cid:12) $u^{-1}(y)$ (cid:12) (cid:12) (cid:19) (cid:12) $dy \int_a^b |u^{-1}(y)|dy$ ($i = 0, 1, 2, \dots, m - 1$).

Lemma 5 Let u be a C -continuous function on $[a, b]$ satisfying (4b). (i) If f satisfies (Hf) ($i = 0, 1, 2, \dots, n$), then (cid:18)(cid:18) $b - a$ (cid:19) $(1 +) \cdot \max$ (cid:90) $b - a$ (cid:26)(cid:18) $b - a$ (cid:19) $\int_a^b |u(x)|dx$ (cid:90) $b - a \int_a^b |u(x)|dx$ (cid:19) $(m-i)$ (cid:18) $b - a$ (cid:19) $(m-i)$ (cid:27) (cid:90) $b - a \int_a^b |u(x)|dx$. (ii) If ψ satisfies (H), then for $i = 0, 1, 2, \dots, n$, there holds (cid:18)(cid:18) $b - a$ (cid:19) $(1 +) \cdot \max$ $b - a$ (cid:26)(cid:18) $b - a$ (cid:90) $b - a$ (cid:19) $\int_a^b |u(x)|dx$ (cid:90) $b - a \int_a^b |u(x)|dx$ (cid:19) $(m-i)$ (cid:19) $(m-i)$ (cid:27) (cid:18) $b - a$ (cid:90) $b - a \int_a^b |u(x)|dx$.

Proof. We only prove (i). Indeed, by Lemma 3 (i)-(iii), we get (cid:18)(cid:18) $b - a$ (cid:19) (cid:26)(cid:18) $b - a$ (cid:90) $b - a$ (cid:19) $(m-i) \max \max$

$F(|u|)dx$ (cid:19) $(m-i)$ (cid:27)(cid:19)

Note that $k > 0$, $1 + k > 0$ and $F(|u|) > 0$, arguing as in the proof of (i), we may get $I > 0$. Finally, (5) implies (cid:4) the desired result.

References

- [1] R. Agarwal and A. Özbekler. Lyapunov type inequalities for nth order forced differential equations with mixed nonlinearities. *Communications on Pure and Applied Analysis*, 15(6):2281–2300, 2016.
- [2] D. Çakmak. Lyapunov-type integral inequalities for certain higher order differential equations. *Appl. Math. Comput.*, 216(2):368–373, 2010.
- [3] S. Dhar and Q. Kong. Liapunov-type inequalities for odd order linear differential equations. *Electron. J. Differential Equations*, 2016(243):1–10, 2016.
- [4] X. He and X. Tang. Lyapunov-type inequalities for even order differential equations. 11(2):465–473, 2012.
- [5] G. M. Lieberman. The natural generalization of the natural conditions of Ladyzhenskaya and Ural'tseva for elliptic equations. *Communications in Partial Differential Equations*, 16(2-3):311–361, 1991.
- [6] X. Lin and Z. Zhao. Iterative technique for a third-order differential equation with three-point nonlinear boundary value conditions. *Electronic Journal of Qualitative Theory of Differential Equations*, 2016(12):1–10, 2016.
- [7] A. Lyapunov. *Probleme General de la Stabilite du Mouvement*, in: Ann. Math. Studies, volume 17. Princeton Univ. Press, 1949. Reprinted from Ann. Fac. Sci. Toulouse, 9 (1907) 203–474, Translation of the original paper published in Comm. Soc. Math. Kharkow, 1892.
- [8] S. Panigrahi. Liapunov-type integral inequalities for certain higher order differential equations. *Electron. J. Differential Equations*, 2009(28):571–582.
- [9] N. Parhi and S. Panigrahi. On liapunov-type inequality for third-order differential equations. *J. Math. Anal. Appl.*, 233(2):445–460, 1999.
- [10] N. Parhi and S. Panigrahi. Liapunov-type inequality for higher order differential equations. *Math. Slovaca*, 52(1):31–46, 2002.
- [11] Y. Qi, Y. Peng, and X. Wang. Liapunov-type inequalities for a class third-order differential equations. *Journal of Pure and Applied Mathematics: Advances and Applications*, 17(2):89–103, 2017.
- [12] Y. Wang. Lyapunov-type inequalities for certain higher order differential equations with anti-periodic boundary conditions. *Appl. Math. Letters*, 25(12):2375–2380, 2012.
- [13] X. Yang. On liapunov-type inequality for certain higher-order differential equations. *Appl. Math. Comput.*, 134(2-3):307–317, 2003.

- [14] X. Yang, Y. Kim, and K. Lo. Lyapunov-type inequality for quasilinear systems. *Appl. Math. Lett.*, 34(1):33–36, 2014.
- [15] X. Yang and K. Lo. Lyapunov-type inequality for a class of even-order differential equations. *Appl. Math. Comput.*, 215(11):3884–3890, 2010.
- [16] Q. Zhang and X. He. Lyapunov-type inequalities for a class of even-order differential equations. *Journal of Inequalities and Applications*, 2012. 2012:5.
- [17] J. Zheng and X. Guo. Lyapunov-type inequalities for p -laplacian equations. 2018. chinaXiv:201805.00171.
- [18] X. Zuo and W. Yang. Lyapunov-type inequalities for higher-order differential equations with one-dimensional p -Laplacian. *Journal of Mathematical inequalities*, 8(4):737–746, 2014.

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv –Machine translation. Verify with original.