

Evaluating and modeling the spatiotemporal pattern of regional-scale salinized land expansion in highly sensitive shoreline landscape of southeastern Iran postprint

Authors: Mohammad, SHAFIEZADEH, Hossein, MORADI, Sima, FAKHERAN, Hossein, MORADI

Date: 2018-10-29T00:00:00+00:00

Abstract

Taking an area of about 2.3×10^4 km² of southeastern Iran, this study aims to detect and predict regional-scale salt-affected lands. Three sets of Landsat images, each set containing 4 images for 1986, 2000, and 2015 were acquired as the main source of data. Radiometric, atmospheric and cutline blending methods were used to improve the quality of images and help better classify salinized land areas under the support vector machine method. A set of landscape metrics was also employed to detect the spatial pattern of salinized land expansion from 1986 to 2015. Four factors including distance to sea, distance to sea water channels, slope, and elevation were identified as the main contributing factors to land salinization. These factors were then integrated using the multi-criteria evaluation (MCE) procedure to generate land sensitivity map to salinization and also to calibrate the cellular-automata (CA) Markov chain (CA-Markov) model for simulation of salt-affected lands up to 2030, 2040 and 2050. The results of this study showed a dramatic dispersive expansion of salinized land from 7.7 % to 12.7% of the total study area from 1986 to 2015. The majority of areas prone to salinization and the highest sensitivity of land to salinization was found to be in the southeastern parts of the region. The result of the MCE-informed CA-Markov model revealed that 20.3% of the study area is likely to be converted to salinized lands by 2050. The findings of this research provided a view of the magnitude and direction of salinized land expansion in a past-to-future time period which should be considered in future land development strategies.

Full Text

Preamble

J Arid Land <https://doi.org/10.1007/s40333-018-0104-0>

Evaluating and modeling the spatiotemporal pattern of regional-scale salinized land expansion in highly sensitive shoreline landscape of southeastern Iran

Mohammad SHAFIEZADEH, Hossein MORADI*, Sima FAKHERAN
Department of Natural Resources, Isfahan University of Technology, Isfahan
8415683111, Iran

Abstract: This study examines an area of approximately 2.3×10^4 km² in southeastern Iran to detect and predict regional-scale salt-affected lands. Three sets of Landsat images, each containing four images for 1986, 2000, and 2015, were acquired as the primary data source. Radiometric, atmospheric, and cut-line blending methods were used to improve image quality and facilitate better classification of salinized land areas under the support vector machine method. A set of landscape metrics was also employed to detect the spatial pattern of salinized land expansion from 1986 to 2015. Four factors—distance to sea, distance to seawater channels, slope, and elevation—were identified as the main contributing factors to land salinization. These factors were integrated using the multi-criteria evaluation (MCE) procedure to generate a land sensitivity map to salinization and to calibrate the cellular-automata (CA) Markov chain (CA-Markov) model for simulating salt-affected lands up to 2030, 2040, and 2050. The results showed a dramatic dispersive expansion of salinized land from 7.7% to 12.7% of the total study area between 1986 and 2015. The majority of areas prone to salinization and the highest sensitivity of land to salinization were found in the southeastern parts of the region. The MCE-informed CA-Markov model revealed that 20.3% of the study area is likely to be converted to salinized lands by 2050. These findings provide a view of the magnitude and direction of salinized land expansion from past to future time periods, which should be considered in future land development strategies.

Keywords: soil salinization; remote sensing; CA-Markov; salt land expansion; southeastern Iran

1 Introduction

Over the last few decades, land use/land cover (LULC) change has occurred considerably across the world and caused substantial adverse impacts on natural resources and ecosystem functions (Lambin and Geist, 2008). The impacts of LULC change are heavily dependent on the intensity of both natural and anthropogenic driving forces (Meyfroidt et al., 2013), the affected natural resources, and the resilience of ecosystem functions (Wu et al., 2013; Xu and Grumbine, 2014; Houghton and Nassikas, 2017). At local scales, the magnitude

and spatial direction of LULC changes can be properly investigated through field surveying and analyzing low-altitude aerial photography (Hurskainen and Pellikka, 2004; Benedek and Szirányi, 2009), such as peripheral expansion of an urban area. However, it is a major concern to determine how much and to what extent a particular type of LULC conversion occurs regionally or globally and affects ecological functions (Foley et al., 2005; Lambin and Geist, 2008), such as the loss of Arctic sea ice, snow cover, tropical forest lands, continental surface water supplies, and the expansion of salinized land in arid parts of the world.

Substantial progress in space-borne remote sensing (RS) techniques over the last five decades has marked the dawn of a new era for synoptic and periodic observations of the earth's land surface at regional levels (Lillesand et al., 2014). Currently, a considerable number of RS image-capturing systems can collect and transmit huge quantities of data back to earth, providing scientists with opportunities to conduct regional-scale mapping (DeFries et al., 2000; Thenkabail et al., 2009) and detect changes (Kim et al., 2014; McDowell et al., 2015). A large body of these studies took advantage of Landsat imagery data because of its systematic revisiting and acquisition of images from all parts of the earth's surface dating back to 1972, with the launch of Landsat 1 (formerly TIRS-1) and free access to its data archive (Roy et al., 2014; Zhu et al., 2015). Furthermore, multi-temporal RS images are considered the primary data source for most LULC change models (Bhatta, 2010). These models can dynamically simulate and predict future states of LULC classes in a given region according to transition rules and associated driving forces extracted through analyzing multi-temporal LULC maps (Clarke et al., 1997; Eastman, 2012). Under such a context, it is feasible to accurately and realistically estimate LULC changes at large spatial and temporal scales, such as regional land salinization change detection and simulation (Zhou et al., 2012).

Salinization is ranked as one of the most soil-devastating types of LULC change, causing a wide spectrum of adverse economic and environmental impacts (Shrivastava and Kumar, 2015). Currently, approximately 1.0×10^9 hm^2 (7% of the Earth's land surface) is salinized worldwide, mainly due to improper management of irrigated agricultural practices, low precipitation, and high evaporation in arid and semi-arid regions (Rozema and Flowers, 2008; Shrivastava and Kumar, 2015). It is estimated that more than 50% of land resources and arable regions will be significantly affected by salt by 2050, with an annual soil-salinity growth rate of 10% (Jamil et al., 2011). From an economic perspective, soil salinization decreases land productivity for crop production and threatens social economy (Shahbaz and Ashraf, 2013; Shrivastava and Kumar, 2015). Moreover, it can substantially influence global ecological functions and services, such as triggering vast sand and dust storms in arid and semi-arid regions (Moradi, 2016). Fortunately, RS techniques have facilitated a better understanding of the intensity, impacts, and extent of regional-scale land salinization (Metternicht and Zinck, 2003), especially in sub-Saharan Africa and the Middle East, including the study area of this research, where frequent sand and dust storms originating from salinized soil lands have become a major problem

for people living in these regions and beyond (Moradi, 2016).

Wu et al. (2014) investigated the expansion of salt land areas caused by improper agricultural activities in central Iraq and developed a salinized soil mapping methodology to evaluate salinized land expansion from multi-temporal Landsat images. Shalaby and Tateishi (2007) applied multi-temporal Landsat images from 1987 to 2001 to detect large-scale salt land expansion on the coast of northwestern Egypt, which showed continual expansion of salt land in transitional areas between land and seawater. Lin et al. (2006) and Zhou et al. (2012) attempted to simulate the spatial expansion of salt-affected landscapes using LULC change models. These studies substantially explored and modeled the effects of driving forces in the expansion of salinized land areas.

In southern Iran, salinized land expansion has adversely impacted the economy and environment and disturbed human life (Moradi, 2016). However, no attempt has yet been made to estimate the expansion of large-scale salinized land areas in the region. A national report conducted by Moradi (2016) indicated that the southeastern part of Iran has become a major source of dust storms on the shores of the Oman Sea and even Afghanistan. Hence, it is important to map and detect large-scale salinized land expansion and provide a comprehensive picture of salinized land expansion. Multi-temporal Landsat imagery data from 1986, 2000, and 2015 and the cellular-automata (CA) Markov chain (CA-Markov) model were used to predict the ongoing future pattern and expansion of salinized land areas up to 2030, 2040, and 2050.

2.1 Study Area

The study area is approximately 2.3×10^4 km² located in southeastern Iran (25°03' -26°13' N, 47°40' -61°58' E; Fig. 1 [Figure 1: see original paper]). The altitude of the area ranges from 0 to 1300 m in the south at the northern coast of the Oman Sea up to the north. The area is dominated by the Indian monsoon in the east and Mediterranean climate in the west. The annual mean temperature and mean annual precipitation vary significantly across the region.

Bare lands and rocky surface mountains are the main landscapes in the northern study area, while salt lands expand across the south and southeast. The coastline of the study area is considered one of the major mangrove habitats of western Asia, with an area of about 1500 hm² largely covered by *Avicennia marina* sp. (Fig. 1), supporting a great diversity of marine and terrestrial species (Zahed et al., 2010). Although the region is rich in natural resources, temporary agriculture, small-scale oyster/shrimp farming activities, and a small number of residential areas, which are sporadically distributed near the coastline, are the main land use types of the study area, indicating that this region is poorly planned and managed according to national land use planning policies (Moradi, 2016).

Fig. 1 Layout of the study area

2.2 Data

Landsat imagery was used as the primary source of data to delineate salt land expansion in a 30-year time profile. The salinization process in this research referred to any kind of processes that cause the accumulation of soluble salt in surface soils (Hanin et al., 2016). Such a definition includes visual and spectral changes that were detectable by satellite sensors and users. Due to the availability of ancillary data such as GPS-field collected reference points and historical LULC maps, as well as the magnitude of salt land expansion in the study area, Landsat imagery data was selected to explore salinized lands for the years 1986 (TM sensor), 2000 (ETM+ sensor), and 2015 (OLI sensor). Moreover, in each year, four Landsat images with path and row of 158-42, 157-42, 156-42, and 156-43 were acquired to cover the whole extent of the study area (12 tiles in total). Table 1 provides the main characteristics of the Landsat images processed in this research. Furthermore, a 30-m spacing digital topographic model (DEM) map and detailed digital maps of coastline and seawater channel networks from the national project of Makran region development (Moradi, 2016) were employed to produce the sensitivity of land to salinization and to calibrate the LULC change model.

Table 1 Research data used in this study

2.3 Image Processing and Change Detection

The study area lies within a mosaic of four Landsat tiles with different visiting times (different number of rows) using different Landsat sensors, and pre-processing procedures were conducted to prepare the images for classification of salinized land areas. Atmospheric and radiometric corrections were simultaneously performed using the FLAASH (fast line-of-sight atmospheric analysis of hypercube) commercial module implemented in ENVI v5.3® software (Module, 2009). FLAASH is a moderate-resolution atmospheric transmission radiative transfer-based atmospheric correction tool for visible to 3- m spectral range that estimated aerosol/haze-free reflectance values based on the spectral behavior of dark pixels observed by Kaufman et al. (1997) over water and vegetation covers or employing an automated linear regression method over dry soil regions (Cooley et al., 2002; Module, 2009). Input Landsat spectral bands were transformed to radiance values in band-interleaved-by-line (BIL) format (Module, 2009) to execute the FLAASH module. Mid-latitude summer atmospheric and maritime with Kaufman-Tanre aerosol retrieval models were also employed to effectively retrieve column water vapor and aerosol amount of dark-land pixels over the image scenes (Module, 2009). Finally, using additional required parameters from Landsat header files such as flight date and time, as well as the selected atmospheric correction models, all Landsat tiles were transformed to a common atmospheric and radiometric datum.

For image mosaicking, the cutline blending (feathering) technique (Herman et al., 2000) with edge-to-edge polylines crossing the images between salinized

land polygons was utilized to smooth the overlapping area of the images. For a detailed description of image feathering techniques, please refer to Tutorial (2010). The reflectance values of the lookup table derived from the overlapping areas were not significantly different in the images with the equal row number of 42, and their data ranges were not balanced through the histogram matching process. The 156-43 images, which contained a large number of dark pixels located over the Oman Sea, revealed different spectral values than the above-located 156-42 images in the overlapping regions. The spectral values of the 156-43 images were effectively adjusted to the 156-42 image scenes to overcome this problem by assigning a linear histogram stretch value of 4%.

With a set of mosaicked and atmospherically corrected image tiles covering the whole extent of the study area, the support vector machine (SVM) technique was employed to identify and map salinized land areas for 1986, 2000, and 2015. SVM is a common and well-documented supervised classification technique that can obtain accurate classification results, especially under a limited number of training sample units (Mountrakis et al., 2011). Such ability is of great help in LULC classification for large areas where it is not practically possible to collect a broad number of sample units. Approximately 1000 pixels related to salinized land areas and 5000 pixels for other LULC types were collected through interpretation of a set of False Color Composite (FCC) images to train the SVM algorithm. The spectral profile of every training sample pixel was intuitively compared with the spectral library of salinized soils collected by Moradi (2016) through extensive field in-situ measurement in southeastern Iran.

Spectral signatures of salinized soils against Landsat 8 spectral bands are presented in Figure 2 [Figure 2: see original paper], providing a better view of training sample units. The grey part of the graph covered all kinds of mildly to strongly salinized soils across the study area. Although various definitions of soil salinization have been suggested, in line with other remote sensing studies of salinized land identification and classification, this paper relies on the spectral signatures of salinized soils detected through interpretation of the spectral behavior of soils to explore salinized lands. The premise of this approach is based on the observation of spectral signatures illustrated in Figure 2; however, it mainly refers to any kind of processes that cause the accumulation of soluble salt in surface soils (Hanin et al., 2016).

A pixel-based procedure was used to evaluate the accuracy of the classification process based on overall accuracy and Kappa coefficient statistics. Reference sample points were equally distributed between marginal and interior pixels of salinized land polygons to avoid biased accuracy results, especially the effect of positional error (see Lunetta and Lyon, 2004). A small set of landscape metrics was employed, including the number of patches (NP), patch density (PD), mean patch size (MPS), mean nearest neighbor distance (MNN), and largest patch index (LPI). NP and MPS metrics were chosen to investigate the degree of dispersive/organic (edge growth) pattern of salt land expansion. Spatial continuity of salt land expansion was also explored through quantification of PD

and MNN. LPI was used to estimate the proportional growth rate of the largest salt land patch located in the southeastern study area. Table 2 provides a brief explanation of the selected landscape metrics in this research.

Table 2 Description of the selected landscape metrics (McGarigal and Marks, 1995)

2.4 CA-Markov Model

We simulated the ongoing probable picture of salt land expansion according to the immediately preceding spatiotemporal changes of salinized land areas from 1986 to 2015. The MCE (multi-criteria evaluation)-guided CA-Markov land change model implemented in Idrisi Selva (Eastman, 2012) was used to project salinized land expansion up to 2030, 2040, and 2050. The CA-Markov model is a combined dynamic approach that can realistically simulate LULC changes of any given landscape under both biophysical and anthropogenic driving forces (Hyandye and Martz, 2017; Palmate et al., 2017), such as prediction of regional-scale salinized land expansion (Lin et al., 2006; Zhou et al., 2012).

In this model, Markov chain estimates the transition probability of a given state (a LULC class) to any other states without considering neighborhood effects and spatial dimensions (Arsanjani et al., 2013; Brémaud, 2013). A CA model with a standard 5\$×\$5 contiguity filter was integrated (Eastman, 2012) to spatially allocate the probable LULC changes in desired areas (pixels). The CA-Markov model systematically produces a set of suitability maps (also known as conditional maps; El-Hallaq and Habboub, 2015) to express the probability of a land parcel (pixel) of a given LULC class changing to the targeted LULC class. By using these suitability maps, the CA-Markov model often fails to incorporate the effects of driving forces and constraints (Mahiny and Clarke, 2012). However, it was possible for users to calibrate the model by introducing their own best set of suitability maps. In doing so, MCE was used as a well-documented linear combination method in the process of land suitability analysis to effectively prioritize every pixel of a given region based on a set of weighted () and standardized factors (fi) and constraints (Ci) through the following equation:

$$1MCEniiii fwC == \sum \Pi$$

In this research, four transition probability matrices were built using the Markov chain model to estimate the magnitude of salinized land expansion for 2015 (by interpreting changes from 1986 to 2000 (calibration phase)), 2030, 2040, and 2050 (by implementing the 1986 and 2015 maps as the earlier and later LULC maps, respectively (prediction phase)). Four factors—distance to sea, distance to seawater channels, slope, and elevation—were identified as the main driving forces imposing salinized land expansion throughout the study area. The elevation and distance to sea express the potential effects of saline groundwater of the Oman Sea, which controls the upstream freshwater through extensive

dam construction (Moradi, 2016). These two factors substantially represent the effects of groundwater that we were not practically able to map at the regional scale. The seawater channel represents an anthropogenic driving force providing local communities with the opportunity to conduct agricultural activities with saline seawater. The slope is a general factor expressing the sensitivity of soil affected by large-scale salinization. All factor maps were transformed to a scale-independent range of 0 to 255, where values close to 255 indicate higher sensitivity of land to salinization using different types of fuzzy membership functions (Table 3). All factors were compared in pairs, and the corresponding weights were extracted using the analytical hierarchy process method (Saaty, 2008; Eastman, 2012). The weighted and standardized factor maps were linearly combined using Equation 1 to produce the final land suitability map. The resulting map from the MCE procedure was referred to as the land sensitivity map to salinization.

Table 3 Factor maps, fuzzification method, and weighting score used to evaluate the sensitivity of land to salinization

With the land sensitivity map to salinization, the transition probability area matrix of the probable salinized land change between 2000 and 2015, as well as the predefined 5×5 contiguity filter and the 2000 map as the initial land cover map, the CA model was executed to simulate the 2015 salt land expansion using the relative operating characteristic (ROC) metric (Pontius and Schneider, 2001). The resultant simulation map was compared with the reference Landsat-derived salinized land areas to evaluate model performance in simulating salt land expansion. The prediction phase was also performed to predict salt land expansion up to 2030, 2040, and 2050 based on the corresponding transition probability area matrices derived through Markovian analysis and the 2015 map as the initial land cover map.

3 Results

The results of image classification and accuracy assessment are presented in Figure 3 [Figure 3: see original paper] and Table 4. All maps obtained overall accuracy (OA) and Kappa coefficient (KC) of more than 0.94 and 0.90, respectively, indicating acceptable accuracy of the image classification procedure. Based on the results, salinized land areas expanded by more than 1.19×10^5 hm² between 1986 and 2015. In 1986, salinized land areas occupied approximately 1.79×10^5 hm² of the region (7.7% of the total area). With an annual growth rate of 0.28%, salinized land areas reached 2.30×10^5 hm² in 2000 (9.8% of the total area). The study area was dramatically salt-affected between 2000 and 2015, where salinized land areas expanded with an annual growth rate of more than 0.29% and reached 2.98×10^5 hm² in 2015 (12.7% of the total area).

Landscape metric quantification provided detailed information about salinized land expansion in the study area (Table 5). NP, PD, MPS, and LPI increased,

while MNN decreased from 1986 to 2015. The increase in NP indicated the dispersive pattern of salinized land expansion from 1986 to 2015, in which new salt-affected land patches were developed and expanded. Under such a pattern, the pre-existing and newly developed salinized land patches grew rapidly, resulting in a substantial increase in MPS. The decreasing MNN values confirmed the dispersive pattern of salinized land expansion. As represented in Figure 3, the dispersive pattern of salt land expansion was limited to specific regions, and the PD values increased from 1.01 to 1.12. The largest salinized land patch, located in the southeastern study area, also experienced considerable area expansion and merged into large-sized patches, and the LPI value increased dramatically from 1.35 to 11.85 (777.0% increase) from 1986 to 2015.

Table 5 Temporal landscape metric values of salinized land expansion in 1986, 2000, and 2015

The MCE procedure was used to quantitatively assess the sensitivity of land to explore the salinization probability and model calibration. Factor maps of land sensitivity analysis—including distance to sea, distance to seawater channels, slope, and elevation—as well as the corresponding standardized maps, are illustrated in Figure 4 [Figure 4: see original paper]. The map of land sensitivity to salinization was finally produced through a linear combination of the weights and the standardized maps, as illustrated in Figure 5 [Figure 5: see original paper]. The sensitivity of land to salinization varied significantly from south to north. The highest sensitivity scores were detected in the south, southwest, and southeast of the study area where salt land areas had historically expanded. Conversely, sensitivity scores decreased with increasing distance to sea, distance to seawater channels, and elevation (up to 1300 m a.s.l.) towards the northern part of the study area.

The result of model calibration is graphically illustrated in Figure 6 [Figure 6: see original paper]. Black-colored regions with an area of approximately $2.6 \times 10^5 \text{ hm}^2$ (88% of total salinized land area in 2015) denote that the salt-affected regions were accurately predicted. The error areas, which were mistakenly classified through the modeling procedure, were categorized into two classes: salinized land mapped but not predicted (area in green, $1.64 \times 10^4 \text{ hm}^2$; 4.6% of the predicted salinized land areas) and salinized land predicted but not mapped (area in red, $1.21 \times 10^4 \text{ hm}^2$; 3.4% of the predicted salinized land areas). The former class was mainly distributed in the eastern study area, while the latter was located in central and western parts. The salinized land areas were estimated to be $2.0 \times 10^5 \text{ hm}^2$, approximately 7330 hm^2 less than the actual salinized land expansion of 2015, by the 1986–2000 transition area matrix generated through the calibration process. The validity of the MCE-informed CA-Markov model was assessed by comparing the projected and actual salt-affected land generated for 2015. The ROC for the salinized land area was 0.74, indicating good agreement between observed and actual salinized land areas.

The CA-Markov prediction phase was conducted to simulate salinized land ex-

pansion up to 2030, 2040, and 2050 (Fig. 7 [Figure 7: see original paper]). The salt-affected land areas for 2030, 2040, and 2050 were estimated to be around 3.7×10^5 , 4.1×10^5 , and 4.7×10^5 hm² (15.7%, 17.8%, and 20.3% of the total area), respectively. In total, it was estimated that the study area has been and will face more than 3.0×10^5 hm² (12.6% of the total area) of salinization from 1986 to 2050.

4 Discussion

In southeastern Iran, most ecosystem functions and processes have crossed their ecological thresholds and are deeply threatening human life and health in various ways, such as severe water scarcity (Foltz, 2002). Hence, land use managers and decision-makers have been evaluating the suitability of natural land resources to migrate settlers from environmentally devastated ecosystems of central Iran to these regions (Moradi, 2016), especially to the Oman Sea shorelines—the study area. However, this area also faces large-scale soil salinization due to direct and indirect anthropogenic driving forces. A high degree of evaporation, extensive freshwater diversions, and upstream blocking by dams have substantially decreased the humidity of the salinized soils (Moradi, 2016). Frequent storms and strong winds carry a considerable amount of dried salinized dust over the area. According to USA Today, storms originating from this region made Zabol the most polluted city in the world in 2015 (Bacon, 2016), highlighting the magnitude of impact of salinized land expansion in this region.

Processed images successfully revealed the magnitude and spatial trend of salt land expansion in southeastern Iran. The results showed that the major sources of soil salinization were in the southeast (including the largest salinized land patch) and western parts of the study area. Temporal analysis of landscape metric values also suggested a dispersive pattern of salinized land expansion since 1986, with increases in NP and MPS as well as decreases in MNN values. Although following an outward-oriented growth pattern, salt-affected areas were significantly limited to areas near sea shoreline and water channels in low-latitude and gentle-slope lands with increased PD values. The expansion of the largest salinized land patch in the southeastern study area was investigated through quantification of the LPI metric. A substantial increase in LPI values, particularly during the period 2000–2015, was due to the merging of the two largest salinized patches.

A further attempt was made to quantitatively explore the sensitivity of land to salinization in the study area. The assumption behind this effort was based on identifying those factors that control regional-scale soil salinization over wide time spans—i.e., over decades. Zhou et al. (2012) found that the driving forces imposing salinization, both biophysical and human-induced factors, were spatiotemporally dynamic at regional scale. The research by Lin et al. (2006) and Zhou et al. (2012) revealed the ability of the CA-Markov model in simulating and predicting salinized land expansion. This model was employed to project ongoing salinized land expansion for 2030, 2040, and 2050. The results of this

research indicate that southern Iran is experiencing an accelerated rate of salt land expansion. According to past changes delineated in this research, salinized lands will expand significantly and will likely occupy the majority of plain regions. This expansion will form a series of environmental crises. For instance, a very large salt patch predicted to grow in the southeastern study area would have a high capability to trigger regional dust storms. Although the results of this research provide a holistic view into the magnitude and direction of salt land expansion, further attempts are needed to increase the reliability of these predictions. In terms of criteria selection, one should be careful about the temporal variability of driving forces. As Zhou et al. (2012) outlined, driving forces contributing to salt land expansion are highly dynamic and have variable properties that can change even annually. Keeping this matter in mind, it is of high importance to take into account the probability of changes in driving forces as well as the growing speed of salinization for the prediction node and, in turn, for adopting proper land use planning strategies.

5 Conclusions

This study evaluated and modeled the spatial expansion of salt lands in southeastern Iran. The results showed that salt lands have been growing across the entire plain regions of the study area. More importantly, it is predicted that such expansion will lead to the generation of a very large salt patch in the southeastern part with potentially significant regional destructive effects. The analyses carried out in this research were all based on spatial analysis of salt lands quantified from processing Landsat 8 imagery data at three time steps. Furthermore, the predictions were supported by a small set of highly dynamic variables. Despite the successful application of the simulation model, it is recommended to restrict the area under investigation and shorten the prediction spans to better scrutinize how and to what degree the salinized lands are expanding.

Acknowledgments

The authors are pleased to thank the personnel of the Chabahar Department of Environment, especially Mr. Ashrafali HOSSEINI, for providing valuable data and facilitating field surveys, and Mr. Ali ASGARIAN for improving the English writing.

References

- Arsanjani J J, Helbich M, Kainz W, et al. 2013. Integration of logistic regression, Markov chain and cellular automata models to simulate urban expansion. *International Journal of Applied Earth Observation and Geoinformation*, 21: 265-275.
- Bacon J. 2016. The most polluted is? Hint: It' s not in China. *USA Today*. [2016-12-19]. <https://www.usatoday.com/story/news/world/2016/12/19/most-polluted-city-is-not-in-china/95606914/#>.

- Benedek C, Szirányi T. 2009. Change detection in optical aerial images by a multilayer conditional mixed Markov model. *IEEE Transactions on Geoscience and Remote Sensing*, 47(10): 3416–3430.
- Bhatta B. 2010. *Analysis of Urban Growth and Sprawl from Remote Sensing Data*. Berlin: Springer-Verlag Berlin Heidelberg.
- Brémaud P. 2013. *Markov Chains: Gibbs Fields, Monte Carlo Simulation, and Queues*. New York: Springer Science & Business Media, 445.
- Clarke K C, Hoppen S, Gaydos L. 1997. A self-modifying cellular automaton model of historical urbanization in the San Francisco Bay area. *Environment and Planning B*, 24(2): 247–261.
- Cooley T, Anderson G P, Felde G W, et al. 2002. FLAASH, a MODTRAN4-based atmospheric correction algorithm, its application and validation. In: IEEE, *IEEE International Geoscience and Remote Sensing Symposium*. Toronto: IEEE.
- DeFries R S, Hansen M C, Townshend J R G, et al. 2000. A new global 1-km dataset of percentage tree cover derived from remote sensing. *Global Change Biology*, 6(2): 247–254.
- Eastman J R. 2012. *IDRISI Selva*. Worcester: Clark University, 354.
- El-Hallaq M A, Habboub M O. 2015. Using Cellular Automata-Markov Analysis and Multi Criteria Evaluation for Predicting the Shape of the Dead Sea. *Advances in Remote Sensing*, 4(1): 83.
- Foley J A, DeFries R, Asner G P, et al. 2005. Global consequences of land use. *Science*, 309(5734): 570–574.
- Foltz R C. 2002. Iran' s water crisis: cultural, political, and ethical dimensions. *Journal of Agricultural & Environmental Ethics*, 15(4): 357–380.
- Hanin M, Ebel C, Ngom M, et al. 2016. New Insights on plant salt tolerance mechanisms and their potential use for breeding. *Frontiers in Plant Science*, 7.
- Herman J R, Bergen J R, Peleg S, et al. 2000. Method and apparatus for mosaic image construction: Google Patents. [2000-06-13]. <https://www.google.com/patents/US6075905>.
- Houghton R A, Nassikas A A. 2017. Global and regional fluxes of carbon from land use and land cover change 1850–2015. *Global Biogeochemical Cycles*, 31(3): 456–470.
- Hurskainen P, Pellikka P. 2004. Change detection of informal settlements using multi-temporal aerial photographs—the case of Voi, SE-Kenya. In: *Proceedings of the 5th African Association of Remote Sensing of the Environment Conference*, 18–21 October 2004. Nairobi: African Association of Remote Sensing of the Environment.

- Hyandye C, Martz L W. 2017. A Markovian and cellular automata land-use change predictive model of the Usangu Catchment. *International Journal of Remote Sensing*, 38(1): 64-81.
- Jamil A, Riaz S, Ashraf M, et al. 2011. Gene expression profiling of plants under salt stress. *Critical Reviews in Plant Sciences*, 30(5): 435-458.
- Kaufman Y J, Wald A E, Remer L A, et al. 1997. The MODIS 2.1-/spl mu/m channel-correlation with visible reflectance for use in remote sensing of aerosol. *IEEE Transactions on Geoscience and Remote Sensing*, 35(5): 1286-1298.
- Kim D-H, Sexton J O, Noojipady P, et al. 2014. Global, Landsat-based forest-cover change from 1990 to 2000. *Remote Sensing of Environment*, 155: 178-193.
- Lambin E F, Geist H J. 2008. *Land-Use and Land-Cover Change: Local Processes and Global Impacts*. Berlin: Springer Science & Business Media, 221.
- Lillesand T, Kiefer R W, Chipman J. 2014. *Remote Sensing and Image Interpretation*. New York: John Wiley & Sons, 721.
- Lin Z, Zhou D, Liu L. 2006. Regional-Scale Assessment and Simulation of Land Salinization Using Cellular Automata-Markov Model. In: ASABE/CSBE North Central Intersection Meeting. Michigan: American Society of Agricultural and Biological Engineers, RRV12110.
- Lunetta R S, Lyon J G. 2004. *Remote sensing and GIS accuracy assessment*. Florida: CRC Press, 394.
- Mahiny A S, Clarke K C. 2012. Guiding SLEUTH land-use/land-cover change modeling using multicriteria evaluation: towards dynamic sustainable land-use planning. *Environment and Planning B*, 39(5): 925-944.
- McDowell N G, Coops N C, Beck P S, et al. 2015. Global satellite monitoring of climate-induced vegetation disturbances. *Trends in Plant Science*, 20(2): 114-123.
- McGarigal K, Marks B J. 1995. FRAGSTATS: spatial pattern analysis program for quantifying landscape structure. *General Technical Report*. PNW-GTR-351. Portland, USA.
- Metternicht G I., Zinck J A. 2003. Remote sensing of soil salinity: potentials and constraints. *Remote Sensing of Environment*, 85(1): 1-20.
- Meyfroidt P, Lambin E F, Erb K-H, et al. 2013. Globalization of land use: distant drivers of land change and geographic displacement of land use. *Current Opinion in Environmental Sustainability*, 5(5): 438-444.
- Module F. 2009. *Atmospheric Correction Module: QUAC and FLAASH User's Guide* (ver. 4). Boulder: Harris Geospatial Co.
- Moradi H. 2016. *Identification of Environmental Resources and Spatial zoning of Makran Coastal Area, Southeastern Iran*. In: Landuse & Land Cover Change

Report (1st ed.). Department of Environment, Iran.

Mountrakis G, Im J, Ogole C. 2011. Support vector machines in remote sensing: A review. *ISPRS Journal of Photogrammetry and Remote Sensing*, 66(3): 247-259.

Palmate S S, Pandey A, Mishra S K. 2017. Modelling spatiotemporal land dynamics for a trans-boundary river basin using integrated Cellular Automata and Markov Chain approach. *Applied Geography*, 82: 11-23.

Pontius R G, Schneider L C. 2001. Land-cover change model validation by an ROC method for the Ipswich watershed, Massachusetts, USA. *Agriculture, Ecosystems & Environment*, 85(1-3): 239-248.

Roy D P, Wulder M A, Loveland T R, et al. 2014. Landsat-8: Science and product vision for terrestrial global change research. *Remote Sensing of Environment*, 145: 154-172.

Rozema J, Flowers T. 2008. Crops for a salinized world. *Science*, 322(5907): 1478-1480.

Saaty T L. 2008. Decision making with the analytic hierarchy process. *International Journal of Services Sciences*, 1(1): 83-98.

Shahbaz M, Ashraf M. 2013. Improving salinity tolerance in cereals. *Critical Reviews in Plant Sciences*, 32(4): 237-249.

Shalaby A, Tateishi R. 2007. Remote sensing and GIS for mapping and monitoring land cover and land-use changes in the northwestern coastal zone of Egypt. *Applied Geography*, 27(1): 28-41.

Shrivastava P, Kumar R. 2015. Soil salinity: A serious environmental issue and plant growth promoting bacteria as one of the tools for its alleviation. *Saudi Journal of Biological Sciences*, 22(2): 123-131.

Thenkabail P S, Biradar C M, Noojipady P, et al. 2009. Global irrigated area map (GIAM), derived from remote sensing, for the end of the last millennium. *International Journal of Remote Sensing*, 30(14): 3679-3733.

Tutorial E-Z. 2010. ENVI user guide. Colorado Springs, CO: ITT, 590.

Wu K Y, Ye X Y, Qi Z F, et al. 2013. Impacts of land use/land cover change and socioeconomic development on regional ecosystem services: The case of fast-growing Hangzhou metropolitan area, China. *Cities*, 31: 276-284.

Wu W, Mhaimed A S, Al-Shafie W M, et al. 2014. Mapping soil salinity changes using remote sensing in Central Iraq. *Geoderma Regional*, 2-3: 21-31.

Xu J, Grumbine R E. 2014. Building ecosystem resilience for climate change adaptation in the Asian highlands. *Wiley Interdisciplinary Reviews: Climate Change*, 5(6): 709-718.

Zahed M A, Rouhani F, Mohajeri S, et al. 2010. An overview of Iranian mangrove ecosystems, northern part of the Persian Gulf and Oman Sea. *Acta Ecologica Sinica*, 30(4): 240-244.

Zhou D, Lin Z, Liu L. 2012. Regional land salinization assessment and simulation through cellular automaton-Markov modeling and spatial pattern analysis. *Science of the Total Environment*, 439: 260-274.

Zhu Z, Woodcock C E, Holden C, et al. 2015. Generating synthetic Landsat images based on all available Landsat data: Predicting Landsat surface reflectance at any given time. *Remote Sensing of Environment*, 162: 67-83.

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv –Machine translation. Verify with original.