

## Impact of large-scale vegetation restoration project on summer land surface temperature on the Loess Plateau, China Postprint

**Authors:** HE, Guohua, ZHAO, Yong, WANG, Jianhua, WANG, Qingming, ZHU, Yongnan, ZHAO, Yong

**Date:** 2018-10-29T00:00:00+00:00

### Abstract

A large-scale afforestation project has been carried out since 1999 in the Loess Plateau of China. However, vegetation-induced changes in land surface temperature (LST) through the changing land surface energy balance have not been well documented. Using satellite measurements, this study quantified the contribution of vegetation restoration to the changes in summer LST and analyzed the effects of different vegetation restoration patterns on LST during both daytime and nighttime. The results show that the average daytime LST decreased by 4.3°C in the vegetation restoration area while the average nighttime LST increased by 1.4°C. The contributions of the vegetation restoration project to the changes in daytime LST and nighttime LST are 58% and 60%, respectively, which are far greater than the impact of climate change. The vegetation restoration pattern of cropland (CR) converting into artificial forest (AF) has a cooling effect during daytime and a warming effect at nighttime, while the conversion of CR to grassland has an opposite effect compared with the conversion of CR to AF. Our results indicate that increasing evapotranspiration caused by the vegetation restoration on the Loess Plateau is the controlling factor of daytime LST change, while the nighttime LST change is affected by soil humidity and air humidity.

### Full Text

#### Preamble

#### Impact of Large-Scale Vegetation Restoration Project on Summer Land Surface Temperature on the Loess Plateau, China

HE Guohua, ZHAO Yong\*, WANG Jianhua, WANG Qingming, ZHU Yongnan  
State Key Laboratory of Simulation and Regulation of Water Cycle in River

Basin, China Institute of Water Resources and Hydropower Research, Beijing 100038, China

**Abstract:** A large-scale afforestation project has been carried out since 1999 in the Loess Plateau of China. However, vegetation-induced changes in land surface temperature (LST) through the changing land surface energy balance have not been well documented. Using satellite measurements, this study quantified the contribution of vegetation restoration to the changes in summer LST and analyzed the effects of different vegetation restoration patterns on LST during both daytime and nighttime. The results show that the average daytime LST decreased by 4.3°C in the vegetation restoration area while the average nighttime LST increased by 1.4°C. The contributions of the vegetation restoration project to the changes in daytime LST and nighttime LST are 58% and 60%, respectively, which are far greater than the impact of climate change. The vegetation restoration pattern of cropland (CR) converting into artificial forest (AF) has a cooling effect during daytime and a warming effect at nighttime, while the conversion of CR to grassland has an opposite effect compared with the conversion of CR to AF. Our results indicate that increasing evapotranspiration caused by the vegetation restoration on the Loess Plateau is the controlling factor of daytime LST change, while the nighttime LST change is affected by soil humidity and air humidity.

**Keywords:** vegetation restoration project; summer; land surface temperature; climate change; Loess Plateau

## Introduction

Arid and semi-arid regions cover 41% of the Earth's land surface, with 38% of the human population living in these regions. These areas are ecologically vulnerable and sensitive to climate change and human activity, which can affect human health (Reynolds et al., 2007; Yang et al., 2015, 2016). A series of large-scale ecological restoration projects have been carried out worldwide to address potentially serious environmental and ecological issues, and China has also made significant efforts in recent decades (Yi et al., 2009; Allan et al., 2013). In 1999, the Chinese government launched the Grain for Green (GFG) program, a large-scale vegetation restoration project that represents the largest eco-environmental construction project and most important nature conservation program in the arid and semi-arid regions of China (Liu et al., 2008; Feng et al., 2013). GFG involves returning croplands on steep slopes to forest lands and grasslands through a government payment scheme, which engages millions of rural households as core agents of project implementation (Lü et al., 2012; Feng et al., 2013). As the core region of GFG, the Loess Plateau is an ecologically vulnerable area sensitive to climate factors (e.g., periodic heavy rainfall) and intensified human activities (e.g., over-grazing and coal mining) (Zhang et al., 2008; Liang et al., 2013). However, the vegetation coverage of the Loess Plateau has gradually improved since the implementation of GFG (Xin et al., 2008; Zhou et al., 2009), and over  $6 \times 10^4$  km<sup>2</sup> of croplands have been converted into forest

lands and grasslands during 1999–2010 (Li et al., 2016).

Numerous studies have attempted to evaluate the impact of the vegetation restoration project on the Loess Plateau from different perspectives. For example, Gao et al. (2017) used the Budyko framework to simulate the water cycle on the Loess Plateau during 1990–2014. With an increase in vegetation coverage, both regional precipitation and actual evapotranspiration showed a significant increasing trend, and the increasing rate of precipitation was greater than evapotranspiration. Deng et al. (2016) found that the storage of soil carbon increased by about 21.3% in the topsoil (0–20 cm) of the re-vegetated lands. Zhai et al. (2015) investigated the relationship between vegetation restoration and land surface albedo on the Loess Plateau for the period 2000–2010 and concluded that human-induced land use change is the control factor for albedo reduction. In addition, some studies discussed the effects of vegetation change on energy and material cycles from different perspectives, including energy budgets (Chapin et al., 2005; McVicar et al., 2007), soil organic matter (Xiao et al., 2017), soil humidity (Jian et al., 2015), and soil erosion (Zhou et al., 2006; Zhang et al., 2016). Based on these studies, we conclude that the vegetation restoration project has greatly affected the environment and material recycling processes on the Loess Plateau. However, the changes in land surface temperature (a dominant factor in fragile ecological regions) caused by the increased vegetation coverage have not been well documented.

As the direct driving force in the exchange of long-wave radiation and turbulent heat fluxes at the surface-atmosphere interface, land surface temperature (LST) is one of the most important parameters in the physical processes of surface energy and water balance at scales from local to global (Anderson et al., 2008; Karnieli et al., 2010). On the one hand, changes in LST will directly affect soil moisture content and soil evaporation, thereby indirectly affecting regional vegetation growth, agricultural production, and the water cycle (Zhang et al., 2014; Harris et al., 2017). On the other hand, LST is an important indicator for predicting drought, climate change, and earthquakes (Choudhury et al., 2007; Amiri et al., 2009; Vancutsem et al., 2010). LST is the joint outcome of sensible heat and latent heat, which is influenced by albedo, evapotranspiration, surface roughness, and other factors (Xue et al., 2017). According to Peng et al. (2014), vegetation restoration changes the albedo and thus changes the amount of absorbed solar radiation at the ground surface. At the same time, vegetation restoration generally increases evapotranspiration (ET) and causes a cooling effect (Jackson et al., 2008). Therefore, vegetation restoration may result in both cooling and warming effects, which is related to the vegetation restoration pattern and latitude (Lee et al., 2011). The biophysical effects of vegetation restoration on local climate can be much larger than the small global cooling effect resulting from the uptake of CO<sub>2</sub> by growing vegetation (Lee et al., 2011). However, these biophysical effects are also complex and depend on “background” climate (Wang et al., 2011). In general, although the large-scale vegetation restoration project has been conducted for more than 15 years, there is a complex relationship between vegetation restoration and LST, and as such,

the effects of vegetation change on LST on the Loess Plateau are not clear and need to be further elucidated.

In this study, we investigated how vegetation restoration affects LST on the Loess Plateau using satellite-derived LST datasets from MODIS instruments in summer (June to August) during 2000 to 2013. These LST data depend on the radiative properties of the land surface (Wan, 2008; Zhou et al., 2012) and therefore have a larger diurnal amplitude than observation data from meteorological stations (Jin and Dickinson, 2010). The objectives of this study are: (1) to reveal the characteristics and controlling factors of LST change in the vegetation restoration area; and (2) to quantify the effects of different vegetation restoration patterns on LST during both daytime and nighttime.

## 2.1 Study Area

The Jinghe River Basin, a small region with an area of  $4.54 \times 10^4$  km<sup>2</sup>, was chosen as the study area to minimize the effect of altitude and precipitation on LST. The basin is located in the southern Loess Plateau of China (34°14'–38°10' N, 105°49'–108°58' E; Fig. 1 [Figure 1: see original paper]), dominated by a semi-arid continental monsoon climate. During the period 1980–2015, the annual summer precipitation varied from 190 mm in the northwest to 250 mm in the southeast part of the basin, with a mean value of 225 ( $\pm 30$ ) mm. It has a typical water-limited landscape, with actual evapotranspiration accounting for 85% of the precipitation; this percentage increases along the precipitation gradient (Jin and Dickinson, 2010). The main soil types in the basin are Loessial soil and dark Loessial soil, which form thick soil layers (50–80 m), loose in structure, easily collapsed and eroded. The land-use types are primarily cropland, grassland, forest, and sandy land. In the Jinghe River Basin, a large number of ecological restoration projects have been implemented in the past 30 years since the 1980s. However, these costly efforts yielded little success during the first 20 years (Peng et al., 2015). In the past 10 years, vegetation restoration has begun to achieve certain effects, and the average summer NDVI has increased from 0.4 in 2000 to 0.7 in 2013.

Fig. 1 Location map and change in NDVI in the study area. (a) NDVI in 2000, (b) NDVI in 2013, and (c) NDVI change from 2000 to 2013.

## 2.2 Data

We used the MODIS collection-3 products downloaded from NASA (<https://www.nasa.gov/topics/earth/index.html>) for LST, surface albedo, and ET. These products have been used extensively in a variety of areas and are proven to be of high quality (Wan, 2008; Zhou et al., 2012). For LST, we used the 8-day average LST from Terra-MODIS (MOD11A2) products in summer (June to August) from 2000 to 2013. The LST data cover both daytime (10:30 am) and nighttime (22:30 pm). In this study, we corrected the noise resulting from cloud contamination and topographic differences, and only

pixels with LST error  $<1$  K were used. The MODIS surface albedo products (MCD43A3) include white sky albedo and black sky albedo, with a 0.5-km spatial resolution and a 16-day time resolution from 2000 to 2013. We used the white sky albedo shortwave broadband in this study, and the bias of the MODIS albedo was mostly less than 5% (Liu et al., 2009). The MODIS ET data (MOD16A2) uses land cover, albedo, air temperature, radiation, humidity, and leaf area index as input data, with a 0.5-km spatial resolution and an 8-day interval. The mean bias of the MODIS ET is 0.3 mm/d compared to ET observations from the eddy flux tower (Mu et al., 2011).

The NDVI data and land cover map (2000, 2005, 2010 and 2013) at 1-km resolution were downloaded from the Resource and Environment Data Platform (<http://www.resdc.cn/DataList.aspx>). The NDVI data come from vegetation sensor with monthly interval and land cover data come from Landsat TM and Landsat ETM. In this study, we adopted the China land resource classification system for the land cover data (Zhang et al., 2010). Meteorological data, such as relative air humidity, observed land surface temperature, and other data needed to calculate the net radiation were collected from six national weather stations. The relative soil humidity was collected from 57 national agricultural meteorological stations in the study area from 2000 to 2013. Meteorological data and relative soil humidity data were downloaded from the Meteorological Data Center of China Meteorological Administration (<http://data.cma.cn/>).

### 2.3 Methods

In the study area, we defined the region where the average NDVI increased by more than 0.2 during 2000 to 2013 as the vegetation restoration area (VRA) (blue regions in Fig. 1c) (Cao et al., 2017) and the area where the NDVI change was less than 0.2 as the non-afforestation area (NAA) (yellow regions in Fig. 1c). In addition, irrigated areas, mountains (slope  $>15^\circ$ ; Deng et al., 2016), and urban areas were excluded (the areas of VRA and NAA account for 52% and 46% of the basin area, respectively). The average elevation difference between VRA and NAA is 178 m, less than the calculation limit of 200 m (Peng et al., 2014).

Because the LST in VRA was under the influence of vegetation restoration and climate, the LST change in NAA without the influence of vegetation was set as LST\_climate, the LST in VRA as LST\_VRA, and the effect of vegetation on LST as LST\_vegetation. The relationship between LST\_VRA and LST\_vegetation is shown as Equation 1. (1) Remote LST was defined as the average of the daytime LST and nighttime LST. All daily meteorological data were calculated to obtain a monthly average, which was finally calculated to obtain the seasonal average data for each year. The changing trend in the time series data was analyzed with the Mann-Kendall method.

### 3.1 LST Change in VRA

The LST in VRA significantly decreased during the summer and exhibited a clear correlation with NDVI (correlation coefficient=0.7). We can clearly see in Figure 2a [Figure 2: see original paper] that the average NDVI in VRA increased from 0.40 in 2003 to 0.67 in 2013. With the increase in NDVI, summer LST in VRA decreased from 25.2°C in 2000 to 23.5°C in 2013. LST is the average value of daytime LST and nighttime LST, and these two factors together determine the changes in LST. The Mann-Kendall trend analysis indicates that the daytime LST significantly decreased from 2000 to 2013 (Test  $Z=-2.29$ , 95% confidence level), and the average temperature decrease was 0.3°C/a.

The nighttime LST slightly increased during the same period (Test  $Z=1.20$ , did not pass the significance test of 95%), with an average rise of 0.1°C/a. The daytime cooling effect is significantly higher than the nighttime warming effect in VRA, meaning that the significant decreasing trend in LST from 2000 to 2013 is mainly attributed to the daytime LST change. We can also see from Figures 2b and c that the daytime LST and nighttime LST in VRA both showed large fluctuations from 2000 to 2013, which may be attributed to climate change and the low time resolution (8-day) of the LST product.

### 3.2 Effects of Vegetation Restoration on LST

The LST change in VRA is affected by vegetation restoration and climate factors. To reveal the impact of vegetation change on LST, we assume that the LST in NAA is only affected by climatic factors, and LST\_climate is used to reflect the influence of climatic factors on the LST of VRA. Based on remote LST data, we plotted the inter-annual variation curve of LST\_VRA, LST\_climate, and LST\_vegetation in the summer from 2000 to 2013 (Figs. 3a and b).

In daytime, LST\_climate shows a slight downward trend (Test  $Z=-0.99$ , did not pass the significance test of 95%), with an average decrease of 0.124°C/a. However, LST\_VRA decreased significantly during the same period, with an average decrease of 0.311°C/a, which is far greater than the annual decrease trend of LST\_climate. These results suggest that the significant decreasing trend in daytime LST\_VRA could be mainly attributed to vegetation restoration, and this cooling effect became stronger with the increase in NDVI (Fig. 3a [Figure 3: see original paper]). Based on Equation 1, LST\_vegetation experienced an extremely significant increase from -3.99°C in 2000 to -6.13°C in 2013 (Test  $Z=3.28$ , 99% confidence level), and the cumulative cooling effect exceeded 2.5°C in summer from 2000 to 2013 (Fig. 3a). By comparing the effect of LST\_vegetation and LST\_climate on LST\_VRA, we conclude that vegetation restoration contributed 58% of the decrease in daytime LST\_VRA, which was far greater than the contribution by climate change (42%).

At nighttime, both LST\_VRA and LST\_climate show a slight upward trend from 2000 to 2013, and the upward trend in LST\_VRA is greater than that in LST\_climate. This result means that vegetation change helps increase the night-

time LST in VRA. Based on Equation 1, vegetation change contributes a little to the warming effect at nighttime, with an average increase of  $0.06^{\circ}\text{C}/\text{a}$  during 2000-2013 (Fig. 3b). Based on the trend in LST\_vegetation, LST\_climate, and LST\_VRA, we can also conclude that vegetation restoration contributed 60% of the increase in nighttime LST\_VRA.

Fig. 3 Summer LST\_VRA, LST\_vegetation, and LST\_climate change in vegetation restoration area during 2000-2013. (a) daytime and (b) nighttime.

### 3.3 Effects of Different Land Cover Types on LST

Converting slope cropland (CR) into artificial forest (AF) and grassland (GR) are the main patterns of vegetation restoration on the Loess Plateau. This study used remote LST data and land cover data to analyze the impact of CR, AF, and GR on daytime and nighttime LST in VRA (Fig. 4 [Figure 4: see original paper]). The average LST of AF was significantly lower than those of CR and GR during the daytime, at about  $2.2^{\circ}\text{C}$  between AF and CR and  $2.4^{\circ}\text{C}$  between AF and GR. The difference in daytime LST between CR and GR is small; however, the LST of GR is always higher than that of CR, with an average gap of  $0.21^{\circ}\text{C}$ . During nighttime, although the differences in LST among AF, CR, and GR are only  $0.07^{\circ}\text{C}$ - $0.63^{\circ}\text{C}$ , the nighttime LST of AF is always slightly higher than those of CR and GR, with differences of  $0.56^{\circ}\text{C}$  and  $0.63^{\circ}\text{C}$ , respectively. The land surface temperature of different land cover is affected by both climate and vegetation factors. According to Equation 1, the average LST of AF is  $-5.6^{\circ}\text{C}$ , which is significantly smaller than those of GR ( $-3.2^{\circ}\text{C}$ ) and CR ( $-3.4^{\circ}\text{C}$ ). This indicates that AF has a stronger cooling effect than CR and GR in daytime.

Although AF, CR, and GR all show a small warming effect at nighttime, the warming effect of AF ( $0.70^{\circ}\text{C}$ ) is always greater than those of CR ( $0.12^{\circ}\text{C}$ ) and GR ( $0.05^{\circ}\text{C}$ ). Compared with other years, the LST varied significantly under different land covers during daytime and nighttime in 2005. This is because the LST in 2005 had larger fluctuations than in other years, which is mainly attributed to climate factors (Fig. 3).

The LST changes of daytime and nighttime in different vegetation restoration patterns were analyzed in this study (Fig. 5 [Figure 5: see original paper]). For CR-AF (conversion from CR to AF), the cooling effect was observed in the daytime and the warming effect occurred at nighttime, which is consistent with the result of the change in LST in VRA. However, for CR-GR (conversion from CR to GR), the diurnal change in LST has the opposite effect compared with CR-AF. In daytime, the cooling effect of LST of CR-AF is  $2.2^{\circ}\text{C}$ . In contrast, the average annual nighttime LST between CR and AF was  $0.2^{\circ}\text{C}$ , indicating that the vegetation restoration of CR-AF warms the land surface during nighttime but with a smaller magnitude than the daytime cooling effect. The warming and cooling effect of GR are always lower than those of CR (Fig. 4), so the vegetation restoration of CR-GR increases LST during the day ( $0.7^{\circ}\text{C}$ ) and decreases LST

at nighttime ( $0.1^{\circ}\text{C}$ ).

#### 4.1 Influence Factors of LST Change in Daytime and Nighttime

Vegetation restoration has a cooling effect in daytime and a warming effect at nighttime in VRA (Fig. 2). The asymmetric diurnal cycle of LST results from different energy balance processes (Dai et al., 1999; Zhou et al., 2007). LST is determined by sensible heat (Allan et al., 1998); therefore, LST can be expressed as a function of sensible heat (H) in the study area (Eq. 2). (2) Based on the surface energy balance equation, H is given as Equation 3. (3) where H is the sensible heat flux ( $\text{MJ}/(\text{m}^2 \cdot \text{d})$ ), ET (mm) and G ( $\text{MJ}/(\text{m}^2 \cdot \text{d})$ ) are the evapotranspiration and soil heat flux, respectively, and L is the latent heat of vaporization of water, which is a constant (L is about  $2.45 \text{ MJ}/\text{kg}$ ) (Allan et al., 1998).  $R_n$  is the net radiation ( $\text{MJ}/(\text{m}^2 \cdot \text{d})$ ).  $R_n$  was calculated using the Penman-Monteith equation: (4) where  $R_{ns}$  and  $R_{nl}$  are net short-wave radiation ( $\text{MJ}/(\text{m}^2 \cdot \text{d})$ ) and net long-wave radiation ( $\text{MJ}/(\text{m}^2 \cdot \text{d})$ ), respectively.  $R_{ns}$  and  $R_{nl}$  can be given as Equations 5 and 6 (Allan et al., 1998). (5) (6) The meanings and values of each coefficient in Equations 5 and 6 can be found in Liu and McVicar (2012). The albedo ( $\alpha$ ) can be set as a different value for different NDVI (Zhai et al., 2015), and changes in the albedo will further affect  $R_n$  and LST (Peng et al., 2014). Therefore, based on MODIS albedo data, we used the annual summer surface albedo from 2000 to 2013 to reflect the effect of vegetation change on LST.

Daytime LST is mainly controlled by  $R_n$  and  $L \times ET$  (Eq. 3;  $R_n$  and  $L \times ET$  are much larger than G in daytime and this study ignores the effect of G on sensible heat flux) (Allan et al., 1998). Summer extra-terrestrial solar radiation is assumed to be similar in VRA during 2000–2013 ( $17.94\text{--}21.91 \text{ MJ}/(\text{m}^2 \cdot \text{d})$ ). Hence, the surface albedo determines the amount of net radiation under the same climate conditions (Eqs. 5 and 6). According to Zhai et al. (2015), there is a significant negative correlation between NDVI and albedo; thus, vegetation restoration will increase  $R_n$  and bring warming effect in daytime (Eqs. 5 and 6). During the same time, the increase in vegetation will also increase evapotranspiration and bring a cooling effect (Zhai et al., 2015). So, the effect of vegetation change on daytime LST is determined by both the warming effect (caused by albedo) and cooling effect (caused by ET). Based on MODIS data, this study analyzed the change in albedo and ET in the VRA from 2000 to 2013. With the increase in vegetation, ET in summer showed an extremely significant increase ( $Z=4.38$ , 99% confidence level), and the annual increase trend is  $0.1 \text{ mm}/\text{d}$  (Fig. 6a [Figure 6: see original paper]). During the same period, the albedo significantly decreased from 0.193 to 0.152 ( $Z=-4.05$ , 99% confidence level), a decrease of 21% and an average annual decrease of 0.0023 (Fig. 6b). The effects of albedo and ET change on heat flux are calculated based on Equations 3–6 (Fig. 7 [Figure 7: see original paper]). We can clearly see in Figure 7 that vegetation restoration led to an increase in both net radiation and latent

heat in VRA; however, the increase trend in latent heat is significantly greater than the net radiation, and the gap between latent heat and net radiation gradually expands, indicating that the cooling effect caused by the ET increase is greater than the warming effect caused by the albedo decrease. Thus, ET is the control factor of daytime LST change in VRA.

In general, land surfaces absorb and store energy from the atmosphere during the day and release energy during nighttime. ET is negligible in nighttime and LST is closely related to the energy stored during daytime and the near-surface atmosphere boundary layer (Wan, 2008; Lee et al., 2011). Based on observation data from national weather stations and agricultural meteorological stations, the change in relative soil humidity and relative air humidity during 2000 to 2013 was analyzed. During this period, the relative soil humidity in VRA increased from 58% to 67%, with an annual average increase of 0.48% (Fig. 8a [Figure 8: see original paper]). Increased soil humidity resulted in more daytime heat storage and more nighttime heating. The increase in air humidity near the surface means more downward long-wave radiation received from the atmosphere and a reduction in the outgoing long-wave radiation from the land surface (Fig. 8b). This long-wave radiative imbalance has a stronger effect during nighttime when the boundary layer is thinner and more stable (Zhou et al., 2009).

## 4.2 Influence Factors of Vegetation Restoration Patterns on LST

Different afforestation patterns have different effects on ET and albedo (Liu et al., 2009). The ET of AF is indeed higher than that of CR in 50% of the VRA and higher than that of GR in 68% of the VRA, while the albedo is lower in 8% of CR and 9% of GR, respectively (Fig. 9 [Figure 9: see original paper]). This clearly indicates that AF across the VRA absorbs more incoming radiation and dissipates more energy as latent heat. Thus, artificial forest induces cooling during daytime. The surface incoming solar radiation in summer is between 17.94 and 21.91 MJ/(m<sup>2</sup>·d) on the Loess Plateau. The average difference in albedo is 0.014 between CR and AF, and 0.003 between GR and CR. Hence, the extra solar energy absorbed by the change in vegetation cover from CR to AF (CR-AF) is 0.25-0.31 MJ/(m<sup>2</sup>·d), while the solar energy dissipated by the change from CR to GR (CR-GR) is -0.05 to -0.07 MJ/(m<sup>2</sup>·d). The ET change resulting from these two vegetation restoration patterns is 0.72 and -0.16 mm/d respectively, which is equal to an extra energy dissipation through latent heat of about 1.76 and -0.39 MJ/(m<sup>2</sup>·d), respectively. So, based on Equations 3-6, vegetation restoration patterns of CR-AF will result in a cooling effect during the day due to more energy dissipation, while CR-GR will result in a warming effect due to more energy absorption.

The impacts of different land use types on soil humidity and air humidity are different at nighttime. According to Xu et al. (2008), under the same climatic conditions, the relative soil humidity of AF in the study area is about equal to

CR, which is 1.5 to 2.0 times that of GR. The relative air humidity (150 cm above vegetation) of AF is 13% more than CR in summer, while the relative air humidity of CR and GR is similar (Li et al., 2010). This result indicates that CR-AF will increase the ability to absorb radiation (increase in air humidity), while CR-GR will decrease the ability to store heat (decrease in soil humidity). Thus, the vegetation restoration of CR-AF increases LST at nighttime, while CR-GR reduces LST. In addition, the nighttime warming effect of CR-AF could be magnified as a result of reduced atmosphere turbulence from a more stable stratification over vegetation (Lee et al., 2011). This reduces the heat dissipation from AF cover types compared with the open lands of CR and GR.

### 4.3 Uncertainty Analysis

The effect of vegetation on LST between the Jinghe River Basin and other regions is compared to evaluate the accuracy of the calculated results (Table 1). Table 1 shows that our results match well with Hao et al. (2016), Peng et al. (2014) and Potchter et al. (2008). However, in the Tibetan Plateau, vegetation greening increases the local LST due to the huge decrease in albedo (Shen et al., 2015). In general, effects of vegetation cover on LST vary with the vegetation restoration pattern and latitude (Lee et al., 2011).

Table 1 Comparison between results from other studies

Location	Period	Vegetation type	Average LST change (°C)	Source
Jinghe River Basin		Restoration	-1.65	This study
Tarim River Basin		Natural	-3.02	Hao et al. (2016)
China		Restoration	-0.90	Peng et al. (2014)
Southern Israel		Natural	-2.00	Potchter et al. (2008)
Tibetan Plateau		Natural	+0.78	Shen et al. (2015)

To test the rationality of MODIS LST data, the study analyzed the remote LST data and observed LST data from six national weather stations in VRA during 2000 to 2013. The remote sensing data has a good linear correlation with the observed data (Fig. 10a [Figure 10: see original paper]). Although the MODIS data is usually 0.68°C-1.02°C lower than the actual LST, they both exhibit a

similar trend. Therefore, the MODIS data in VRA can meet the requirements of the study.

Fig. 10 Scatterplot of uncertainty analysis. (a) Remote LST and observed LST and (b) estimated Rn and observed Rn.

In our research, we obtained the summer Rn from the Penman-Monteith equation. The observed data from the weather station (there is only one weather station in VRA that can observe the net radiation and observation time starts from 2006) was analyzed to test the rationality of the estimated Rn. Figure 10b shows that the correlation coefficient between the estimated Rn and the observed Rn reached 0.81, and the error between observed data and estimated data is 4%–9%. Therefore, the estimated Rn can meet the requirement of the study.

Although many factors are considered, this study still has limitations regarding the effects of vegetation changes on LST. For example, vegetation greening can lead to the increase in surface roughness and more solar energy could be absorbed by the surface accordingly. Our results may overestimate the cooling effect caused by vegetation change. This work also ignores the effect of human activities on LST, such as irrigation. In addition, it must be noted that the Loess Plateau is a very large and geodiverse region; thus, the research results in the Jinghe River Basin may not apply to the entire Loess Plateau.

Vegetation restoration in the Jinghe River Basin has gradually reduced the LST in summer. Because the temperature of the deep soil is relatively stable, the decrease in LST will directly reduce the soil temperature gap, thus reducing soil evaporation and increasing the soil moisture content (Kidron and Kronenfeld, 2016; Ren et al., 2017). Increased soil moisture caused by vegetation change is an important reason for reduced runoff on the Loess Plateau (Fu et al., 2003), and this could lead to uncertainty in regional water resources management as well (Feng et al., 2016).

## 5 Conclusions

Based on the remote LST data and land cover data, we concluded that the cooling effect and warming effect of PF, CR and GR are gradually decreasing. Therefore, the vegetation restoration pattern of CR-PF resulted in the cooling effect in the daytime and the warming effect at nighttime, while the vegetation restoration pattern of CR-GR has the opposite effect.

Based on the surface energy balance equation, this paper discussed the effect of vegetation restoration on energy budget. With vegetation change, net radiation (inflected by albedo) and latent heat (inflected by ET) in VRA show an increasing trend; however, the average increase trend of latent heat ( $-2.4 \text{ MJ}/(\text{m}^2 \cdot \text{d})$ ) was significantly greater than the net radiation ( $0.09 \text{ MJ}/(\text{m}^2 \cdot \text{d})$ ). Thus, the daytime LST in VRA was decreased. For vegetation restoration patterns of CR-AF and CR-GR, CR converted into AF means that about  $1.5 \text{ MJ}/(\text{m}^2 \cdot \text{d})$

d) of energy is released, while CR converted into GR means that about 0.3 MJ/(m<sup>2</sup> · d) of energy is absorbed. Therefore, vegetation restoration pattern of CR-AF decreased LST during the day, while CR-GR increased LST. The increase of nighttime LST in VRA can be attributed to the increase of relative soil humidity and relative air humidity.

Although many factors are considered in this study for LST attribution analysis, there are still some limitations. It should be noted that the Loess Plateau is a very large and geodiverse region; as such, the research results in the Jinghe River Basin may not apply to the entire Loess Plateau.

**Acknowledgements** This work was funded by the National Key Research and Development Program of China (2016YFC0401306), the National Science Fund for Distinguished Young Scholars (51625904), and the International Science & Technology Cooperation Program of China (2016YFE0102400). The authors are grateful to the anonymous reviewers for their detailed comments.

## References

- Allan J D, McIntyre P B, Smith S D P, et al. 2013. Joint analysis of stressors and ecosystem services to enhance restoration effectiveness. *Proceedings of the National Academy of Sciences of the United States of America*, 110(1): 372-377.
- Allan R G, Pereira L S, Raes D, et al. 1998. Crop evapotranspiration-Guidelines for computing crop water requirements-FAO irrigation and drainage paper 56. FAO. Rome, Italy.
- Amiri R, Weng Q H, Alimohammadi A, et al. 2009. Spatial-temporal dynamics of land surface temperature in relation to fractional vegetation cover and land use/cover in the Tabriz urban area, Iran. *Remote Sensing of Environment*, 113(12): 2603-2615.
- Anderson M C, Norman J M, Kustas W P, et al. 2008. A thermal-based remote sensing technique for routine mapping of land-surface carbon, water and energy fluxes from field to regional scales. *Remote Sensing of Environment*, 112(12): 4227-4241.
- Cao Z, Li Y R, Liu Y C, et al. 2017. When and where did the loess plateau turn 'green' ? Analysis of the tendency and breakpoints of the normalized difference vegetation index. *Land Degradation & Development*, 29(1): 162-175.
- Chapin III F S, Sturm M, Serreze MC, et al. 2005. Role of land-surface changes in arctic summer warming. *Science*, 310(5748): 657-660.
- Choudhury S, Panda S K, Saraf A K, et al. 2007. MODIS land surface temperature data detects thermal anomaly preceding 8 October 2005 Kashmir earthquake. *International Journal of Remote Sensing*, 28(20): 4587-4596.

- Dai A, Trenberth K E, Karl T R. 1999. Effects of clouds, soil moisture, precipitation, and water vapor on diurnal temperature range. *Journal of Climate*, 12: 2451-2473.
- Deng J, Sun P S, Zhao F Z, et al. 2016. Soil C, N, P and its stratification ratio affected by artificial vegetation in subsoil, loess plateau china. *PLoS ONE*, 11(3): e0151446.
- Feng X, Fu B, Lu N, et al. 2013. How ecological restoration alters ecosystem services: an analysis of carbon sequestration in China's Loess Plateau. *Scientific Reports*, 3: 2846.
- Feng X M, Fu B J, Piao S L, et al. 2016. Revegetation in China's loess plateau is approaching sustainable water resource limits. *Nature Climate Change*, 6: 1019-1022.
- Fu B, Wang J, Chen L D, et al. 2003. The effects of land use on soil moisture variation in the Danangou catchment of the Loess Plateau, China. *Catena*, 54(1-2): 197-213.
- Gao X R, Sun M, Zhao Q, et al. 2017. Actual ET modelling based on the Budyko framework and the sustainability of vegetation water use in the loess plateau. *Science of the Total Environment*, 579: 1550-1559.
- Hao X M, Li W, Deng H J. 2016. The oasis effect and summer temperature rise in arid regions - case study in Tarim Basin. *Scientific Reports*, 6: 35418.
- Harris P P, Folwell S S, Gallego-Elvira B, et al. 2017. An evaluation of modeled evaporation regimes in Europe using observed dry spell land surface temperature. *Journal of Hydrometeorology*, 18: 1453-1470.
- Jackson R B, Randerson J T, Canadell J G, et al. 2008. Protecting climate with forests. *Environmental Research Letters*, 3(4): 044006.
- Jian S Q, Zhao C Y, Fang S M, et al. 2015. Effects of different vegetation restoration on soil water storage and water balance in the Chinese Loess Plateau. *Agricultural and Forest Meteorology*, 206: 85-96.
- Jin M L, Dickinson R E. 2010. Land surface skin temperature climatology: benefitting from the strengths of satellite observations. *Environmental Research Letters*, 5(4): 44004.
- Karnieli A, Panov N, Goldberg A, et al. 2010. Use of NDVI and land surface temperature for drought assessment: merits and limitations. *Journal of Climate*, 23: 618-633.
- Kidron G J, Kronenfeld R. 2016. Temperature rise severely affects pan and soil evaporation in the Negev Desert. *Ecohydrology*, 9(6): 1130-1138.
- Lee X, Goulden M L, Hollinger D Y, et al. 2011. Observed increase in local cooling effect of deforestation at higher latitudes. *Nature*, 479: 384-387.

- Li Q Y, Yu X X, Xin Z B, et al. 2010. Analysis on soil physical properties of different land uses in a typical watershed of the Loess Plateau. *Research of Soil and Water Conservation*, 17(6): 106-110, 114. (in Chinese)
- Li S, Liang W, Fu B, et al. 2016. Vegetation changes in recent large-scale ecological restoration projects and subsequent impact on water resources in China' s Loess Plateau. *Science of the Total Environment*, 569-570: 1032-1039.
- Liang K, Liu C M, Liu X M, et al. 2013. Impacts of climate variability and human activity on streamflow decrease in a sediment concentrated region in the Middle Yellow River. *Stochastic Environmental Research and Risk Assessment*, 27(7): 1741-1752.
- Liu J, Schaaf C, Strahler A, et al. 2009. Validation of moderate resolution imaging spectroradiometer (MODIS) albedo retrieval algorithm: dependence of albedo on solar zenith angle. *Journal of Geophysical Research Atmospheres*, 114(D1): 1106-1117.
- Liu J G, Li S X, Ouyang Z Y, et al. 2008. Ecological and socioeconomic effects of China' s policies for ecosystem services. *Proceedings of the National Academy of Sciences of the United States of America*, 105(28): 9477-9482.
- Liu Q, McVicar T R. 2012. Assessing climate change induced modification of Penman potential evaporation and runoff sensitivity in a large water-limited basin. *Journal of Hydrology*, 464-465: 352-362.
- Lü Y H, Fu B J, Feng X M, et al. 2012. A policy-driven large scale ecological restoration: quantifying ecosystem services changes in the loess plateau of China. *PLoS ONE*, 7: e31782.
- McVicar T R, Niel T G V, Li L T, et al. 2007. Spatially distributing monthly reference evapotranspiration and pan evaporation considering topographic influences. *Journal of Hydrology*, 338(3-4): 196-220.
- Mu Q Z, Zhao M S, Running S W. 2011. Improvements to a MODIS global terrestrial evapotranspiration algorithm. *Remote Sensing of Environment*, 115(8): 1781-1800.
- Peng H, Jia Y W, Tague C, et al. 2015. An Eco-Hydrological Model-Based assessment of the impacts of soil and water conservation management in the Jinghe River Basin, China. *Water*, 7(11): 6301-6320.
- Peng S S, Piao S L, Zeng Z, et al. 2014. Afforestation in China cools local land surface temperature. *Proceedings of the National Academy of Sciences of the United States of America*, 111(8): 2915-2920.
- Potchter O, Goldman D, Kadish D, et al. 2008. The oasis effect in an extremely hot and arid climate: The case of southern Israel. *Journal of Arid Environments*, 72(9): 1721-1733.
- Ren R, Ma J J, Cheng Q Y, et al. 2017. An investigation into the effects of temperature gradient on the soil Water-Salt Transfer with evaporation. *Water*,

9(7): 456.

Reynolds J F, Smith D M S, Lambin E F, et al. 2007. Global desertification: building a science for dryland development. *Science*, 316(5826): 847-851.

Shen M G, Piao S L, Jeong S J, et al. 2015. Evaporative cooling over the Tibetan Plateau induced by vegetation growth. *Proceedings of the National Academy of Sciences of the United States of America*, 112(30): 9299-9304.

Vancutsem C, Ceccato P, Dinku T, et al. 2010. Evaluation of MODIS land surface temperature data to estimate air temperature in different ecosystems over Africa. *Remote Sensing of Environment*, 114(2): 449-465.

Wan Z. 2008. New refinements and validation of the MODIS Land-Surface Temperature/Emissivity products. *Remote Sensing of Environment*, 112(1): 59-74.

Wang Y P, Phipps S J, de Noblet N. 2011. Importance of background climate in determining impact of land-cover change on regional climate. *Nature Climate Change*, 1: 472-475.

Xiao K C, He T G, Chen H, et al. 2017. Impacts of vegetation restoration strategies on soil organic carbon and nitrogen dynamics in a karst area, southwest China. *Ecological Engineering*, 101: 247-254.

Xin Z B, Xu J X, Zheng W. 2008. Spatiotemporal variations of vegetation cover on the Chinese Loess Plateau (1981-2006): Impacts of climate changes and human activities. *Science in China Series D: Earth Science*, 51(1): 67-78.

Xu L P, Yang G H, Jiang Y, et al. 2008. Comparison of microclimate effects under three Forest-Grass rehabilitation of Artificial Vegetation in Loess Plateau. *Journal of Soil and Water Conservation*, 22: 163-173. (in Chinese)

Xue Y N, Liang H, Zhang Y, et al. 2017. Spatial and Temporal Variations of Land Surface Temperature of the Loess Plateau. *Earth and Environment*, 45(5): 500-507. (in Chinese)

Yang Y T, Long D, Guan H D, et al. 2015. GRACE satellite observed hydrological controls on interannual and seasonal variability in surface greenness over mainland Australia. *Journal of Geophysical Research Biogeosciences*, 119: 2245-2260.

Yang Y T, Guan H D, O Batelaan, et al. 2016. Contrasting responses of water use efficiency to drought across global terrestrial ecosystems. *Scientific Reports*, 6: 23284.

Yi H, Cai J L, He Y, et al. 2009. Correlation of precipitation to temperature variation in the Huanghe River (Yellow River) basin during 1957-2006. *Journal of Hydrology*, 372(1-4): 1-8.

Zhai J, Liu R, Liu J Y, et al. 2015. Human-Induced landcover changes drive a diminution of land surface albedo in the loess Plateau (China). *Remote Sensing*,

7(3): 2926-2941.

Zhang B, He C, Burnham M, et al. 2016. Evaluating the coupling effects of climate aridity and vegetation restoration on soil erosion over the Loess Plateau in China. *Science of the Total Environment*, 539: 436-449.

Zhang C, Xie G, Li S, et al. 2010. The productive potentials of sweet sorghum ethanol in China. *Applied Energy*, 87(7): 2360-2368.

Zhang D J, Tang R L, Zhao W, et al. 2014. Surface soil water content estimation from thermal remote sensing based on the temporal variation of land surface temperature. *Remote Sensing*, 6(4): 3170-3187.

Zhang X P, Zhang L, Zhao J, et al. 2008. Responses of streamflow to changes in climate and land use/cover in the Loess Plateau, China. *Water Resources Research*, 44(7): 2183-2188.

Zhou H J, van Rompaey A, Wang J. 2009. Detecting the impact of the “Grain for Green” program on the mean annual vegetation cover in the Shaanxi province, China using SPOT-VGT NDVI data. *Land Use Policy*, 26(4): 954-960.

Zhou L, Tian Y, Roy S B, et al. 2012. Impacts of wind farms on land surface temperature. *Nature Climate Change*, 2: 539-543.

Zhou L M, Dai A, Dai Y J, et al. 2009. Spatial dependence of diurnal temperature range trends on precipitation from 1950 to 2004. *Climate Dynamics*, 32(2-3): 429-440.

Zhou L M, Dickinson R E, Tian Y H, et al. 2007. Impact of vegetation removal and soil aridation on diurnal temperature range in a semiarid region: application to the sahel. *Proceedings of the National Academy of Sciences of the United States of America*, 104(46): 17937-17942.

Zhou Z C, Shangguan Z P, Zhao D. 2006. Modeling vegetation coverage and soil erosion in the Loess Plateau Area of China. *Ecological Modelling*, 198(1-2): 263-268.

*Note: Figure translations are in progress. See original paper for figures.*

*Source: ChinaXiv – Machine translation. Verify with original.*