

## Sub-parallel ridge-trench interaction and an alternative model for the Silurian-Devonian archipelago in Western Junggar and North-Central Tianshan in NW China

**Authors:** Zhang, Ji'en, Chen, Yichao, Xiao, Wenjiao, Wakabayashi, John, Windley, Brian F., Yin, Jiyuan, Zhang, Ji'en

**Date:** 2021-06-22T00:00:00+00:00

### Abstract

Plate boundary evolution and interpreted paleogeography commonly involve the interaction between oceanic ridges and trenches. Western Junggar and Central-North Tianshan, NW China, have previously been regarded as independent orogens, but this model was challenged by the discovery of their similar tropic-subtropic coral assemblages and of Precambrian and Early Paleozoic clasts transmitted from the Central Tianshan in Early to Middle Paleozoic sediments in SW Junggar, suggesting they had been in contact by then. The presence of Late Silurian-Early Devonian adakitic rocks in northern SW Junggar was explained by slab roll-back, which required an active subducting slab in SW Junggar. However, this model is inconsistent with the cessation of arc magmatism immediately after the adakitic magmatism indicating there was no subduction in SW Junggar at those times. By re-evaluating the relevance of four diagnostic features of modern Californian ridge subduction, we propose a new model to explain the Late Silurian-Early Devonian orogenic framework of NW China in terms of sub-parallel ridge-trench interaction. The transmitted clasts from Central Tianshan in Early Paleozoic sediments in SW Junggar, at the same time as the Late Ordovician and Middle Silurian coral assemblages, and the complementary accretionary complex in SW Junggar and magmatic arc in Central Tianshan, are all contributory evidence for an Early Paleozoic subduction system. This subduction zone was intruded by two 446-380 Ma double magmatic belts: 1. A ridge subduction-generated adakitic gabbro-basalt-andesite-diorite-granite-rhyolite suite that intruded a 504-446 Ma accretionary complex in SW Junggar, and 2. a magmatic arc in Central Tianshan. A buoyant subducting ridge rifted and separated these magmatic belts resulting in the opening of a new 414-325 Ma ocean, which is preserved in OIB- and MORB-type ophiolites in SW Junggar

and North Tianshan. These geological and geochemical relationships resemble those generated by oceanic spreading in the modern Gulf of California, which led to separation of magmatic belts in Baja California and mainland Mexico. The new ocean split the Central Tianshan magmatic arc from the Early Paleozoic Mt. Xiemisitai-Barleik-Mayile accretionary complex in SW Junggar, which changed to a new passive continental margin that led to deposition of Devonian shallow marine-terrestrial sediments, and to cessation of magmatism at 379-349 Ma. These features, including the Middle Paleozoic orogenic architecture and the geochemical properties of specific magmatic rocks, indicate the development of a Late Silurian-Early Devonian sub-parallel-to-trench ridge subduction. This model not only interprets phenomena mentioned in the above earlier models, but also reconciles unresolved spatial relationships between Western Junggar and the Tianshan. Subsequent closure of the new ocean and subduction of its ridge in the Late Paleozoic gave rise to the archipelagic framework that we see today in NW China. An analysis of worldwide examples of sub-parallel ridge-trench interaction demonstrates that a ridge can undergo multiple episodes of subduction before a terminal ocean closure. In this context our new tectonic model for the western Junggar and Tianshan regions may provide an improved understanding of the structure and evolution of ridge-subduction events in ancient orogenic belts.

## Full Text

### Preamble

Silurian-Devonian sub-parallel ridge-trench interaction in Western Junggar and North-Central Tianshan in NW China: Alternative genesis of archipelagic architecture

Ji'en Zhang<sup>1</sup>, Yichao Chen<sup>2</sup>, Wenjiao Xiao<sup>1</sup>, John Wakabayashi<sup>3</sup>, Brian F. Windley<sup>1</sup>, Shuaihua Song<sup>1,2</sup>, and Jiyuan Yin

**1 State Key Laboratory of Lithospheric Evolution, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing 100029 China**

**3 Department of Earth and Environmental Sciences, California State University, Fresno, CA 93740 USA**

### ABSTRACT

Western Junggar and North-Central Tianshan in NW China comprise a double magmatic belt that evolved through 446-380 Ma SSZ-type gabbro-basalt-andesite-diorite-granite-rhyolite magmatism, which intruded a 504-446 Ma accretionary complex in SW Junggar and a coeval magmatic arc in Central Tianshan. This orogenic framework is interpreted as a product of sub-parallel ridge-trench interaction that generated the double magmatic belt together with

adakitic intrusions in the older accretionary complex. In this model, a buoyant subducted ridge stalled and separated the double magmatic belts, resulting in the opening of a new 414–325 Ma intra-arc ocean represented by Nb-depleted OIB- and MORB-type ophiolites. Mafic rocks generated by seafloor spreading in the modern Gulf of California record a similar evolution and chemistry. This new ocean split the northern accretionary complex along the Mt. Xiemisitai-Barleik-Mayile line, leading to deposition of Devonian shallow marine-terrestrial sediments and cessation of magmatism at 380–349 Ma; this evolution also resembles that of the late Cenozoic passive margins of Baja California. Subsequent removal of the new ocean and its ridge-subduction gave rise to an archipelagic framework in the Late Paleozoic. A worldwide analysis of published examples of sub-parallel ridge-trench interaction indicates that a ridge jump can lead to multiple episodes of subduction, which could occur long before terminal ocean closure.

## INTRODUCTION

A spreading ridge can interact with a trench at a range of convergence angles (Sisson et al., 2003). Sub-parallel subduction can generate a short-lived magmatic belt in a forearc setting, adjacent to a magmatic arc, comparable to relations in Eocene Alaska, modern western California and the Chile triple junction (Sisson et al., 2003); we term these synchronous sub-parallel magmatic belts as “double magmatic belts.”

Stalling of the subduction of a buoyant ridge (van Wijk et al., 2001) rifts the upper plate to develop a passive continental margin, and then forms new oceanic crust above upwelling sub-ridge asthenosphere, separating the two magmatic belts, as has taken place in Baja California and the Gulf of California (Michaud et al., 2006). Such a plate boundary configuration can evolve further into an archipelagic architecture with continued oceanic spreading and subduction.

The western Altaids (Fig. 1a [Figure 1: see original paper]), one of the world’s classic orogens, was generated by subduction of the western segment of the Paleo-Asian Ocean (Xiao et al., 2015). This orogen evolved along a major subduction zone with local archipelagic configurations, which generally formed via forearc and backarc spreading with one magmatic belt (Windley et al., 2007; Xiao et al., 2015). However, existing models cannot explain the 446–380 Ma double magmatic belt, between which developed 414–325 Ma oceanic crust in SW Junggar and Chinese North-Central Tianshan, an eastern extension of the Yili block (Fig. 1b). This complex configuration requires an alternative tectono-magmatic model.

This paper will present a new tectonic model for the evolution of SW Junggar and North-Central Tianshan that includes mid-Paleozoic sub-parallel ridge-subduction leading to a Late Paleozoic archipelago configuration in the western Altaids.

## TECTONIC UNITS

Western Junggar is divisible into SW Junggar and NW Junggar units along a boundary north of the Tacheng basin-Mt. Xiemisitai-Hongguleleng (Fig. 1b). In the northern and western parts of SW Junggar, 472–531 Ma OIB- and MORB-type ophiolites are present, while 379–414 Ma Nb-depleted OIB- and MORB-type ophiolites crop out in the Baijiantan area and along the Darbut Fault (Figs. 1b, 2a) (Zhang et al., 2011; Zhang et al., 2018).

In the Mt. Xiemisitai-Mayile area of SW Junggar, 380–446 Ma SSZ-type gabbro-basalt-andesite-diorite-granite-rhyolite intrudes an Early Paleozoic accretionary complex (Chen et al., 2010; Yin et al., 2017). Magmatism of 300–349 Ma age is widespread to the north and south of the Late Paleozoic accretionary complex (Figs. 1b, 2a). NW Junggar includes Late Paleozoic Saur arc rocks and an accretionary complex with Carboniferous turbidites juxtaposed with Ordovician Kujibai ophiolite (Figs. 1b, 2d) (Chen et al., 2017).

The Central Tianshan exposes Precambrian basement, a long-lived 300–476.5 Ma magmatic belt, and Silurian-Carboniferous mylonite, gneiss, and amphibolites (Figs. 1b, 2e) (Wang et al., 2012). In the North Tianshan, situated between the Central Tianshan and SW Junggar, an accretionary complex consists of Carboniferous turbidites and 325–386 Ma Nb-depleted OIB- and MORB-type ophiolites (Figs. 1b, 2e) (Xu et al., 2006).

## METHODOLOGY

Constraining the orogenic processes of a short-lived archipelagic architecture with a dismembered accretionary complex and paired igneous arc is generally difficult because many evolutionary pathways can produce this framework. To address this issue, we first remove complicated younger orogenic units and affiliate them with an older episode that has only one accretionary complex and a contemporaneous arc constituting a subduction zone; we then link this subduction zone to the evolving spatial relationships of subsequent associated subduction episodes. It is noteworthy that deduced results of each orogenic stage should match the facts and data collected from field work and indoor analysis.

## EARLY PALEOZOIC SUBDUCTION SYSTEM

Li et al. (2006) proposed that the North-Central Tianshan and SW Junggar in China (separated by the Aibihu Fault) evolved independently (Fig. 1b). However, paleomagnetic data show that SW Junggar underwent 35° counterclockwise post-Permian rotation (Fig. 1b) (Choulet et al., 2011), suggesting that it was previously aligned parallel with the North-Central Tianshan.

In SW Junggar, a 479–446 Ma accretionary complex containing 458–504 Ma blueschist and Ordovician-Silurian turbidites lacks a coeval island arc. In the Central Tianshan, a 476.5–439 Ma continental arc lacks a paired accretionary complex (Fig. 1b).

Moreover, SW Junggar and Central Tianshan were in contact by the Early-Middle Paleozoic, based on the following evidence: (1) Quartzitic schist from the Laba River at the southern end of SW Junggar (Fig. 1b and DR2) and at Wenquan in the Central Tianshan (across the Aibihu Fault) share similar clastic components (Fig. 1b). (2) Silurian-Devonian strata from SW Junggar have detrital zircon age peaks at 380-500, 600, 900-1000, 1500-1700, and 2400-2500 Ma, similar to those in the Central Tianshan (Figs. 1b, 2b, f, DR3a-b). (3) These strata contain Precambrian and 446-479 Ma clasts whose sources are absent in SW Junggar (Figs. 2a, b, e), indicating provenance from the Central Tianshan.

Accordingly, the evidence suggests that the SW Junggar accretionary complex and Central Tianshan arc constitute a 476.5-446 Ma trench-arc system (Fig. 3a [Figure 3: see original paper]).

## MIDDLE PALEOZOIC SUB-PARALLEL RIDGE-TRENCH INTERACTION

In the mid-Paleozoic, continuously evolving subduction generated 380-446 Ma magmatic rocks that intruded the Early Paleozoic accretionary complex in SW Junggar (Chen et al., 2010) and a coeval magmatic arc in the Central Tianshan, forming a double magmatic belt (Fig. 2 [Figure 2: see original paper]). We interpret the magmatic rocks that intrude the accretionary complex—including adakitic granodiorites with positive Nd(t) in the Mt. Xiemisitai area (Fig. 3b) (Yin et al., 2017)—as products of a slab window beneath SW Junggar caused by either (a) a subducted spreading ridge or (b) a slab tear in old subduction lithosphere (Calmus et al., 2011). Between the double magmatic belts, younger 325-414 Ma Nb-depleted MORB-type ophiolites occur in SW Junggar and Chinese North Tianshan, which may have developed in (c) a rift basin above a buoyant subducting slab or (d) a back-arc/forearc basin.

If the MORB-type ophiolites developed in a sizable back-arc or forearc basin (model d), a coeval 349-380 Ma trench and island arc should be located in SW Junggar, in contrast to the actual field relationship (Fig. 1b).

The presence of 325-414 Ma ophiolites in the Late Paleozoic SW Junggar and Chinese North Tianshan accretionary complexes makes the slab-tear model (b) unlikely, because they are younger than the time of the double magmatic belts (446-414 Ma).

Therefore, we prefer hypothesis (a, c) that a Middle Paleozoic ridge parallel to a trench subducted beneath SW Junggar (Fig. 3c), as this best agrees with field relationships and has modern analogues. In our model, a buoyant subducted ridge stalled, resulting in rifting of the upper plate and opening of a new SW Junggar/Chinese North Tianshan ocean, similar to the extant Gulf of California (Michaud et al., 2006).

The trench migrated to the southern margin of this ocean in the Tianshan (Fig.

3c), enabling 414–386 Ma oceanic crust in SW Junggar to be accreted before the 386–325 Ma ophiolites in the Chinese North Tianshan (Figs. 1b, 2a, e). This resulted in generation of the Central Tianshan continental arc by a subduction system analogous to those in modern Alaska and Chile (Thorkelson et al., 2011) and produced 357–375 Ma deformation and metamorphism (Fig. 3c).

Subduction of the spreading ridge may have resulted in magmatic interaction with an overlying metasomatized mantle wedge, or it may have promoted partial melting of this mantle wedge to generate Nb-depleted MORB-type oceanic crust (Zhang et al., 2011); this would be similar to mafic rocks in the Gulf of California (Calmus et al., 2011).

Devonian shallow marine-terrestrial sediments with corals, brachiopods, sea lilies, and plants (Gong and Zong, 2015) were deposited on the northern margin of the Early Paleozoic accretionary complexes along the Mt. Xiemisitai-Barleik-Mayile line (Fig. 1b), and magmatism ceased from 380 to 349 Ma (Liu et al., 2017). These sedimentary-kinematic relations are similar to those in the young passive continental margin above a remnant slab in Baja California (Paulssen and de Vos, 2017). Moreover, Carboniferous sandstones in SW Junggar contain 349–380 Ma detrital zircons suggesting a source in the Central Tianshan (Fig. 2a, c, e, 3c).

## ARCHIPELAGIC ARCHITECTURE OF THE WESTERN ALTAIDS

The new ocean evolved to form Late Paleozoic intra-oceanic subduction systems with a continental arc in an overall archipelago configuration. The ocean subducted southward to generate the Chinese North Tianshan accretionary complex (Xu et al., 2006) and the long-lived Central Tianshan continental arc (Fig. 3d).

A double-vergent subduction system developed in SW Junggar (Zhang et al., 2011). Southward subduction generated an intra-oceanic arc at Guai 10 (Fig. 1b) in the NW Junggar Basin and an accretionary complex containing Carboniferous turbidites and the Baijiantan-Baikouquan ophiolites (Figs. 1b, 3d). Northward subduction resulted in formation of the Mt. Xiemisitai-Mayile-Barleik magmatic arc and an accretionary wedge including the Darbut ophiolite (Figs. 1b, 3d). Subduction of the young ridge (Fig. 3d) generated the Baogutu adakites, high-Mg# sanukitoids, and the Miaoergou charnockite (Zhang et al., 2011).

In NW Junggar, the Paleo-Asian Ocean subducted northward to form a Late Paleozoic accretionary complex that includes Carboniferous trench sandstones juxtaposed with the Ordovician Kujibai ophiolite (Fig. 3d) (Chen et al., 2017).

Our new model suggests that such an archipelagic architecture in the western Altaids evolved as a consequence of sub-parallel ridge subduction. This type of subduction mechanism and subsequent evolution leads to complex rifting and

amalgamation. The recognition of double magmatic belts provides a means of tracking such complex histories in orogenic belts.

## IMPLICATIONS

Sub-parallel ridge-trench interactions characterize modern subduction zones, so such interactions would be expected in the formation of some ancient orogenic belts such as the western Altaiids. It is noteworthy that a spreading ridge can undergo multiple episodes of subduction, resulting in generation of an archipelagic framework.

## ACKNOWLEDGEMENTS

We thank the referees for helpful reviews. This work has been supported by the National Key R & D Program of China (2017YFC0601206), the Strategic Priority Research Program (B) of CAS (XDB18030103, XDB18020203), and the National Natural Science Foundation of China (41472208).

## REFERENCES CITED

- Calmus, T., Pallares, C., Maury, R. C., Aguillón-Robles, A., Bellon, H., Benoit, M., and Michaud, F., 2011, Volcanic Markers of the Post-Subduction Evolution of Baja California and Sonora, Mexico: Slab Tearing Versus Lithospheric Rupture of the Gulf of California: *Pure and Applied Geophysics*, v. 168, no. 8, p. 1303-1330.
- Chen, J. F., Han, B. F., Ji, J. Q., Zhang, L., Xu, Z., He, G. Q., and Wang, T., 2010, Zircon U-Pb ages and tectonic implications of Paleozoic plutons in northern West Junggar, North Xinjiang, China: *Lithos*, v. 115, no. 1-4, p. 137-152.
- Chen, Y. C., Xiao, W. J., Windley, B. F., Zhang, J. E., Zhou, K. F., and Sang, M., 2017, Structures and detrital zircon ages of the Devonian-Permian Tarbagatay accretionary complex in west Junggar, China: imbricated ocean plate stratigraphy and implications for amalgamation of the CAO: *International Geology Review*, v. 59, no. 9, p. 1097-1115.
- Choulet, F., Chen, Y., Wang, B., Faure, M., Cluzel, D., Charvet, J., Lin, W., and Xu, B., 2011, Late Paleozoic paleogeographic reconstruction of Western Central Asia based upon paleomagnetic data and its geodynamic implications: *Journal of Asian Earth Sciences*, v. 42, no. 5, p. 867-884.
- Gong, Y. M., and Zong, R. W., 2015, Paleozoic stratigraphic regionalization and paleogeographic evolution in Western Junggar, Northwestern China: *Earth Science*, v. 40, no. 3, p. 461-484 (in Chinese with English abstract).
- Li, J. Y., He, G. Q., Xu, X., Li, H. Q., Sun, G. H., Yang, T. N., Gao, L. M., and Zhu, Z. X., 2006, Crustal tectonic framework of Northern Xinjiang and adjacent

regions and its formation: *Acta Geologica Sinica*, v. 80, no. 1, p. 148-168 (in Chinese with English abstract).

Liu, B., Han, B. F., Chen, J. F., Ren, R., Zheng, B., Wang, Z. Z., and Feng, L. X., 2017, Closure Time of the Junggar-Balkhash Ocean: Constraints From Late Paleozoic Volcano-Sedimentary Sequences in the Barleik Mountains, West Junggar, NW China: *Tectonics*, v. 36, no. 12, p. 2823-2845.

Michaud, F., Royer, J. Y., Bourgois, J., Dymant, J., Calmus, T., Bandy, W., Sossion, M., Mortera-Gutiérrez, C., Sichler, B., Rebolledo-Viera, M., and Pontoise, B., 2006, Oceanic-ridge subduction vs. slab break off: Plate tectonic evolution along the Baja California Sur continental margin since 15 Ma: *Geology*, v. 34, no. 1, p. 13-16.

Paulssen, H., and de Vos, D., 2017, Slab remnants beneath the Baja California peninsula: Seismic constraints and tectonic implications: *Tectonophysics*, v. 719-720, p. 27-

Sisson, V. B., Ravlis, T. L., Roeske, S. M., and Thorkelson, D. J., 2003, Introduction: An overview of ridge-trench interactions in modern and ancient settings, in Sisson, V. B., Roeske, S. M., and Pavlis, T. L., eds., *Geology of a transpressional orogen developed during ridge-trench interaction along the North Pacific margin*: Boulder, Colorado, GSA Special Paper, 371, p. 1-18.

Thorkelson, D. J., Madsen, J. K., and Sluggett, C. L., 2011, Mantle flow through the Northern Cordilleran slab window revealed by volcanic geochemistry: *Geology*, v. 39, no. 3, p. 267-270.

van Wijk, J. W., Govers, R., and Furlong, K. P., 2001, Three-dimensional thermal modeling of the California upper mantle: a slab window vs. stalled slab: *Earth and Planetary Science Letters*, v. 186, no. 2, p. 175-186.

Wang, B., Jahn, B. M., Shu, L. S., Li, K. S., Chung, S. L., and Liu, D. Y., 2012, Middle-Late Ordovician arc-type plutonism in the NW Chinese Tianshan: Implication for the accretion of the Kazakhstan continent in Central Asia: *Journal of Asian Earth Sciences*, v. 49, no. 0, p. 40-53.

Windley, B. F., Alexeiev, D., Xiao, W. J., Kröner, A., and Badarch, G., 2007, Tectonic models for accretion of the Central Asian Orogenic belt: *Journal of the Geological Society, London*, v. 164, p. 31-47.

Xiao, W., Windley, B. F., Sun, S., Li, J., Huang, B., Han, C., Yuan, C., Sun, M., and Chen, H., 2015, A Tale of Amalgamation of Three Permo-Triassic Collage Systems in Central Asia: Oroclines, Sutures, and Terminal Accretion: *Annual Review of Earth and Planetary Sciences*, v. 43, no. 1, p. 477-507.

Xu, X. Y., Li, X. M., Ma, Z. P., Xia, L. Q., Xia, Z. C., and Peng, S. X., 2006, LA-ICPMS zircon U-Pb dating of gabbro from the Bayingou ophiolite in the Northern Tianshan Mountains: *Acta Geologica Sinica*, v. 80, no. 8, p. 1168-1176 (in Chinese with English abstract).

Yin, J. Y., Chen, W., Xiao, W. J., Yuan, C., Windley, B. F., Yu, S., and Cai, K. D., 2017, Late Silurian-early Devonian adakitic granodiorite, A-type and I-type granites in NW Junggar, NW China: Partial melting of mafic lower crust and implications for slab roll-back: *Gondwana Research*, v. 43, p. 55-73.

Zhang, J. E., Xiao, W. J., Han, C. M., Ao, S. J., Yuan, C., Sun, M., Geng, H. Y., Zhao, G. C., Guo, Q. Q., and Ma, C., 2011, Kinematics and age constraints of deformation in a Late Carboniferous accretionary complex in Western Junggar, NW China: *Gondwana Research*, v. 19, no. 4, p. 958-974.

Zhang, J. E., Xiao, W. J., Luo, J., Chen, Y. C., Windley, B. F., Song, D. F., Han, C. M., and Safonova, I., 2018, Collision of the Tacheng block with the Mayile-Barleik-Tangbale accretionary complex in Western Junggar, NW China: Implication for Early-Middle Paleozoic architecture of the western Altaids: *Journal of Asian Earth Sciences*, v. 159, p. 259-278.

## FIGURE CAPTIONS

**Figure 1** (a) Geological map of the West Junggar and Chinese North-Central Tianshan, showing spatial relationships of the main orogenic components, including 380–446 Ma double magmatic belts, separated by 325–414 Ma oceanic crust, intruded into an Early-Middle Paleozoic accretionary complex in SW Junggar and a magmatic arc in the Central Tianshan. See Figure DR1 for further details. (b) Inserted schematic map of East Asia showing the location of the Altaids (Xiao et al., 2015).

**Figure 2** Time-stratigraphic relationships between ophiolites, magmatism, HP metamorphism, and detrital zircons in sedimentary rocks in SW Junggar (a–c), NW Junggar (d) and the Chinese North-Central Tianshan (e–f). Note the development of 380–446 Ma double magmatic belts, which are older than 325–414 Ma ophiolites in the SW Junggar and Chinese North Tianshan (a, e). Detrital zircons of Precambrian and 446–479, 349–380 Ma, which are magmatic gaps in the SW Junggar (a), are dated in Silurian-Devonian (b) and Carboniferous (c) strata. The source of these clasts is in the Central Tianshan (e–f). Light green bars and dark green graphs in b, c, f present detrital zircons of 300–550 and 550–3500 Ma, respectively.

**Figure 3** Four diagrams illustrating the tectonic crustal evolution in the West Junggar and Chinese North-Central Tianshan, showing the progressive development of Early Paleozoic subduction zone, Middle Paleozoic sub-parallel ridge-trench interaction, and a final Late Paleozoic archipelagic framework by 349–316 Ma. Acronyms are defined in Fig. 1. Further discussion in the text.

*Note: Figure translations are in progress. See original paper for figures.*

*Source: ChinaXiv – Machine translation. Verify with original.*