

Potential responses of vegetation to atmospheric aerosols in arid and semi-arid regions of Asia postprint

Authors: JIAO Linlin, WANG Xunming, CAI Diwen, HUA Ting, WANG Xunming

Date: 2021-06-04T00:00:00+00:00

Abstract

Changes in atmospheric aerosols have profound effects on ecosystem productivity, vegetation growth and activity by directly and indirectly influencing climate and environment conditions. However, few studies have focused on the effects of atmospheric aerosols on vegetation growth and activity in the vulnerable arid and semi-arid regions, which are also the source areas of aerosols. Using the datasets of aerosol optical depth (AOD), normalized difference vegetation index (NDVI) and multiple climatic variables including photosynthetically active radiation (PAR), surface solar radiation (SSR), surface air temperature (TEM) and total precipitation (PRE), we analyzed the potential responses of vegetation activity to atmospheric aerosols and their associated climatic factors in arid and semi-arid regions of Asia from 2005 to 2015. Our results suggested that areas with decreasing growing-season NDVI were mainly observed in regions with relatively sparse vegetation coverage, while AOD tended to increase as NDVI decreased in these regions. Upon further analysis, we found that aerosols might exert a negative influence on vegetation activity by reducing SSR, PAR and TEM, as well as suppressing PRE in most arid and semi-arid regions of Asia. Moreover, the responses of atmospheric aerosols on vegetation activity varied among different growing stages. At the early growing stage, higher concentration of aerosol was accompanied with suppressed vegetation growth by enhancing cooling effects and reducing SSR and PAR. At the middle growing stage, aerosols tended to alter microphysical properties of clouds with suppressed PRE, thereby restricting vegetation growth. At the late growing stage, aerosols exerted significantly positive influences on vegetation activity by increasing SSR, PAR and TEM in regions with high anthropogenic aerosols. Overall, at different growing stages, aerosols could influence vegetation activity by changing different climatic factors including SSR, PAR, TEM and PRE in arid and semi-arid regions of Asia. This study not only clarifies the impacts of aerosols on vegetation

activity in source areas, but also explains the roles of aerosols in climate.

Full Text

Preamble

Potential Responses of Vegetation to Atmospheric Aerosols in Arid and Semi-Arid Regions of Asia

JIAO Linlin^{1, 2, 3}, WANG Xunming^{2, 3*}, CAI Diwen^{2, 3}, HUA Ting⁴

¹ College of Mining Engineering, North China University of Science and Technology, Tangshan 063210, China

² Key Laboratory of Water Cycle and Related Land Surface Processes, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China

³ University of Chinese Academy of Sciences, Beijing 100049, China

⁴ Shaanxi Key Laboratory of Earth Surface System and Environmental Carrying Capacity, College of Urban and Environmental Sciences, Northwest University, Xi'an 710127, China

Abstract: Changes in atmospheric aerosols profoundly affect ecosystem productivity, vegetation growth, and activity by directly and indirectly influencing climate and environmental conditions. However, few studies have focused on the effects of atmospheric aerosols on vegetation growth and activity in vulnerable arid and semi-arid regions, which are also the source areas of aerosols. Using datasets of aerosol optical depth (AOD), normalized difference vegetation index (NDVI), and multiple climatic variables including photosynthetically active radiation (PAR), surface solar radiation (SSR), surface air temperature (TEM), and total precipitation (PRE), we analyzed the potential responses of vegetation activity to atmospheric aerosols and their associated climatic factors in arid and semi-arid regions of Asia from 2005 to 2015. Our results suggested that areas with decreasing growing-season NDVI were mainly observed in regions with relatively sparse vegetation coverage, while AOD tended to increase as NDVI decreased in these regions. Upon further analysis, we found that aerosols might exert a negative influence on vegetation activity by reducing SSR, PAR, and TEM, as well as suppressing PRE in most arid and semi-arid regions of Asia. Moreover, the responses of atmospheric aerosols on vegetation activity varied among different growing stages. At the early growing stage, higher concentration of aerosol was accompanied with suppressed vegetation growth by enhancing cooling effects and reducing SSR and PAR. At the middle growing stage, aerosols tended to alter microphysical properties of clouds with suppressed PRE, thereby restricting vegetation growth. At the late growing stage, aerosols exerted significantly positive influences on vegetation activity by increasing SSR, PAR, and TEM in regions with high anthropogenic aerosols. Overall, at different growing stages, aerosols could influence vegetation activity by changing different climatic factors including SSR, PAR, TEM, and PRE in

arid and semi-arid regions of Asia. This study not only clarifies the impacts of aerosols on vegetation activity in source areas, but also explains the roles of aerosols in climate.

Keywords: aerosol optical depth; climatic factors; normalized difference vegetation index; spatiotemporal distribution; indirect effect

*Corresponding author: WANG Xunming (E-mail: xunming@igsnrr.ac.cn)

Received 2019-09-17; revised 2021-01-15; accepted 2021-01-22

1 Introduction

Drylands, including arid and semi-arid regions, cover about 41% of the world's land surface area (Huang et al., 2017). Over the past centuries, drylands have dramatically transformed in response to frequent human activities and climate changes (IPCC, 2013). Vegetation growth is vulnerable to harsh environments in drylands (Peng et al., 2011; Pauli et al., 2012; Zhang et al., 2014; Wang et al., 2017), and climatic factors including solar radiation, temperature, and precipitation may have profound impacts on vegetation dynamics in these regions (Nemani et al., 2003; Park and Sohn, 2010; Mohammat et al., 2013; Kim et al., 2014).

Drylands are also regarded as potential sources of aerosols (Shao et al., 2011). Since the 2000s, atmospheric aerosols have increased rapidly, which would have remarkable influences on the surface energy budget of the Earth (Yoon et al., 2014; Hsu et al., 2017). On the one hand, aerosols could modify the shortwave radiation that the Earth receives by scattering and absorbing solar radiation (Chameides et al., 1999; Steiner et al., 2005; Wang et al., 2018). On the other hand, aerosols could interact with cloud droplets by serving as cloud condensation nuclei (CCN) or ice nuclei (IN), which would change the physical and radiative properties of clouds, and eventually alter the Earth's radiation balance (Chameides et al., 1999; Steiner et al., 2005; Wang et al., 2018).

Not only that, interactions between aerosol-induced radiation changes and microphysical processes of clouds could further produce complex effects on climatic factors (Gu et al., 2016; Ezhova et al., 2018). For instance, aerosols have been reported to cool the Earth surface with a particular reduction in night-time temperature (Nair et al., 2011). Moreover, since aerosols are involved in the microphysical processes of clouds (Rosenfeld et al., 2014; Dong et al., 2019), they would substantially affect regional precipitation, but their effects on precipitation differ among situations with various types of aerosols and clouds (Hui et al., 2008; Min et al., 2009; Yan et al., 2014). Among all types of aerosols, dust aerosols are found to change the number of cloud droplets and cloud water path, thereby affecting low cloud cover, high cloud cover, and water vapor contents, which ultimately alter the precipitation rate (Huang et al., 2006). In addition, dust aerosols may reduce the cloud particle sizes and even 'burn' the clouds (i.e.,

the heating effect; Yin and Chen, 2007), thus suppressing warm rain formation (Rosenfeld et al., 2001; Huang et al., 2009; Qian et al., 2009).

Radiative effects of aerosols could indirectly influence ecosystem productivity and CO₂ exchange through multiple mechanisms (Tie et al., 2016; Yue et al., 2017; Wang et al., 2018), which are found to vary among different ecosystems (Yue et al., 2017). Generally, the response of net ecosystem CO₂ exchange to aerosol loading is negative in grasslands (including semi-arid grasslands), but positive in forests and croplands, possibly due to their different canopy structures (Roderick et al., 2001; Niyogi et al., 2004; Xie et al., 2010).

Arid and semi-arid regions of Asia are considered as one of the major source areas of dust aerosols, which has emitted a total of >30% of global dust aerosols (Zhang et al., 1997; Ginoux et al., 2012). This region covers an area of 1.3×10^7 km² stretching from central and western to northeastern Asia, and receives an annual mean precipitation of <500 mm (Fig. 1 [Figure 1: see original paper]), which makes it extremely sensitive to climatic and environmental changes. For instance, Gessner et al. (2013) found that changes in the seasonal precipitation of arid and semi-arid regions in Asia, especially in Central Asia with an annual precipitation of 100–400 mm, were closely related to the evolution of vegetation. In addition, Zhang et al. (2016) found that increases in winter precipitation and summer temperature may be beneficial for vegetation greening in eastern areas of Asia. Mohammad et al. (2013) found that drought and spring cooling may reduce vegetation growth in extremely arid regions. However, potential effects of aerosols on vegetation activity in arid and semi-arid regions of Asia are still poorly understood.

In order to understand the direct effect of aerosols on climatic factors and their indirect effects on vegetation activity in arid and semi-arid regions of Asia, we used datasets of aerosol optical depth (AOD), normalized difference vegetation index (NDVI), and major climatic factors including photosynthetically active radiation (PAR) (surface visible solar radiation with a spectral range of 400–700 nm), surface solar radiation (SSR), surface air temperature (TEM), and total precipitation (PRE) to analyze the spatiotemporal variations in the correlations between vegetation activity and major climatic factors that are potentially influenced by aerosols. In this study, we aimed not only to identify the ecological effects of aerosols in their source areas, but also to clarify the roles of aerosols in climate and ecosystem.

Fig. 1 shows arid and semi-arid regions and land uses in Asia. Land uses are referenced from NASA (National Aeronautics and Space Administration; <https://lpdaac.usgs.gov>). AERONET, aerosol robotic network. 1, Dushanbe; 2, Issyk-Kul; 3, Semi-Arid Climate and Environment Observatory of Lanzhou University (SACOL); 4, Dalanzadgad.

2.1 Data Sources

NDVI data from 2005 to 2015 were obtained from MODIS (moderate-resolution imaging spectroradiometer)/Terra collection 5.0 Vegetation Index Products (MOD13C1-NDVI), which were downloaded from LAADS (level 1 and atmosphere archive and distribution system) web (<https://ladsweb.nascom.nasa.gov/>). This gridded dataset (with a temporal resolution of 16-d and a spatial resolution of $0.05^{\circ} \times 0.05^{\circ}$) was predominantly designed to provide a globally consistent assessment of vegetation condition (Running et al., 1994; Justice et al., 1998; Salano et al., 2010).

To improve the accuracy of the analysis results, we defined those pixels with a mean growing-season (81–304 d) NDVI of >0.1 as vegetated areas (Zhou et al., 2001). Datasets of MYD04 AOD have been widely used to develop global aerosol pattern as it comprehensively describes the optical properties of aerosols and quantitatively describes aerosol radiative forcing (Levy et al., 2013; Yu et al., 2013). In this study, we used the daily AOD data from 2005 to 2015, with a spatial resolution of $10 \text{ km} \times 10 \text{ km}$, from the MODIS Aqua collection 6.0 Aerosol Product (MYD04). The dataset 'Dark_{{{Target}}}{{{Deep}}}{{{{Blue}}}{{{{Optical}}}{{{{550}}}}_{{{Combined}}}' was used for further analysis, which could select one type of retrievals (if any) or an average of the two retrievals that applied the dark target or deep blue algorithm.

Ground-observation AOD was obtained from aerosol robotic network (AERONET) of NASA (National Aeronautics and Space Administration). CE318 sunphotometer was deployed in the AERONET for detecting the bands at wavelengths of 380, 440, 500, 675, 870, and 1020 nm. AERONET measurement was generally used to validate the satellite-retrieved AOD since it contained the spectral data on solar radiation, from which various aerosol optical properties could be retrieved using an inversion algorithm (Dubovik et al., 2000). The four AERONET measurement sites of AOD data in arid and semi-arid regions of Asia (including Dushanbe (38.55°N , 68.86°E), Issyk-Kul (42.62°N , 76.98°E), Semi-Arid Climate and Environment Observatory of Lanzhou University (SACOL) (35.95°N , 104.14°E), and Dalanzadgad (43.56°N , 104.42°E)) were used to validate the MYD04-AOD data from 2005 to 2015 (Fig. 1).

Since atmospheric aerosols have affected vegetation growth primarily by altering the quantities and qualities of PAR or by changing climatic factors (surface temperature and precipitation) (Mike, 2002), the relationship between NDVI and parameters such as TEM, PRE, SSR, and PAR were analyzed in this study. All climatic factor data were obtained from the ERA-Interim reanalysis from the European Center for Medium-Range Weather Forecasts (ECMWF; Simmons et al., 2006). All variables (TEM, PRE, SSR, and PAR) had a spatial resolution of $0.125^{\circ} \times 0.125^{\circ}$ and a temporal resolution of 12 h (except for TEM with a 3-h resolution).

2.2.1 AOD at 550 nm Wavelength

To compare satellite-retrieved AOD with that from ground observation, it is necessary to obtain AOD at the same wavelength by interpolating AOD from ground observation. In this study, AOD and Angström exponents of 440 and 675 nm channels in the level 2.0 products were interpolated to a wavelength of 550 nm after cloud screening and quality control with the AERONET. The interpolation method can be expressed as:

$$\tau(\lambda) = \beta \lambda^{-\alpha_{ab}}$$

where $\tau(\lambda)$ is the AOD of the λ channel; β is the Angström turbidity coefficient; and α_{ab} is the Angström exponent between wavelengths of a and b nm (here the wavelength range is 440–675 nm).

For the measurements at all sites collected from 2005 to 2015, we extracted the AERONET data at 13:30 (LST) with a 3-h time window (i.e., between 12:00 and 15:00), which just corresponded to the passing time of the MODIS data (Ginoux et al., 2012). AOD (550 nm) values from the AERONET network were then averaged during the satellite overpass time, which could be compared with the regional mean of MODIS retrievals within a 0.5° radius around each ground station (Mi et al., 2007).

2.2.2 Linear Trend in NDVI and AOD at Growing Stages

AOD and NDVI at growing stages were obtained by averaging the values from 81 to 304 d every year. Then, linear trends in NDVI and AOD at growing stages were calculated in each pixel using ordinary least square (OLS) regression method:

$$y = \alpha + \beta x + \varepsilon$$

where y is the annual mean NDVI or AOD; α is the intercept; β is the slope; x is the monitored year; and ε is the error term.

The slope was interpreted as a measure of linear trends in NDVI and AOD over time, and t-test was used to determine the significance of linear trends in NDVI and AOD for each pixel.

2.2.3 Partial Correlation Between NDVI and Climatic Factors

To examine the potential impact of atmospheric aerosols on vegetation activity, we estimated the partial correlation between NDVI and climatic factors. Partial correlation is used to measure the linear dependence between two variables when the influence of possible controlling variables is excluded (Engström and Ekman,

2010). In the case of k variables $(x_1, x_2, \dots, x_k)$, the partial correlation is defined as:

$$r_{ij|g} = \frac{r_{ij|g-1} - r_{ig|g-1}r_{jg|g-1}}{\sqrt{(1 - r_{ig|g-1}^2)(1 - r_{jg|g-1}^2)}}$$

where $r_{ij|g-1}$, $r_{ig|g-1}$, and $r_{jg|g-1}$ are the remaining correlations between x_i and x_j , between x_i and x_{lg} , and between x_j and x_{lg} , respectively, assuming no $g-1$ controlling variables; and $r_{ij|g}$ is the remaining correlation between x_i and x_j , assuming no g controlling variables.

In this study, all datasets (AOD, NDVI, and climatic factors) were preprocessed to have the same resolution (temporal resolution of 16-d and spatial resolution of $0.125^\circ \times 0.125^\circ$). Then, we calculated the partial correlation between NDVI and climatic factors in two different scenarios: (1) scenario a was used to calculate the partial correlation between NDVI and each climatic factor with the other three factors as controlling variables; and (2) scenario b was used to get the partial correlation between NDVI and each climatic factor with both the other climatic factors and AOD as controlling variables. Significance of partial correlation could be given using a two-tailed t-test. Finally, we compared the spatial patterns of inter-annual partial correlation between NDVI and four climatic factors for each 16-d period in different scenarios.

To further explore the influence of aerosols on the correlation between NDVI and climatic factors at different growing stages (including early (81–144 d), middle (145–240 d), and late growing stage (241–304 d)), we selected five typical regions with significant NDVI change trends at growing stages, including the Caspian Coastal Plain (region A1), the Indus River Plain (region A2), eastern Mongolian Plateau (region A3), the Ordos Plateau (region A4), and eastern Kazakhskiy Melkosopochnik (region A5) (Fig. 1). In addition, percentages of entire region and five typical regions showing significantly positive correlations were also calculated for each 16-d period to represent the evolutions of partial correlation at different growing stages.

3.1 Comparison of AERONET AOD and MODIS AOD

In arid and semi-arid regions of Asia, comparison of AOD values from AERONET observations and MODIS retrievals showed general agreements at four sites ($P < 0.05$; Fig. 2 [Figure 2: see original paper]), especially for the Dushanbe site, which had a correlation coefficient as high as 0.79. Compared with in situ observation, MODIS AOD values at SACOL, Dushanbe, and Dalanzadgad sites were underestimated, but it was slightly overestimated at Issyk-Kul site. Although there were some MODIS-retrieved AOD values that lay beyond the error range of AERONET observation, the MODIS AOD values could still represent the regional AOD characteristics in arid and semi-arid regions of Asia.

Fig. 2 shows a scatter plot of MODIS (moderate-resolution imaging spectroradiometer) retrieved AOD (aerosol optical depth) against AERONET (aerosol robotic network) AOD.

3.2 Temporal Variations of NDVI and AOD at Growing Stages

NDVI climatology at growing stages from 2005 to 2015 showed that across arid and semi-arid regions of Asia, higher NDVI (>0.4) at growing stages was primarily found in eastern Inner Mongolian Plateau, northern Mongolian Plateau, and eastern Kazakhskiy Melkosopochnik (Fig. 3a [Figure 3: see original paper]), which were mainly covered by forests, croplands, and grasslands (Fig. 1), whereas NDVI at growing stages ranging between 0.1 and 0.2 mainly scattered in the Caspian Sea Plain, Turan and Indus River Plains with relatively sparse vegetation. From 2005 to 2015, about 13.2% of arid and semi-arid regions of Asia showed significantly increasing NDVI trends at growing stages (Fig. 3b), which were mainly distributed in northeastern Mongolia and the Ordos Plateau, with a maximum increasing rate of $>0.02/a$. By comparison, only 4.5% of arid and semi-arid regions of Asia showed significantly decreasing NDVI trends at growing stages ($P<0.05$), with a decreasing rate of $>0.01/a$.

Fig. 3b shows areas passing the significance test ($P<0.05$) in a gray square grid.

Average AOD at growing stages from 2005 to 2015 showed that in arid and semi-arid regions of Asia, regions with higher AOD were mainly located in the Indus River Plain, Tarim Basin, and the Caspian Coastal Plain, whereas lower AOD was mainly found in northern Mongolian Plateau and eastern Kazakhskiy Melkosopochnik (Fig. 4a [Figure 4: see original paper]). In arid and semi-arid regions of Asia, regions with higher AOD basically corresponded to those areas with lower NDVI, except for few regions such as northern Mongolian and northeastern Inner Mongolian Plateau. From 2005 to 2015, about 50.0% of the entire study area showed increases in AOD (Fig. 4b), but only 6.8% of those regions were significant ($P<0.05$) and were mainly located in lower-elevation regions such as the Caspian Coastal Plain (with an increasing rate of $>0.05/a$). On the other hand, 7.0% of total areas experienced significantly decreasing AOD trends at growing stages ($P<0.05$), which were mainly located at higher-elevation regions, particularly in parts of eastern Mongolian Plateau, with a maximum decreasing rate of $>0.05/a$.

Fig. 4b shows areas passing the significance test ($P<0.05$) in a gray square grid.

3.3.1 Effect of PAR on Vegetation Activity

Partial correlation between PAR and NDVI in scenario b (i.e., taking AOD, PRE, TEM, and SSR as the controlling variables) showed that when excluding the influence of AOD in addition to the other three climatic factors, area percentages with significantly positive correlation were much larger than those in

scenario a (i.e., merely excluding the influence of PRE, TEM, and SSR). This phenomenon was more obvious at the early growing stage than at any other stages (Table 1).

Moreover, five typical regions were selected to show the detailed differences in partial correlation between scenarios a and b. For instance, at both early and late growing stages, area percentage with significantly positive correlation of all regions was larger in scenario b than in scenario a, indicating that when excluding the influence of AOD, partial correlation between PAR and NDVI had increased in arid and semi-arid regions of Asia. By comparison, at the middle growing stage, there was little change in area percentage between two scenarios.

The evolution of partial correlation between PAR and NDVI at different growing stages (Figs. 5 and S1) showed that when excluding the influence of AOD (scenario b), area percentage with significantly positive correlation was found in the Caspian Coastal Plain and Turan Plain at the early growing stage, in eastern Kazakhskiy Melkosopochnik and southeastern Mongolian Plateau at the middle growing stage, and southern Mongolian Plateau at the late growing stage. This phenomenon might be due to the obviously different responses of vegetation with different canopy structures on PAR at three growing stages.

3.3.2 Effect of PRE on Vegetation Activity

Partial correlation between PRE and NDVI in scenario b (i.e., excluding the influence of both AOD and other three climatic factors) showed that area percentage with significantly positive correlation was larger than that in scenario a (i.e., excluding the influence of other three climatic factors), especially at the middle growing stage (Table 2), implying the adverse impact of AOD on NDVI. Although five typical regions (A1–A5) showed similar changes, there were much more notable negative effects of atmospheric aerosols on vegetation growth over the Ordos Plateau (A4) at the early growing stage and over eastern Mongolian Plateau (A3) at the late growing stage, which is probably because atmospheric aerosols inhibited vegetation growth by suppressing precipitation via ‘burning’ the cloud droplets (Table 2; Figs. 6 and S2).

Among five typical regions, there were still spatial differences in the response of vegetation activity at growing stages on precipitation when the effect of atmospheric aerosols was eliminated (i.e., changes from scenario a to scenario b). For instance, at the early growing stage, there was an evident increase in area percentage with a significant correlation (PRE vs. NDVI) over the Ordos Plateau when influence of AOD was excluded, whereas there was little change between two scenarios over eastern Mongolian Plateau. This difference at the early growing stage was possibly because of their different ways of water supply to vegetation growth: plants over the Ordos Plateau were strongly constrained by precipitation, and atmospheric aerosols could influence local vegetation by altering precipitation; while area percentage on eastern Mongolian Plateau was

more affected by spring snow-melt water, thus making the role of precipitation in vegetation activity less important so that the influence of AOD was diminished. By comparison, from the middle to late growing stage, there was an evident increase in areas with significant correlations over eastern Mongolian Plateau rather than the Ordos Plateau from scenarios a to b. During these stages, precipitation became the dominant factor limiting vegetation growth over eastern Mongolian Plateau due to its continental arid climate, whereas the Ordos Plateau was under the control of summer monsoon with relatively ample precipitation, making local vegetation less vulnerable to precipitation variations so that the influence of AOD on NDVI was relatively less important.

3.3.3 Effect of SSR on Vegetation Activity

Partial correlation between SSR and NDVI showed that area percentage of the entire study area with significantly positive correlation was larger in scenario b than in scenario a, particularly at the early and middle growing stages (Table 3), indicating that the relationship between SSR and NDVI was closer without the influence of AOD. This was basically true for five typical regions, with the only exception of the Indus River Plain (A2). When AOD was used as the control variable (scenario b), NDVI was significantly and positively correlated with SSR over eastern Turan Plain and eastern Kazakhskiy Melkosopochnik at the early growing stage, and eastern and central Mongolian Plateau at the late growing stage (Figs. 7 and S3). However, area percentage with a significantly positive correlation was smaller over the Indus River Plain (A2) and the Ordos Plateau (A3) at the early growing stage.

3.3.4 Effect of TEM on Vegetation Activity

When the influence of aerosols was eliminated in partial correlation, area percentage with a significantly positive correlation between TEM and NDVI generally increased (i.e., from scenario a to b), which was more pronounced at the middle growing stage than at the other growing stages (Table 4). A similar change pattern was also found in five typical regions (Table 4). When AOD was used as the control variable that was eliminated in partial correlation, there were significantly positive correlations over the Turan Plain, eastern Kazakhskiy Melkosopochnik, and the Caspian Coastal Plain at the early growing stage; and over northern Kazakhskiy Melkosopochnik and the Indus River Plain at the late growing stage (Figs. 8 and S4).

4 Discussion

Atmospheric aerosols affect terrestrial vegetation activity indirectly via changing climate systems and ecosystems, including enhancing the scattering and absorption of solar radiation, changing atmospheric temperature, altering the probability and quantity of precipitation, and providing nutrients to some extent (Ramanathan et al., 2001; Steiner et al., 2005; Wild, 2009). In addition, atmo-

spheric aerosols may influence vegetation production and activity through the modulation of their photosynthesis, respiration, metabolism, and water utilization (Tie et al., 2016). In the present study, our results showed that atmospheric aerosols had clear effects on vegetation activity by influencing SSR, PAR, TEM, and PRE in arid and semi-arid regions of Asia.

4.1 Relationship Between Major Climatic Factors and Vegetation Activity When Excluding the Influence of Aerosols

Over the arid and semi-arid regions of Asia, a large proportion of areas exhibited significantly positive correlations between major climatic factors and NDVI when the influence of AOD was excluded. Firstly, area percentage with a significantly positive correlation between vegetation activity and PAR became higher when the influence of AOD was excluded, which could be attributable to the decreases in total PAR that could have facilitated vegetation growth. There were two plausible mechanisms behind the decreased PAR: (1) one explanation was that in the majority of study area, the regions with sparsely covered vegetation had emitted relatively higher amounts of dust aerosols and resultant decreasing PAR, which inhibited vegetation growth; and (2) the other explanation was that total PAR was still decreasing in areas with dense vegetation (such as north-eastern Inner Mongolia) due to the high concentration of aerosols from their source areas and consequently the decreasing PAR reduced the areas in which PAR facilitates vegetation growth (Xie et al., 2010).

Secondly, area percentage with a significantly positive correlation between SSR and NDVI became smaller when the influence of AOD was considered, which was possibly because of the negative forcing of higher aerosol emissions on SSR in arid and semi-arid regions of Asia with sparse vegetation coverage (Lau et al., 2006; Zhang et al., 2009; Huang et al., 2014). Generally, the surface cooling effects of aerosols (Du et al., 2014; Huang et al., 2014) would delay the onset of vegetation phenology in spring with reduced material accumulation of vegetation, and would trigger a higher risk of frost with restrained vegetation growth and decreased nutrient availability at the entire growing stage (Bokhorst et al., 2008; Bennie et al., 2010; He et al., 2015).

Thirdly, area percentage with a significantly positive correlation between PRE and NDVI was larger when eliminating the effects of AOD, possibly because aerosols may suppress precipitation by altering the cloud particle size and cloud temperature (Yin and Chen, 2007; Rosenfeld et al., 2014; Huang et al., 2016), thus restraining vegetation activity in arid and semi-arid regions (Schimel et al., 2001; Fang et al., 2004; Chuai et al., 2013). Moreover, enhanced aerosol emissions from potential source areas were generally accompanied by reduced TEM and suppressed precipitation (Piao et al., 2006; Gessner et al., 2013), which would be unfavorable for vegetation activity. It should be noted that the Inner Mongolian Plateau did not show significantly positive correlation between NDVI and major climatic factors when AOD was used as a control variable, which was possibly because vegetation of this region might be more attributable

to anthropogenic activities than to atmospheric aerosols. For instance, the afforestation program has been proven to promote vegetation restoration in recent years in China (He et al., 2015; Liu et al., 2015).

4.2 Spatiotemporal Variation in the Relationship Between Major Climatic Factors and Vegetation Activity

There were obvious spatial and temporal differences in responses of vegetation to climatic factors under the influence of aerosols. For instance, at the early growing stage, there were significantly positive correlations between NDVI and climatic factors including PAR, SSR, and TEM over the Caspian Coastal Plain and the Turan Plain when the influence of aerosols was excluded, which were all correlated with the higher concentration of spring aerosols that had substantially reduced PAR, SSR, and TEM via altering the incident (diffuse and direct) solar radiation (Cheng et al., 2015), leading to decreased contributions of these factors to vegetation growth (Mohammat et al., 2013; Huang et al., 2014; He et al., 2015; Bi et al., 2017).

At the middle growing stage, area percentages with significantly positive correlations of NDVI with PAR, SSR, and TEM were only found in central and western Kazakhskiy Melkosopchnik and southeastern Mongolian Plateau when the influence of aerosols was not considered, which could be attributed to the following reasons: firstly, photosynthesis of vegetation over these regions with a simple canopy structure would easily reach its saturation points in summer, even though dramatic reduction of aerosol concentrations in these regions could increase both PAR and SSR (Niyogi et al., 2004; Strada et al., 2015); secondly, the increased surface temperature resulting from descending aerosol concentrations was not any more the key climatic factor influencing vegetation growth in hot summers (Gessner et al., 2013; Mohammat et al., 2013). However, there was still a significantly positive correlation between PRE and NDVI with an extra exclusion of the influence of aerosols in most regions because dust aerosols (the main type of atmospheric aerosols) remained an important factor in suppressing PRE in summer: dust aerosols embedded in the cloud layer would reduce cloud cover (especially the lower cloud) and water vapor content, and would increase the evaporation of cloud droplets, thus eventually constraining vegetation growth (Huang et al., 2006; Huang et al., 2014; Li et al., 2018).

At the late growing stage, area percentages with significantly positive correlations between NDVI and PAR as well as SSR evidently increased over the Indus River Plain when excluding the influence of aerosols, possibly because the aerosols emitted from this region were primarily generated by anthropogenic activities, which would absorb SSR and PAR more easily than other types of aerosols (Dey et al., 2004). In addition, area percentage with a significantly positive correlation between TEM and NDVI also increased on the Indus River Plain probably due to the enhanced soil evapotranspiration driven by increased temperature, which was associated with an increased proportion of anthropogenic activities (Latha et al., 2018). By comparison, a significantly positive correla-

tion between PRE and NDVI was observed in most of the study area with the only exception of the Indus River Plain, possibly due to the suppressed precipitation that excessive inorganic (dust) aerosol nuclei could reduce the volume of droplets (Qian et al., 2009; Li et al., 2011; Rosenfeld et al., 2014).

5 Conclusions

From 2005 to 2015, there were decreasing trends in NDVI over the regions with sparse vegetation coverage, such as the Caspian Coastal Plain, which had a maximum decreasing rate of $>1 \times 10^{-2} \text{ } \$/\text{a}$. However, AOD tended to increase as NDVI decreased in most regions. When there was an extra exclusion of the influence of aerosols, the responses of vegetation to climatic factors showed clear spatiotemporal differences. Generally, at different growing stages, aerosols could influence vegetation activity by changing different climatic factors. At the early growing stage, the higher concentration of aerosols would reduce PAR, SSR, and TEM, thereby constraining vegetation activity, which was obvious over northern Kazakhskiy Melkosopchnik. At the middle growing stage, decreased concentration of aerosols had little effects on PAR, SSR, and TEM, but greatly suppressed precipitation and eventually regulated vegetation activity. At the late growing stage, the potential effects of aerosols on vegetation activity were evident in areas with high anthropogenic activities, such as in the Indus River Plain. Different responses of vegetation activity to aerosols and their related climatic factors would be due to differences in the vegetation canopy structures, their light utility efficiency, as well as regional environmental and climatic conditions. Our studies on the impact of aerosols on vegetation activity in arid and semi-arid regions of Asia would not only clarify the ecological effects of the aerosols in the source areas, but also explain the roles of aerosols in climate systems and ecosystems.

Acknowledgements: This work was supported by the National Key Research and Development Program of China (2016YFA0601900), the Key Frontier Program of the Chinese Academy of Sciences (QYZDJ-SSW-DQC043), the National Natural Science Foundation of China (41771012), and the Applied and Basic Research Program from Tangshan Science and Technology Bureau, China (20130202b).

References

- Bennie J J, Wiltshire A J, Joyce A N, et al. 2010. Characterizing inter-annual variation in the spatial pattern of thermal microclimate in a UK upland using a combined empirical-physical model. *Agricultural and Forest Meteorology*, 150(1): 1-15.
- Bi J R, Huang J P, Shi J S, et al. 2017. Measurement of scattering and absorption properties of dust aerosol in a Gobi farmland region of northwestern China—a potential anthropogenic influence. *Atmospheric Chemistry and Physics*, 17: 7775-7792.

- Bokhorst S, Bjerke J W, Bowles F W, et al. 2008. Impacts of extreme winter warming in the sub-Arctic: growing season responses of dwarf shrub heathland. *Global Change Biology*, 14(11): 2603–2612.
- Chameides W L, Yu H, Liu S C, et al. 1999. Case study of the effects of atmospheric aerosols and regional haze on agriculture: An opportunity to enhance crop yields in China through emission controls. *Proceedings of the National Academy of Sciences of the United States of America*, 96(24): 13626–13633.
- Cheng S J, Bohrer G, Steiner A L, et al. 2015. Variations in the influence of diffuse light on gross primary productivity in temperate ecosystems. *Agricultural and Forest Meteorology*, 201: 98–110.
- Chuai X W, Huang X J, Wang W J, et al. 2013. NDVI, temperature and precipitation changes and their relationships with different vegetation types during 1998–2007 in Inner Mongolia, China. *International Journal of Climatology*, 33(7): 1696–1706.
- Dey S, Tripathi S N, Singh R P. 2004. Influence of dust storms on the aerosol optical properties over the Indo-Gangetic Basin. *Journal of Geophysical Research*, 109: D20211.
- Dong B, Wilcox L J, Highwood E J, et al. 2019. Impacts of recent decadal changes in Asian aerosols on the East Asian summer monsoon: roles of aerosol-radiation and aerosol-cloud interactions. *Climate Dynamics*, 53(5): 3235–3256.
- Du J, He Z B, Yang J J, et al. 2014. Detecting the effects of climate change on canopy phenology in coniferous forests in semi-arid mountain regions of China. *International Journal of Remote Sensing*, 35(17): 6490–6507.
- Dubovik O, King M D. 2000. A flexible inversion algorithm for retrieval of aerosol optical properties from sun and sky radiance measurements. *Journal of Geophysical Research*, 105(D16): 20673–20696.
- Engström A, Ekman A M L. 2010. Impact of meteorological factors on the correlation between aerosol optical depth and cloud fraction. *Geophysical Research Letters*, 37(18): 1480–1493.
- Ezhova E, Ylivinkka I, Kuusk J, et al. 2018. Direct effect of aerosols on solar radiation and gross primary production in boreal and hemiboreal forests. *Atmospheric Chemistry and Physics*, 18: 17863–17881.
- Fang J Y, Piao S L, He J S, et al. 2004. Increasing terrestrial vegetation activity in China, 1982–1999. *Science China Series C*, 47(3): 229–240.
- Gessner U, Naeimi V, Klein I, et al. 2013. The relationship between precipitation anomalies and satellite-derived vegetation activity in Central Asia. *Global Planetary Change*, 110: 74–87.
- Ginoux P, Prospero J M, Gill T E, et al. 2012. Global-scale attribution of anthropogenic and natural dust sources and their emission rates based on MODIS deep blue aerosol products. *Review of Geophysics*, 50(3): RG3005.

- Gu L, Pallardy S G, Hosman K P, et al. 2016. Impacts of precipitation variability on plant species and community water stress in a temperate deciduous forest in the central US. *Agricultural and Forest Meteorology*, 217: 120–136.
- He B, Chen A F, Wang H L, et al. 2015. Dynamic response of satellite-derived vegetation growth to climate change in the Three North Shelter Forest Region in China. *Remote Sensing*, 7(8): 9998–10016.
- Hsu N C, Gautam R, Sayer A M, et al. 2017. Global and regional trends of aerosol optical depth over land and ocean using SeaWiFS measurements from 1997 to 2010. *Atmospheric Chemistry and Physics*, 17: 8037–8053.
- Huang J P, Minnis B, Lin T, et al. 2006. Possible influences of Asian dust aerosols on cloud properties and radiative forcing observed from MODIS and CERES. *Geophysical Research Letters*, 33(6): L06824.
- Huang J P, Fu Q, Su J, et al. 2009. Taklimakan dust aerosol radiative heating derived from CALIPSO observations using the Fu-Liou radiation model with CERES constraints. *Atmospheric Chemistry and Physics*, 9: 4011–4021.
- Huang J P, Wang T H, Wang W C, et al. 2014. Climate effects of dust aerosols over East Asian arid and semiarid regions. *Journal of Geophysical Research*, 119: 11398–11416.
- Huang J P, Li Y, Fu C, et al. 2017. Dryland climate change: Recent progress and challenges. *Review of Geophysics*, 55(3): 719–778.
- Hui W J, Cook B I, Ravi S, et al. 2008. Dust-rainfall feedbacks in the West African Sahel. *Water Resources Research*, 44(5): W05202.
- IPCC. 2013. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. New York: Cambridge University Press, 1–12.
- Justice C O, Vermote E, Townshend J R G, et al. 1998. The moderate resolution imaging spectroradiometer (MODIS): Land remote sensing for global change research. *IEEE Transactions on Geoscience and Remote Sensing*, 36(4): 1228–1249.
- Kim Y, Kimball J S, Didan K, et al. 2014. Response of vegetation growth and productivity to spring climate indicators in the conterminous United States derived from satellite remote sensing data fusion. *Agricultural and Forest Meteorology*, 194: 132–144.
- Latha R, Vinayak B, Murthy B S. 2018. Response of heterogeneous vegetation to aerosol radiative forcing over a northeast Indian station. *Journal of Environmental Management*, 206: 1224–1232.
- Lau K M, Kim M K, Kim K M. 2006. Asian summer monsoon anomalies induced by aerosol direct forcing: The role of the Tibetan Plateau. *Climate Dynamics*, 26: 855–864.

- Levy R C, Mattoo S, Munchak L A, et al. 2013. The collection 6 MODIS aerosol products over land and ocean. *Atmospheric Measurement Techniques*, 6: 2989–3034.
- Li X Q, Ting M F, Lee D E. 2018. Fast adjustments of the Asian summer monsoon to anthropogenic aerosols. *Geophysical Research Letters*, 45(2): 1001–1010.
- Li Z Q, Niu F, Fan J W, et al. 2011. Long-term impacts of aerosols on the vertical development of clouds and precipitation. *Natural Geoscience*, 4: 888–894.
- Liu X F, Zhu X F, Li S S, et al. 2015. Changes in growing season vegetation and their associated driving forces in China during 2001–2012. *Remote Sensing*, 7(11): 15517–15535.
- Mi W, Li Z Q, Xia X G, et al. 2007. Evaluation of the Moderate Resolution Imaging Spectroradiometer aerosol products at two aerosol robotic network stations in China. *Journal of Geophysical Research*, 112: D22S08.
- Mike B. 2002. *The Influence of Aerosols on Plant Growth*. Honolulu: Georgia Institute of Technology, 5–10.
- Min Q L, Li R, Lin B, et al. 2009. Evidence of mineral dust altering cloud microphysics and precipitation. *Atmospheric Chemistry and Physics*, 9(9): 3223–3231.
- Mohammat A, Wang X H, Xu X T, et al. 2013. Drought and spring cooling induced recent decrease in vegetation growth in Inner Asia. *Agricultural and Forest Meteorology*, 178–179: 21–30.
- Nair U S, Mcnider R T, Patadia F, et al. 2011. Sensitivity of nocturnal boundary layer temperature to tropospheric aerosol surface radiative forcing under clear-sky conditions. *Journal of Geophysical Research*, 116(D2): D02205.
- Nemani R R, Keeling C D, Hashimoto H, et al. 2003. Climate-driven increases in global terrestrial net primary production from 1982 to 1999. *Science*, 300(5625): 1560–1563.
- Niyogi D, Chang H I, Saxena V K, et al. 2004. Direct observations of the effects of aerosol loading on net ecosystem CO₂ exchanges over different landscapes. *Geophysical Research Letters*, 31(20): L20506.
- Park H S, Sohn B J. 2010. Recent trends in changes of vegetation over East Asia coupled with temperature and rainfall variations. *Journal of Geophysical Research*, 115: D14101.
- Pauli H, Gottfried M, Dullinger S, et al. 2012. Recent plant diversity changes on Europe's mountain summits. *Science*, 336(6079): 353–355.
- Peng S, Chen A, Xu L, et al. 2011. Recent change of vegetation growth trend in China. *Environmental Research Letters*, 6(4): 044027.

- Piao S L, Mohammat A, Fang J Y, et al. 2006. NDVI-based increase in growth of temperate grasslands and its responses to climate changes in China. *Global Environmental Change*, 16: 340–348.
- Qian Y, Gong D Y, Fan J W, et al. 2009. Heavy pollution suppresses light rain in China: Observations and modeling. *Journal of Geophysical Research*, 114: D00K02.
- Ramanathan V, Crutzen P, Kiehl J, et al. 2001. Aerosols, climate, and the hydrological cycle. *Science*, 294(5549): 2119–2124.
- Roderick M L, Farquhar G D, Berry S L, et al. 2001. On the direct effect of clouds and atmospheric particles on the productivity and structure of vegetation. *Oecologia*, 129(1): 21–30.
- Rosenfeld D, Rudich Y, Lahav R, et al. 2001. Desert dust suppressing precipitation: A possible desertification feedback loop. *Proceedings of the National Academy of Sciences of the United States of America*, 98(11): 5975–5980.
- Rosenfeld D, Wood R, Donner L J, et al. 2013. Aerosol cloud-mediated radiative forcing: Highly uncertain and opposite effects from shallow and deep clouds. New York: Springer Science+Business Media, 105–149.
- Rosenfeld D, Andreae M O, Asmi A, et al. 2014. Global observations of aerosol-cloud-precipitation-climate interactions. *Reviews of Geophysics*, 52(4): 750–808.
- Running S W, Justice C O, Salomonson V, et al. 1994. Terrestrial remote sensing science and algorithms planned for EOS/MODIS. *International Journal of Remote Sensing*, 15(17): 3587–3620.
- Schimel D S, House J I, Hibbard K A, et al. 2001. Recent patterns and mechanisms of carbon exchange by terrestrial ecosystems. *Nature*, 414: 169–172.
- Shao Y P, Wyrwoll K H, Chappell A, et al. 2011. Dust cycle: An emerging core theme in Earth system science. *Aeolian Research*, 2(4): 181–204.
- Solano R, Didan K, Jacobson A, et al. 2010. MODIS Vegetation Index Users' Guide (MOD13 Series) Collection 5. Tucson: University of Arizona, 35–42.
- Simmons A, Uppala S, Dee D, et al. 2006. ERA-interim: New ECMWF reanalysis products from 1989 onwards. *ECMWF News*, 110: 25–35.
- Steiner A L, Chameides W L. 2005. Aerosol-induced thermal effects increase modelled terrestrial photosynthesis and transpiration. *Tellus Series B: Chemical and Physical Meteorology*, 57(5): 404–411.
- Strada S, Unger N, Yue X. 2015. Observed aerosol-induced radiative effect on plant productivity in the eastern United States. *Atmospheric Environment*, 122: 463–476.
- Tie X, Huang R J, Dai W, et al. 2016. Effect of heavy haze and aerosol pollution on rice and wheat productions in China. *Scientific Reports*, 6(1): 29612.

- Wang X, Wu J, Chen M, et al. 2018. Field evidences for the positive effects of aerosols on tree growth. *Global Change Biology*, 24: 498-499.
- Wang X M, Ma W Y, Hua T, et al. 2017. Variation in vegetation greenness along China' s land border. *Science China Earth Sciences*, 60: 1-8.
- Wild M. 2009. Global dimming and brightening: A review. *Journal of Geophysical Research*, 114: D00D16.
- Xie J, Huang J B, Wang G Y, et al. 2010. The effects of clouds and aerosols on net ecosystem CO₂ exchange over semi-arid Loess Plateau of Northwest China. *Atmospheric Chemistry and Physics*, 10: 8205-8218.
- Yan Z B, Lin Z H, Zhang H. 2014. The relationship between the East Asian subtropical westerly jet and summer precipitation over East Asia as simulated by the IAP AGCM4.0. *Atmospheric Oceanic Science Letters*, 7(6): 487-492.
- Yin Y, Chen L. 2007. The effects of heating by transported dust layers on cloud and precipitation: A numerical study. *Atmospheric Chemistry and Physics*, 7: 3497-3505.
- Yoon J, Burrows J P, Vountas M, et al. 2014. Changes in atmospheric aerosol loading retrieved from space-based measurements during the past decade. *Atmospheric Chemistry and Physics*, 14: 26001-26041.
- Yu H B, Chin M, West J J, et al. 2013. A multi-model assessment of the influence of regional anthropogenic emission reductions on aerosol direct radiative forcing and the role of intercontinental transport. *Journal of Geophysical Research*, 118(2): 716-736.
- Yue X, Unger N, Harper K, et al. 2017. Ozone and haze pollution weakens net primary productivity in China. *Atmospheric Chemistry and Physics*, 17(9): 6073-6089.
- Zhang D F, Zakey A S, Gao X J, et al. 2009. Simulation of dust aerosol and its regional feedbacks over East Asia using a regional climate model. *Atmospheric Chemistry and Physics*, 9: 1095-1110.
- Zhang R, Ouyang Z T, Xie X, et al. 2016. Impact of climate change on vegetation growth in arid Northwest of China from 1982 to 2011. *Remote Sensing*, 8(5): 364.
- Zhang X Q, Lei Y C, Pang Y, et al. 2014. Tree mortality in response to climate change induced drought across Beijing, China. *Climatic Change*, 124(1): 179-190.
- Zhang X Y, Arimoto R, An Z S. 1997. Dust emission from Chinese desert sources linked to variations in atmospheric circulation. *Journal of Geophysical Research*, 102: 28-41.
- Zhou L M, Tucker C J, Kaufmann R K, et al. 2001. Variations in northern vegetation activity inferred from satellite data of vegetation index during 1981

to 1999. Journal of Geophysical Research, 106: 20069–20083.

Appendix

Fig. S1 shows the spatial pattern in the number of 16-d periods with a significantly positive correlation between NDVI (normalized difference vegetation index) and PAR (photosynthetically active radiation) when considering AOD (aerosol optical depth) as the control variable (right panel) or not (left panel) over the arid and semi-arid regions of Asia during growing stages (each having four or six 16-d periods). Areas with insignificant ($P > 0.05$) correlations and those with missing AOD values are not shown. Abbreviations are the same as in Figures S2–S4.

Fig. S2 shows the spatial pattern in the number of 16-d periods with a significantly positive correlation between NDVI and PRE (total precipitation) when considering AOD as the control variable (right panel) or not (left panel) over the arid and semi-arid regions of Asia during growing stages (each having four or six 16-d periods). Areas with insignificant ($P > 0.05$) correlations and those with missing AOD values are not shown.

Fig. S3 shows the spatial pattern in the number of 16-d periods with a significantly positive correlation between NDVI and SSR (surface solar radiation) when considering AOD as the control variable (right panel) or not (left panel) over the arid and semi-arid regions of Asia during growing stages (each having four or six 16-d periods). Areas with insignificant ($P > 0.05$) correlations and those with missing AOD values are not shown.

Fig. S4 shows the spatial pattern in the number of 16-d periods with a significantly positive correlation between NDVI and TEM (surface air temperature) when considering AOD as the control variable (right panel) or not (left panel) over the arid and semi-arid regions of Asia during growing stages (each having four or six 16-d periods). Areas with insignificant ($P > 0.05$) correlations and those with missing AOD values are not shown.

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv – Machine translation. Verify with original.