

Promoting the production of salinized cotton field by optimizing water and nitrogen use efficiency under drip irrigation (Postprint)

Authors: LIN En, LIU Hongguang, LI Xinxin, LI Ling, Sumera ANWAR, LIU Hongguang

Date: 2021-08-06T00:00:00+00:00

Abstract

Cotton is the main economically important crop in Xinjiang, China, but soil salinization and shortage of water and nutrients have restricted its production. A field experiment was carried out in the salinity-affected arid area of Northwest China from 2018 to 2019 to explore the effects of nitrogen and water regulation on physiological growth, yield, water and nitrogen use efficiencies, and economic benefit of cotton. The salinity levels were 7.7 (SL) and 12.5 dS/m (SM). Drip irrigation was used with low, medium and adequate irrigation levels representing 60%, 80% and 100% of cotton crop water demand, respectively, and three nitrogen applications, i.e., 206, 275 and 343 kg/hm², accounting for 75%, 100% and 125% of local N application, respectively were used. The multi-objective optimization based on spatial analysis showed that, at SL salinity, water use efficiency (WUE), nitrogen use efficiency (NUE), economic benefit and yield simultaneously reached more than 85% of their maxima at 379.18–398.32 mm irrigation and 256.69–308.87 kg/hm². At SM salinity, WUE, yield and economic benefit simultaneously reached more than 85% of their maxima when irrigation was 351.24–376.30 mm and nitrogen application was 230.18–289.89 kg/hm². NUE, yield and economic benefit simultaneously reached their maxima at 428.01–337.72 mm irrigation, and nitrogen application range was 222.14–293.93 kg/hm². The plants at SL salinity had 21.58%–46.59% higher WUE rates, 14.91%–34.35% higher NUE rates and 20.71%–35.34% higher yields than those at SM salinity. The results are of great importance for the nutrient and water management in cotton field in the arid saline area.

Full Text

Preamble

LIN En^{1,2}, LIU Hongguang^{1,2*}, LI Xinxin^{1,2}, LI Ling^{1,2}, Sumera AN-WAR³

¹ College of Water Conservancy & Architectural Engineering, Shihezi University, Shihezi 832000, China

² Xinjiang Production & Construction Group Key Laboratory of Modern Water-Saving Irrigation, Shihezi 832000, China

³ Institute of Molecular Biology and Biotechnology, The University of Lahore, Lahore 54660, Pakistan

Abstract

Cotton is the main economically important crop in Xinjiang, China, but soil salinization and shortages of water and nutrients have restricted its production. A field experiment was carried out in the salinity-affected arid area of Northwest China from 2018 to 2019 to explore the effects of nitrogen and water regulation on physiological growth, yield, water and nitrogen use efficiencies, and economic benefit of cotton. The salinity levels were 7.7 (SL) and 12.5 dS/m (SM). Drip irrigation was used with low, medium and adequate irrigation levels representing 60%, 80% and 100% of cotton crop water demand, respectively, and three nitrogen applications of 206, 275 and 343 kg/hm², accounting for 75%, 100% and 125% of local N application, respectively. Multi-objective optimization based on spatial analysis showed that, at SL salinity, water use efficiency (WUE), nitrogen use efficiency (NUE), economic benefit and yield simultaneously reached more than 85% of their maxima at 379.18–398.32 mm irrigation and 256.69–308.87 kg/hm². At SM salinity, WUE, yield and economic benefit simultaneously reached more than 85% of their maxima when irrigation was 351.24–376.30 mm and nitrogen application was 230.18–289.89 kg/hm². NUE, yield and economic benefit simultaneously reached their maxima at 428.01–337.72 mm irrigation, and nitrogen application range was 222.14–293.93 kg/hm². The plants at SL salinity had 21.58%–46.59% higher WUE rates, 14.91%–34.35% higher NUE rates and 20.71%–35.34% higher yields than those at SM salinity. These results are of great importance for nutrient and water management in cotton fields in arid saline areas.

Keywords: cotton growth; multi-objective optimization; soil salinization; water and nitrogen regulation; spatial analysis

1 Introduction

Saline soil is distributed in nearly 100 countries worldwide, covering a total area of about 1×10^9 hm², which accounts for approximately 25% of the total land area. Additionally, about 3×10^5 hm² of arable land is developing toward secondary salinization every year. Soil salinization has become one of

the main causes of land degradation in arid areas, seriously affecting sustainable agricultural development and the stability of oasis ecological environments [?, ?, ?]. Xinjiang, located in the arid and semi-arid areas of Northwest China, has low precipitation and high evaporation, with salinized area accounting for about one-third of the cultivated land area and a low land utilization rate that seriously restricts agricultural development [?, ?, ?]. Because of its high salt tolerance and effective use of saline-alkali land resources, cotton has become the largest cash crop in Xinjiang, accounting for 78% of the country's total agricultural area and 85% of the country's output [?, ?].

Water and nitrogen are important factors affecting cotton yield and quality. Reasonable irrigation and nitrogen application can significantly improve cotton yield, quality and economic benefit [?, ?]. Due to excessive soil salinization resulting in cotton growth inhibition and exacerbating premature senescence, farmers increase their nitrogen application to ensure normal growth of cotton, which causes problems such as waste of resources and environmental pollution. Therefore, in the process of developing and utilizing saline-alkali land in arid and semi-arid areas, how to give full play to the synergistic coupling effect of water and fertilizer on crop growth and improve cotton planting technology without further deteriorating the ecological environment is an important scientific issue.

The integration of water and fertilizer in drip irrigation transports water and fertilizer to the root zone of crops through the drip irrigation system for absorption and utilization. This irrigation method not only increases crop yield but also effectively reduces losses of water and fertilizer and greatly improves their utilization efficiency. Drip irrigation is an important modern agricultural technology that can alleviate the shortage of soil and water resources in arid areas [?, ?]. In recent years, many scholars in China and elsewhere have performed much research on the regulation of water and nitrogen in cotton. Within a certain threshold range, increasing water and nitrogen application has a synergistic effect on increasing cotton yield. Outside this range, yield may increase marginally or even decline [?, ?, ?, ?].

A reasonable ratio of water and nitrogen not only helps to stabilize yield but also avoids waste of water and fertilizer resources and reduces the impact of excess fertilizer on the environment. [?, ?] and [?, ?] found that water-nitrogen interaction in mild saline-alkali soil had a very significant effect on the net photosynthetic rate, transpiration rate and stomatal conductance of cotton at each growth stage, and considered that the optimal amount of nitrogen application under irrigation was 3740 m^3/hm^2 . [?, ?] found that a low irrigation level of 60% of crop water demand (ET_c) significantly inhibited any effect of fertilizer; however, an irrigation level of 90% ET_c and nitrogen application of 278 kg/hm^2 maximized cotton seed yield, WUE and net income. [?, ?] studied the water-nitrogen coupling effect on cotton under different irrigation levels in the cotton area of Gansu Province, and developed a water-nitrogen model for efficient utilization of water and nitrogen to promote cotton growth.

[?, ?] studied the coupling effect of different water and fertilizer rates on cot-

ton seed yield, WUE, fertilizer use efficiency and economic benefit in northern Xinjiang. They found that an irrigation range of 213–463 mm and fertilizer application rate (N-P₂O₅-K) of 212.59–367.51 kg/hm² resulted in cotton seed yield, economic benefit and WUE simultaneously reaching 90% of their maximum values. [?, ?] proposed an irrigation range of 328–395 mm and a nitrogen application range of 250–300 kg/hm² as the best irrigation and nitrogen levels for cotton planting. [?, ?] studied the effects of water-nitrogen coupling on cotton yield and water-nitrogen transformation in southern Xinjiang; irrigation of 395 mm and nitrogen application of 300 kg/hm² significantly increased WUE, NUE and yield of cotton. Cotton is a crop with a high nitrogen requirement, and inappropriate nitrogen application causes overgrowth and premature senescence, which affects yield and quality. [?, ?] suggested that the application of N fertilizer (0–360 kg/hm²) could alleviate salt damage, promote cotton growth and increase both cotton yield and WUE.

Salt stress on cotton results in reduced leaf area, thinner stems and lowered dry matter. [?, ?] found that with the increase in soil salinity content, cotton dry matter, boll number per plant, boll weight and lint yield decreased. [?, ?] found that cotton yield began to decrease at a soil salinity threshold of 7.7 dS/m, and decreased by 5.2% with an increase in salinity of 1.0 dS/m. [?, ?] studied the effect of water-nitrogen coupling on sunflower growth in salinized soil and found that water was the main factor affecting growth, followed by salt stress and then nitrogen. Although many studies have investigated cotton planting and growth in saline-alkali soil [?, ?], they have focused on improving soil and fertilizer use efficiencies [?, ?, ?]. Studies of water-nitrogen coupling have mainly focused on the main food crops in non-salinized areas, and there are few studies on the physiological growth of cotton under salt stress in arid areas. For cotton cultivated on saline land with limited water resources, effective irrigation and fertilization are very important for physiological growth, dry matter, quality, WUE, PFP (nitrogen partial factor productivity), NUE, yield and economic benefit.

In this paper, using spatial analysis [?, ?, ?], we carried out the horizontal projection of three-dimensional surfaces obtained by multiple regression equations of each index to find the overlap of acceptable regions of each target (\$85% maximum) and obtained the optimal combination range of each index. Based on multiple regression analysis of cotton growth indices under drip irrigation and different salt stress, this paper aimed to (1) comprehensively evaluate the yield, quality, water-saving, fertilizer saving and high efficiency of cotton combined with spatial analysis method; (2) propose an optimal water-nitrogen coupling model for the above indices; and (3) reveal the response of cotton growth to water and nitrogen regulation under salt stress.

2.1 Study Area

The field experiment was carried out in Beiquan Town (43°21'–45°20' N, 84°43'–86°35' E), Shihezi City, Xinjiang Uygur Autonomous Region, China during the

cotton-growing period of May–November in 2018 and 2019. The elevation of the site is 412 m, the average surface slope is 6% and the depth of groundwater is more than 5 m. Average annual precipitation is 198 mm, average mean temperature is 7.5°C–8.2°C, annual average wind speed is 1.5 m/s, average annual sunshine time is 2865 h, and the accumulated temperature for $\geq 10^{\circ}\text{C}$ is 3463°C. The average evaporation is 1340 mm. The soil is sandy loam with 43% saturation moisture content, 30% water holding capacity and 42% soil porosity. The physical properties of 0–100 cm soil depth are shown in Table 1.

2.2 Experimental Design and Soil Sampling

The average salinity content of 0–100 cm soil layer was measured from different areas of fields in the test area. Two plots (S1 and S2 in Figure 1 [Figure 1: see original paper]) with soil salinities of 7.4 and 12.1 dS/m were selected representing low (SL) and medium salinity (SM), respectively.

There were 18 treatments in the experiment, including two salinity levels, three N rates and three irrigation levels with three replicates per treatment. The irrigation levels used in the experiment were: low (I1), medium (I2), and adequate (I3), representing 60%, 80% and 100% of cotton ET_c , respectively. The plants were fertilized using three nitrogen levels: 206 (N1), 275 (N2), and 343 (N3) kg/hm², accounting for 75% (low), 100% (medium), and 125% (high) of local N application, respectively. Each plot size was 4.6 m $\times$ 6.0 m (width $\times$ length).

The drip irrigation under plastic mulching method was used in the field as shown in Figure 2 [Figure 2: see original paper]. The drip irrigation laterals used were single-wing labyrinth emitters with an internal diameter of 16.0 mm, 0.3 m emitter interval, and 3.2 L/h average flow rate of the emitter. A water meter, ball valve and fertilizer tank were separately installed at the head of each field experiment to accurately control water and nitrogen consumption. Cotton variety ‘Xinluzao 7’ was sown at 3–4 cm depth on 15 April 2018 and 1 May 2019. The planting density of cotton was 181,909 plants/hm².

Fertilizers of phosphate (P_2O_5 ; 100 kg/hm²), potassium (K_2O ; 100 kg/hm²) and 30% of N (20% of conventional urea and 10% of ^{15}N -urea) were applied, and the remaining 70% of N fertilizer was applied with irrigation at different times during cotton growth stages (Table 2). ^{15}N -urea, with an abundance of 5.16%, was purchased from Shanghai Research Institute, China.

In this experiment, water requirement of cotton was calculated using the following formula:

$$\text{ET}_c = \text{ET}_0 \times K_c$$

where ET_c is the water demand (mm) of the crop in a certain period; ET_0 is the reference crop evapotranspiration (mm); and K_c is the crop coefficient for the corresponding growth period. In this experiment, K_c values for cotton seedling,

flower bud, flower and boll, and boll opening stages were taken as 0.35, 0.76, 1.18 and 0.60, respectively [?, ?]. ET_0 was calculated using the Penman-Monteith equation [?, ?]:

$$ET_0 = \frac{0.408\Delta(R_n - G) + \gamma \frac{900}{T+273} u_2 (e_s - e_a)}{\Delta + \gamma(1 + 0.34u_2)}$$

where R_n is the ground net radiation evaporation equivalent ($MJ/(m^2 \cdot d)$); G is the soil heat flux ($MJ/(m^2 \cdot d)$); γ is the psychrometric constant ($kPa/^\circ C$); T is the average temperature ($^\circ C$); u_2 is the wind speed 2 m above the ground (m/s); e_s is the saturated vapor pressure (kPa); e_a is the actual vapor pressure (kPa); and Δ is the slope of the saturated vapor pressure temperature curve ($kPa/^\circ C$). All parameters in the formula were provided by the Shihezi Meteorological Bureau, China.

2.3.1 Dry Matter

Four cotton plants were randomly taken at different growth stages from each treatment. The roots, stems, leaves, buds and bolls were dried in an oven at $75^\circ C$ to a constant weight. Dry matter was determined using precision electronic weighing. Finally, cotton dry matter was calculated as follows:

dry matter per hectare = average dry matter of five cotton plants $\times$ planting density per hectare

[?, ?]

2.3.2 Boll Weight and Yield

At the harvest period, 30, 40 and 30 bolls were collected from 7 upper branches, 4-6 middle branches and 1-3 lower branches, respectively. The 100 cotton bolls were weighed and actual yield was calculated for each plot. Cotton seeds were weighed and yield per hectare was calculated by multiplying planting density per hectare [?, ?].

2.3.3 Harvest Index (HI)

The formula for HI is as follows:

$$HI = \frac{Y}{DMW}$$

where Y is the cotton seed yield (kg/hm^2); and DMW is the above-ground dry matter (kg/hm^2) [?, ?].

2.3.4 WUE

The formula for WUE is as follows:

$$WUE = \frac{Y}{ET}$$

where Y is the cotton seed yield (kg/hm²); and ET is the actual water consumption of the crop (mm):

$$ET = Pr + U + I - D - R \pm \Delta W$$

where Pr is the precipitation during growth period (mm); U is the groundwater recharge (mm); I is the irrigation (mm); D is the deep seepage amount (mm); R is the runoff (mm); and ΔW is the change in soil water storage before sowing and after harvest (mm). Before cotton planting and after harvesting, soil samples (0–100 cm) were collected from middle and bare lands of each test plot. Soil moisture was measured by soil drying method, and average value was taken. Due to the depth of groundwater, flat terrain and low rainfall, U, R and D were ignored during growth period [?, ?].

2.3.5 NUE

¹⁵N isotope tracer method was used to calculate NUE [?, ?]. The formulas for NUE are as follows:

$$N_{\text{uptake by plant part}} = \text{dry weight of plant part} \times \text{total N content in test plant part}$$

$$N_{\text{uptake by crops}} = N_{\text{uptake of aboveground part}} + N_{\text{root uptake}}$$

$$^{15}\text{N atom percentage excess} = ^{15}\text{N abundance of samples or in } ^{15}\text{N-urea} - \text{natural abundance of } ^{15}\text{N}$$

$$Ndf = \frac{^{15}\text{N atom percentage excess of test plant part}}{^{15}\text{N atom percentage excess of } ^{15}\text{N-urea}}$$

$$^{15}\text{N-urea N uptake by test plant part} = N_{\text{uptake by test plant part}} \times Ndf_{\text{test plant part}}$$

$$^{15}\text{N-urea N uptake by crops} = ^{15}\text{N-urea N of aboveground parts} + ^{15}\text{N-urea N uptake of root}$$

$$NUE = \frac{{}^{15}\text{N-urea N uptake by crops}}{{}^{15}\text{N-urea N application rate}} \times 100\%$$

Total N content of sample was determined by the Kjeldahl method and ${}^{15}\text{N}$ abundance was determined by stable isotope mass spectrometer (MAT-253).

2.3.6 Nitrogen Partial Factor Productivity (PFP)

The formula for PFP is as follows [?, ?]:

$$PFP = \frac{Y}{N}$$

where N is the total amount of applied nitrogen (kg/hm^2).

2.3.7 Cotton Fiber Quality

Cotton samples (about 20 g) were randomly collected during cotton boll opening period from each plot and micronaire value was determined at the Cotton Quality Testing Center, Ministry of Agriculture, Xinjiang Uygur Autonomous Region, China. The cotton fiber samples were analyzed using a high volume instrument fiber detector in a constant temperature and humidity laboratory [?, ?].

2.3.8 Economic Benefit (E)

The formula for E is as follows [?, ?]:

$$E = G - W - F - K - L$$

where G is the gross profit; W is the irrigation cost; F is the fertilizer cost; K is the manual machinery cost; and L is the land leasing cost (all units are USD/hm^2).

2.4 Data Processing

The value of each index was the average of three replicates in each treatment. Data Processing System was used for analysis of variance and Duncan's test, with significance considered at the $P < 0.05$ level. MATLAB 2017 and Origin 2019 were used to perform multiple regression, extreme value solution and drawing.

3.1 Boll Weight, Dry Matter, Yield and HI of Cotton

For both years, irrigation-salinity interaction had a significant effect on boll weight, dry matter, yield and HI of cotton, and nitrogen-salinity interaction had significant effects on dry matter and yield. Irrigation-nitrogen interaction had significant effects on boll weight and HI (Table 3). At SL salinity for both years, there was a significant difference in boll weight among N1, N2 and N3 treatments under I1 irrigation, but no significant difference among N1, N2 and N3 treatments under I2 or I3 irrigation. Under the same nitrogen level, boll weight initially increased and then decreased with the increase in irrigation (Fig. 3a [Figure 3: see original paper] and b). At SM salinity for both years, there were significant differences among N1, N2 and N3 treatments under I2 irrigation, and boll weight was significantly higher for N2 than for N3 treatment.

Dry matter was much lower at SM salinity than at SL salinity (Fig. 3c and d). For both years, dry matter increased with the increase in irrigation at the same salinity level. The maximum dry matter for both years at SL salinity was about 1.66-fold higher than that at SM salinity. At SL salinity for both years, dry matter initially increased and then decreased with the increase in nitrogen application at the same irrigation level, and there was a significant difference in dry matter between I1 and I3 for N1, N2 and N3 treatments. At SM salinity for both years, under all irrigations, dry matter was significantly higher for N1 than for N3 treatment, indicating that under the double stresses of water and salt, higher nitrogen application did not alleviate salt stress but aggravated stress on cotton. At the same irrigation level, the difference in dry matter among nitrogen levels was less in 2019 (Fig. 3d) than in 2018 (Fig. 3c).

At SM salinity for both years, cotton yield under different water and nitrogen treatments was lower than that of SL salinity, indicating that salt stress reduced cotton yield (Fig. 3e and f). At different salinity levels, cotton seed yield was lower under I1 than under other irrigation levels, indicating that water is the main factor restricting cotton yield under salt stress. At SL salinity for both years, yield showed an initial increase followed by a decrease with increasing nitrogen application under I1, indicating that continuous increase in nitrogen application made little contribution to yield increase. In both years, yield was higher under I2 than under I1 irrigation, indicating that water deficit affected fertilizer efficiency.

Irrigation had significant interactions with salinity and nitrogen for HI in 2018 and 2019 (Table 3). HI values at SL and SM salinities ranged from 0.35 to 0.42 and from 0.33 to 0.36, respectively (Fig. 3g and h). At SL salinity for two years, HI value initially increased and then decreased with the increase in irrigation at the same nitrogen level, indicating that the contribution of irrigation was greater to vegetative organs than to seeds. At SM salinity for two years, HI values under I1 and I3 irrigations were lower than that for I2 in 2019, indicating that both water deficit and excess water were not conducive to transport from vegetative organs to reproductive organs under salt stress. For N1 treatment,

total dry matter accumulation of plants was lower and cotton seed yield was lower than that of boll weight.

3.2 Cotton Quality

All interactions among irrigation, nitrogen and salinity significantly affected micronaire values in both years (Table 4). At SL salinity, micronaire value increased with increasing nitrogen application at the same irrigation level. For N2 and N3 treatments, micronaire value decreased with the increase in irrigation. Micronaire value was significantly higher for N3 than for N1 and N2 treatments at SL salinity. Micronaire value was significantly higher for I1 than for I2 and I3 irrigations (except for N1 in 2018). For two years, at the same irrigation level, micronaire value initially increased and then decreased with increasing nitrogen application, and was significantly higher for N2 than for N1 and N3 treatments (Fig. 4a [Figure 4: see original paper] and b). At SL salinity, micronaire value reached the peak range of 5.03–4.71 for N3 treatment (I1N3), while micronaire value reached the peak range of 4.35–4.59 for N2 treatment at SM salinity.

For both years, nitrogen-irrigation coupling had significant effects on fiber length, but the effect of nitrogen application on fiber length was not significant (Table 4). For both salinity levels, there was no significant difference in fiber length between N2 and N3 under I3, but under I1, it was significantly longer for N2 than for N1 and N3 treatments (Fig. 4c and d). Under the same nitrogen application, fiber length initially increased and then decreased with the increase in irrigation under two salinity levels. Too much or too little water and nitrogen hinder plant nitrogen metabolism, leading to premature senescence or reduced cotton fiber quality. Therefore, appropriate amounts of water and nitrogen will optimize fiber quality. For both years, the maximum value of cotton fiber length at SL salinity was 28.25 mm, which was 6.24% higher than the minimum fiber length. At SM salinity, the maximum fiber length was 26.21 mm, which was 8.36% higher than the minimum fiber length.

3.3 WUE, PFP and NUE

For both years, interaction among irrigation, nitrogen and salinity had a significant effect on WUE, PFP and NUE (Table 5). At SL salinity, WUE decreased with the increase in irrigation within the same nitrogen level. The effect of nitrogen application on WUE was $N2 > N1 > N3$ under I1 and I3 irrigations (Fig. 5a [Figure 5: see original paper]). At SM salinity, WUE initially increased and then decreased with rising nitrogen application under I1 and I2 irrigations, and WUE was significantly higher in 2019 than in 2018 (Fig. 5a and b). Under I3 irrigation, WUE was significantly higher for N2 than for N1 and N3 treatments. Among treatments, the highest WUE at SL and SM salinities occurred under I1 and N2 treatments, with values of 0.81–1.68. At the same nitrogen level, only low water and high fertilizer levels improved WUE.

At SL salinity, the maximum value of PFP occurred at I2 and N1 treatment

(Fig. 5c and d). At the same irrigation level, PFP decreased gradually with the increase in nitrogen application. For N1 and N2 treatments, PFP initially increased and then decreased with the increase in irrigation. At SL salinity, there were significant differences in PFP among N1, N2 and N3 treatments under the same irrigation level. High water and low fertilizer were beneficial for improving PFP.

NUE significantly differed under I2 and I3 irrigations at SL salinity (Fig. 5e and f). Under I3 irrigation, NUE increased with an increase in nitrogen application. At the same nitrogen level, NUE initially increased and then decreased with the increase in irrigation. At SM salinity, there were significant differences in NUE among different nitrogen treatments, and the highest NUE was found under I2 and N2 treatments. Proper irrigation can increase NUE of cotton. The appropriate amount of water and nitrogen can increase fertilizer efficiency. Too much nitrogen will increase soil salinity and aggravate secondary salinization; conversely, less nitrogen will not meet the demand for cotton physiological growth.

3.4 Economic Benefit

The gross costs in 2018 and 2019 at SL salinity were 5449.00–7399.36 and 840.89–1141.88 USD/hm², respectively (Table 6). At SL salinity, the highest gross cost was increased by 14.6% and 49.0% compared with the lowest gross cost, the lowest economic benefits were 840.89 and 571.37 USD/hm², and the highest economic benefits were 91691.16 and 2268.70 USD/hm² in 2018 and 2019, respectively. At SM salinity, the lowest economic benefits were 152.63 and 196.13 USD/hm² in 2018 and 2019, respectively, and correspondingly the highest economic benefits were 1252.36 and 1251.64 USD/hm². These results showed that appropriate management of water and fertilizer can affect gross profit and double economic benefit. The difference between the highest and the lowest economic benefits at SM salinity was 4.14 times that of SL salinity. Although the cost of land leasing accounts for a small proportion of the total investment, salt stress leads to great economic losses for cotton, which is one of the main reasons that farmers prefer to rent land with high cost and low salinity content.

4 Multi-Objective Optimization

The economic benefit, WUE, NUE and PFP were selected as optimization objectives for the sustainable development of cotton field due to their important effects on yield. Binary quadratic equations based on the least square method were developed for regression analysis under different salinity levels, taking water and nitrogen inputs as independent variables, and yield, economic benefit, WUE, NUE and PFP as dependent variables (Table 7). The effects of water and nitrogen inputs on all dependent variables were significant, and the determination coefficients were all above 0.86. Taking I1 and I3 as the upper and lower limits of irrigation, respectively, and N1 and N3 as the upper and lower limits

of nitrogen application, respectively, the maximum values of each equation were obtained by MATLAB extremum problem-solving method, and the maximum irrigation and nitrogen application levels were obtained.

At SL salinity, the yield, economic benefit and PFP simultaneously reached the maximum with 410.00–414.00 mm irrigation. At SM salinity, the yield, economic benefit and PFP reached the maximum with 394.00–402.63 mm irrigation. WUE and PFP reached the maximum with 301.12–402.63 mm irrigation and 247.93–264.71 kg/hm² nitrogen application (Table 8). Under the same irrigation level, nitrogen application varied greatly, thus it was not possible to have the above indicators simultaneously reaching their maxima under the same irrigation and nitrogen levels.

Through normalized treatment of each index, we used spatial analysis to evaluate the acceptable ranges of yield, economic benefit, WUE, NUE and PFP. We found the reasonable acceptance range of 85% at SL and SM salinities, and comprehensively analyzed the contour projection of 85% of the maximum values of WUE, quality, NUE, economic benefit and yield (Figs. 6 and 7). At SL salinity, when WUE, NUE, economic benefit and yield reached the maximum, the range for irrigation was 379.18–398.32 mm and nitrogen application was 256.69–308.87 kg/hm² (Fig. 8 [Figure 8: see original paper]). At SM salinity, WUE, yield and economic benefit reached their maxima at the ranges of 351.24–376.30 mm irrigation and 230.18–289.89 kg/hm² nitrogen application. The range of irrigation for the maximum NUE, yield and economic benefit was 415.14–437.87 mm and the range of nitrogen application was 222.14–293.93 kg/hm².

5.1 Boll Weight, Dry Matter Accumulation, Cotton Yield and HI

In arid and semi-arid areas, low water and nitrogen input affects cotton growth; however, excessive irrigation and nitrogen application not only reduce WUE, increase costs and waste resources, but also damage soil structure. Furthermore, excessive fertilization results in groundwater pollution, contributes to soil salinization, and increases the negative effect of soil salinity on plant performance [?, ?]. Water and nitrogen management of cotton can be achieved by coordinating irrigation and nitrogen application to provide the best opportunity to save water and achieve high yield and fiber quality.

At low salinity (7.4 dS/m), higher boll weight, plant dry weight and cotton yield were obtained compared with higher salinity (12.1 dS/m). Seed yields were 27.6% and 27.3% lower at SM than at SL in 2018 and 2019, respectively. Boll weights were 19.6% and 12.2% lower at SM than at SL in 2018 and 2019, respectively. The reduction in boll weight is due to the negative effects of salinity as indicated by the significant negative correlation between soil salinity and cotton yield and its components. Previous studies showed that lower cotton yield was related to decreased lint weight per boll and boll number per plant with the increase in soil salinity content [?, ?]. The fewer bolls per plant might

be attributed to the greater chance of shedding due to nutrient imbalance caused by high soil salinity [?, ?].

In our study, the effect of salt stress on cotton boll weight fluctuated with changes in nitrogen application and irrigation levels. Cotton yield is an important index to evaluate different water and nitrogen levels under salt stress. Nevertheless, our results indicated that the adverse effects of soil salinity on cotton growth could be alleviated by nitrogen application, but excessive nitrogen application at SM salinity inhibited the expansion of cotton growth, leading to a decrease in boll weight and seed yield as indicated by the least dry matter for N3 and the highest dry matter for N2. The highest seed yield was found for N2, with 5003 kg/hm² in 2018 and 5027 kg/hm² in 2019. Seed yield was 9.0% and 6.8% less for N3 than for N1 in 2018 and 2019, respectively. These results suggested that over-fertilization increased the negative effects of soil salinity on plant performance and might contribute to soil salinization [?, ?].

The results of this study were consistent with those of previous studies in which aboveground dry matter initially increased and then decreased with the increase in nitrogen application. The 252–255 kg/hm² nitrogen rate is proposed for cotton under saline soil if the water supply meets crop water requirements [?, ?]. The reason for this phenomenon may be that excessive nitrogen application inhibited the absorption of other nutrients by cotton organs, which then affected the formation of dry matter [?, ?].

To determine the optimal dose of nitrogen, consideration of irrigation is particularly relevant. Similar levels of nitrogen could produce differences in nitrogen utilization and yield depending on water supply [?, ?]. The amount of irrigation needs to be considered before determining the optimal nitrogen requirement of the crop.

The accumulation of cotton dry matter is the basis for obtaining high-yield and high-quality cotton. Under drip irrigation conditions, increasing the amount of irrigation increased the dry matter accumulation of cotton, but too much water leads to overgrowth. Under different salinity levels, seed yield of cotton was lower under I1 than under other irrigation levels, indicating that water is the main factor restricting cotton yield under salt stress. Excessive irrigation lowered the crop rhizosphere soil osmotic potential and raised crop transpiration, plant active transport and nutrient absorption, and thus dry matter of aboveground tissues increased [?, ?]. These results implied that a moderate irrigation application that was 80% of ETc for cotton under mulched drip irrigation provided advantages in terms of yield and economic benefit in the arid region.

Under salt stress in arid areas, the effect of water-nitrogen interaction on cotton yield was significant for dry matter, boll weight and yield. This result may be caused by the coupling effect of fertilizer by irrigation and transferring fertilizer by water. A higher cotton yield can be achieved by promoting the effect of water and nitrogen. Previous results revealed that cotton yield with drip irrigation increased with the increase in irrigation and fertilizer application within

a certain range and then yield declined beyond a certain threshold. Overall, in the two-year study, the maximum dry matter and yield at SL and SM salinities occurred under I2 and N2 treatment. The appropriate combination of water and nitrogen is required to properly utilize resources, and a proper irrigation level could improve plant capability to utilize the applied nitrogen [?, ?]. Furthermore, fiber length, NUE and HI also reached the maximum under I2 and N2 treatment. However, the highest N (I1) and the lowest irrigation (N3) treatment produced the lowest dry matter, boll weight and seed yield, indicating that less water supply reduces the availability of nitrogen to cotton. [?, ?] found that less water supply lessened yield and biomass accumulation due to stomatal closure and decreased photosynthesis. At the same time, under the highest irrigation (I3) and the lowest nitrogen (N1) treatment, the lowest WUE, NUE and micronaire values occurred because high irrigation could cause hypoxia and nutrient leaching. However, dry matter was the highest under I3 and N2 treatment in both years, which indicated that higher irrigation improved vegetative growth and inhibited reproductive growth [?, ?]. Thus, suitable water and fertilizer can increase cotton yield, which is consistent with other results [?, ?].

HI values were lower under I1 and I3 irrigations than under I2 irrigation, indicating that both water deficit and excess irrigation are not conducive to transport nutrients from vegetative organs to reproductive organs of cotton under salt stress [?, ?]. In our study, at SL salinity, the maximum yield (6218.20 kg/hm²) was obtained when the amount of irrigation and nitrogen application was 410.17 mm and 295.47 kg/hm², respectively. At SM salinity, yield reached the maximum of 4705.62 kg/hm² when the amount of irrigation and nitrogen application was 402.63 mm and 264.71 kg/hm², respectively. Cotton yield was 32.14% higher at SL than at SM salinity.

5.2 Cotton Quality

Fiber quality is one of the main indicators determining the economic benefit of cotton besides yield. Cotton quality is mainly composed of micronaire value and fiber length. High micronaire values (>5.0) indicate irregular coarse fibers, while low values (<3.5) show immature fibers [?, ?]. In this study, micronaire value and fiber length decreased with the increase in soil salinity. Previous studies of the effect of soil salinity on final quality of cotton fiber are not consistent. [?, ?] found that soil salinity had a significant positive correlation with micronaire value, but no significant correlation with fiber length. [?, ?] found that fiber length uniformly increased with the increase in soil salinity, but fiber strength and maturity decreased at high salinity. Our study showed that nitrogen increased fiber length and micronaire value at a medium dose, while fiber length and micronaire value decreased with the increase in irrigation. The formation of fiber length is determined by the rate and duration of fiber elongation. [?, ?] found that increasing nitrogen before the flowering stage of cotton increased the duration of fiber elongation and maximized elongation rate. In this study, micronaire value reached the highest under I1 irrigation, but the lowest under I3

irrigation. Previous studies showed that micronaire value and irrigation levels are negatively correlated [?, ?, ?]. Higher irrigation or nitrogen causes immature fiber [?, ?]. This result could be attributed to more vegetative growth and less reproductive growth under higher irrigation level [?, ?].

5.3 WUE, PFP and NUE

Under the same irrigation levels, nitrogen application can effectively improve WUE. Research has shown that increasing the amount of fertilizer can improve WUE within a certain threshold [?, ?]. The reason is that nitrogen application promotes cotton growth and increases available water in the soil. The available water moves to the vicinity of the cotton root system, increasing absorption efficiency and further improving the effective use of soil water [?, ?]. Nitrogen application also had a great effect on PFP. PFP decreased with the increase in nitrogen application under different irrigation levels. Similar results were reported by [?, ?] and [?, ?], who found the maximum PFP occurred at the lowest nitrogen application. Although PFP was highest for N1 treatment under all irrigation levels, it could not meet production requirements.

To obtain higher economic benefit of cotton planting on saline-alkali land, it is necessary to improve NUE and reduce nitrogen loss. Our results showed that NUE was maximum when maximum seed yield and fiber length were achieved at the same irrigation and nitrogen levels. Similarly, [?, ?] found that under low and medium salinities, increasing nitrogen amount up to an appropriate level could significantly increase nitrogen uptake of cotton. However, excessive nitrogen application did not increase uptake. Under salt stress, the absorption and utilization of nitrogen by cotton depended on the amount of water [?, ?]. NUE decreased if water was excessive, possibly because nitrogen was leached into deep soil so that crop roots could not absorb it.

5.4 Economic Benefit

The results showed that in salinized areas, irrigation water accounted for a small proportion of the total cost, and so farmers' awareness of water-saving was not strong, resulting in waste of water resources that was not conducive to sustainable development. Economic benefit was 1.78–3.45 and 1.18–1.53 times higher under I2 irrigation than under I1 and I3 irrigations at SL salinity, and 2.93–4.44 and 1.82–2.05 times higher at SM salinity. Thus irrigation had a great impact on economic benefit. Moderate increase in nitrogen could significantly increase yield under SM salinity and I1 irrigation, but this was not obvious for I3 irrigation, and there was no obvious or even a downward trend at SL salinity with excessive nitrogen application.

6 Conclusions

A significant negative correlation between soil salinity and cotton yield was found in this study. Thus an optimal water and nitrogen combination could

increase cotton dry matter accumulation and increase boll weight, thereby improving cotton yield under salt stress. The results of multiple regression and spatial analysis showed that cotton yield, economic benefit, WUE, NUE and PFP could reach more than 85% of their maximum values. The maximum WUE, NUE, economic benefit and yield were attained at irrigation ranges of 379.18–398.32 mm and 256.69–308.87 kg/hm² nitrogen application at SL salinity, and 428.01–337.72 mm irrigation and 222.14–293.93 kg/hm² nitrogen application at SM salinity.

Acknowledgements

The study was supported by the National Natural Science Foundation of China (U1803244, 51669029, 2020DB01) and the National Key Research and Development Program of China (2016YFC0501406).

References

- \cite{Adiku S G K, Renger M, Wessolek G, et al. 2001. Simulation of the dry matter production and seed yield of common beans under varying soil water and salinity conditions. *Agricultural Water Management*, 47(1): 55–68.}
- [?, ?, ?, ?, ?]
- \cite{Badr M A, Abou-Hussein S D, El-Tohamy W A. 2016. Tomato yield, nitrogen uptake and water use efficiency as affected by planting geometry and level of nitrogen in an arid region. *Agricultural Water Management*, 169: 90–97.}
- [?, ?, ?, ?, ?]
- \cite{Chen W P, Hou Z N, Wu L S, et al. 2010. Effects of salinity and nitrogen on cotton growth in arid environment. *Plant and Soil*, 326(1): 61–73.}
- \cite{Choudhary R L, Minhas P S, Wakchaure G C, et al. 2020. Effect of IW: CPE-based irrigation scheduling and N-fertilization rate on yield, water and N-use efficiency of wheat (*Triticum aestivum*). *Agricultural Research*, 10(2): 1–12.}
- \cite{Dağdelen N, Başal H, Yılmaz E, et al. 2009. Different drip irrigation regimes affect cotton yield, water use efficiency and fiber quality in western Turkey. *Agricultural Water Management*, 96(1): 111–120.}
- \cite{Deng Z, Bai D, Zhai G L, et al. 2013. Effects of water and nitrogen regulation on the yield and water and nitrogen use efficiency of cotton in south Xinjiang, Northwest China under plastic mulched drip irrigation. *Chinese Journal of Applied Ecology*, 24(9): 2525–2532. (in Chinese)}
- \cite{Du H Y, Sun X L, Liu X W, et al. 2017. Effects of optimized fertilization on plant growth and soil nutrients supply in saline fields. *Acta Agriculturae Boreali-Sinica*, 32(1): 220–225. (in Chinese)}

- \cite{Feng L, Zhang J P, Sun C T, et al. 2014. Responses of cotton growth and water physiological indices to salt stress at different growing stages. Chinese Journal of Ecology, 33(5): 1195-1199. (in Chinese)}
- \cite{García-López J, Lorite I J, García-Ruiz R, et al. 2016. Yield response of sunflower to irrigation and fertilization under semi-arid conditions. Agricultural Water Management, 176: 151-162.}
- \cite{He H J, Wang Z H, Zheng X R, et al. 2017. Effects of water-nitrogen coupling on growth and yield of cotton under mulch drip irrigation. Xinjiang Agricultural Sciences, 54(11): 1983-1989. (in Chinese)}
- \cite{Hou Z A, Chen W P, Li X, et al. 2009. Effects of salinity and fertigation practice on cotton yield and ^{15}N recovery. Agricultural Water Management, 96(10): 1483-1489.}
- \cite{Hu Y J, Wang X B, Zhao Q S, et al. 2010. Effects of sea ice water irrigation and different fertilizations on soil salinity dynamics and cotton. Transactions of the Chinese society of Agricultural Engineering, 26(9): 20-27. (in Chinese)}
- \cite{Ierna A, Pandino G, Lombardo S, et al. 2011. Tuber yield, water and fertilizer productivity in early potato as affected by a combination of irrigation and fertilization. Agricultural Water Management, 101(1): 35-41.}
- \cite{Jiang Y J, Zheng D M, Lv S Q, et al. 1995. Influence of soil salt content on cotton yield and quality. Xinjiang Agricultural Sciences, 3: 116-118. (in Chinese)}
- \cite{Kljun A, El-Dessouky H M, Benians T A, et al. 2014. Analysis of the physical properties of developing cotton fibres. European Polymer Journal, 51: 57-68.}
- \cite{Krishnamurthy S L, Gautam R K, Sharma P C, et al. 2016. Effect of different salt stresses on agro-morphological traits and utilisation of salt stress indices for reproductive stage salt tolerance in rice. Field Crops Research, 190: 26-33.}
- \cite{Leal A J, Piatì G L, Leite R C, et al. 2020. Nitrogen and mepiquat chloride can affect fiber quality and cotton yield. Revista Brasileira de Engenharia Agrícola e Ambiental, 24(4): 238-243.}
- \cite{Lin E, Liu H G, He X L, et al. 2019. Water-nitrogen coupling effect on drip-irrigated dense planting of dwarf jujube in an extremely arid area. Agronomy, 9(9): 561.}
- \cite{Li P L, Zhang F C. 2010. Regulation effect of water and nitrogen on cotton biomass and yield under different drip irrigation patterns. Journal of Applied Ecology, 21(11): 2814-2820. (in Chinese)}
- \cite{Li P L, Zhang F C. 2012. Effects of water and nitrogen coupling on cotton nitrogen absorption and utilization under alternate root partition drip.

Transactions of the Chinese Society of Agricultural Engineering, 28(S1): 112-116. (in Chinese)}

[?, ?, ?, ?, ?]

\cite{Luo J, Chen B L, Xiang G R, et al. 2017. Effects of organic manure on dry matter accumulation, nutrient absorption and cotton production on salt affected land. Soil and Fertilizer Sciences in China, (2): 107-113. (in Chinese)}

\cite{Min W, Hou Z A, Ma L J, et al. 2014. Effects of water salinity and N application rate on water-and N-use efficiency of cotton under drip irrigation. Journal of Arid land, 6(4): 454-467.}

\cite{Pinnamaneni S R, Anapalli S S, Ruixiu S U I, et al. 2021. Effects of irrigation and planting geometry on cotton (*Gossypium hirsutum* L.) fiber quality and seed composition. Journal of Cotton Research, 4(1): 1-11.}

\cite{Sobrinho F P C, Guerra H O, Araújo W P, et al. 2015. Fiber quality of upland cotton under different irrigation depths. Revista Brasileira de Engenharia Agrícola e Ambiental, 19(11): 1057-1063.}

\cite{Sui R, Byler R K, Delhom C D. 2017. Effect of nitrogen application rates on yield and quality in irrigated and rainfed cotton. Journal of Cotton Science, 21(2): 113-121.}

\cite{Tang L S, Li Y, Zhang J H. 2005. Physiological and yield responses of cotton under partial root zone irrigation. Field Crops Research, 94(2-3): 214-223.}

\cite{Thompson T L, Doerge T A. 1996. Nitrogen and water interactions in sub-surface trickle-irrigated leaf lettuce II. Agronomic, economic, and environmental outcomes. Soil Science Society of America Journal, 60(1): 168-173.}

\cite{Umbetaev I, Bigaraev O, Baimakhanov K. 2015. Effect of soil salinity on the yield of cotton in Kazakhstan. Russian Agricultural Sciences, 41(4): 222-224.}

\cite{Wang H D, Wu L F, Cheng M H, et al. 2018. Coupling effects of water and fertilizer on yield, water and fertilizer use efficiency of drip-fertigated cotton in northern Xinjiang, China. Field Crops Research, 219: 169-179.}

\cite{Wang R S, Kang Y H, Wan S Q, et al. 2011. Salt distribution and the growth of cotton under different drip irrigation regimes in a saline area. Agricultural Water Management, 100(1): 58-69.}

\cite{Wang R S, Kang Y H, Wan S Q, et al. 2012. Influence of different amounts of irrigation on salt leaching and cotton growth under drip irrigation in an arid and saline area. Agricultural Water Management, 110: 109-117.}

\cite{Wang R S, Wan S Q, Sun J X, et al. 2018. Soil salinity, sodicity and cotton yield parameters under different drip irrigation regimes during saline wasteland reclamation. Agricultural Water Management, 209: 20-31.}

\cite{Wang Z H, Zhu Y K, Zhang J Z, et al. 2018. Effects of water and nitrogen fertilization on physiological characteristics and yield of cotton under drip irrigation in mildly salinized soil. Transactions of the Chinese Society for Agricultural Machinery, 49(6): 296-308. (in Chinese)}

\cite{Wang Z M, Wang B G, Jin M G. 2013. Water and salt stress to cotton in the condition of drip irrigation under mulch. Geological Science and Technology Information, 32(3): 166-170. (in Chinese)}

\cite{Wu L F, Zhang F C, Zhou H M, et al. 2014. Effect of drip irrigation and fertilizer application on water use efficiency and cotton yield in North of Xinjiang. Transactions of the Chinese Society of Agricultural Engineering, 30(20): 137-146. (in Chinese)}

\cite{Wu L F, Zhang F C, Wang H D, et al. 2015. Simulation of cotton leaf area index under deficit irrigation in Xinjiang. Transactions of the Chinese Society for Agricultural Machinery, 46(1): 249-258. (in Chinese)}

\cite{Xue Z, Shi H B, Guo Y, et al. 2007. Effects of coupling of soil water and fertilizer on the growth of sunflower during its seeding period in saline soil. Transactions of the Chinese Society of Agricultural Engineering, (3): 91-94.}

\cite{Ye Q B, Zhang H C, Wei H Y, et al. 2007. Effects of nitrogen fertilizer on nitrogen use efficiency and yield of rice under different soil conditions. Frontiers of Agriculture in China, 1(1): 30-36.}

\cite{Zhang D M, Li W J, Xin C S, et al. 2012. Lint yield and nitrogen use efficiency of field-grown cotton vary with soil salinity and nitrogen application rate. Field Crops Research, 138: 63-70.}

\cite{Zhang H, Li D S, Zhou Z G, et al. 2017. Soil water and salt affect cotton (*Gossypium hirsutum* L.) photosynthesis, yield and fiber quality in coastal saline soil. Agricultural Water Management, 187: 112-121.}

\cite{Zhang L, Chen B L, Zhang G W, et al. 2013. Effect of soil salinity, soil drought, and their combined action on the biochemical characteristics of cotton roots. Acta Physiologiae Plantarum, 35(11): 3167-3179.}

\cite{Zhao W Q, Wang Y H, Zhou Z G, et al. 2012. Effect of nitrogen rates and flowering dates on fiber quality of cotton (*Gossypium hirsutum* L.). Journal of Experimental Agriculture International, 2(2): 133-159.}

[?, ?, ?, ?, ?, ?]

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv – Machine translation. Verify with original.