

Determination of neutron-skin thickness using configurational information entropy

Authors: Ma, Chun-Wang, Liu, Yi-Pu, Wei, Hui-Ling, Pu, Jie, Cheng, Kai-Xuan, Wang, Yu-Ting, Ma, Chun-Wang

Date: 2021-12-31T15:17:25+00:00

Abstract

Configurational information entropy (CIE) theory was employed to determine the neutron skin thickness of neutron-rich calcium isotopes. The nuclear density distributions and fragment cross-sections in 350 MeV/u $^{40-60}\text{Ca} + ^9\text{Be}$ projectile fragmentation reactions were calculated using a modified statistical abrasion-ablation model. CIE quantities were determined from the nuclear density, isotopic, mass, and charge distributions. The linear correlations between the CIE determined using the isotopic, mass, and charge distributions and the neutron skin thickness of the projectile nucleus show that CIE provides new methods to extract the neutron skin thickness of neutron-rich nuclei.

Full Text

Preamble

Determination of neutron-skin thickness using configurational information entropy

Chun-Wang Ma^{1,2,†}, Yi-Pu Liu¹, Hui-Ling Wei¹, Jie Pu¹, Kai-Xuan Cheng¹, and Yu-Ting Wang¹

¹School of Physics, Henan Normal University, Xinxiang 453007, China

²Key Laboratory Optoelectronic Sensing Integrated Application of Henan Province, Henan Normal University, Xinxiang 453007, China

Configurational information entropy (CIE) theory was employed to determine the neutron skin thickness of neutron-rich calcium isotopes. The nuclear density distributions and fragment cross sections in 350 MeV/u $^{40-60}\text{Ca} + ^9\text{Be}$ projectile fragmentation reactions were calculated using a modified statistical abrasion-ablation model. CIE quantities were determined from the nuclear density, isotopic, mass, and charge distributions.

The linear correlations between the CIE determined using the isotopic, mass, and charge distributions and the neutron skin thickness of the projectile nucleus show that CIE provides new methods to extract the neutron skin thickness of neutron-rich nuclei.

Keywords: Neutron-skin thickness, Configurational information entropy, Cross section distribution, Projectile fragmentation

INTRODUCTION

Next-generation radioactive nuclear beam facilities will provide new opportunities to explore extreme nuclei near and beyond the drip lines. Nuclei with a large neutron excess can form exotic neutron skin or halo structures, which have attracted significant interest experimentally and theoretically for the past 30 years. The neutron skin thickness is defined as $\delta_{np} = \delta_n - \delta_p$, which denotes the difference between the point neutron and point proton root-mean-square (RMS) radii of a nucleus. Many methods have been developed to experimentally determine the neutron skin thickness. However, most of these methods involve indirect measurements and are model-dependent. Typical approaches include the reaction cross section (σ_R), charge-changing cross section (σ_{cc}) [?, ?], electric dipole polarizability [?], photon multiplicity [?], the π^-/π^+ ratio or Σ^-/Σ^+ [?, ?], $^3\text{H}/^3\text{He}$ ratio [?], and α decay half-life time [?].

Projectile fragmentation reactions, which constitute the main experimental approach for studying rare isotopes, are particularly suitable for determining neutron skin thickness due to the distinct experimental phenomena induced by the neutron skin structure [?, ?]. Examples include isospin effects in the isotopic cross section [?], neutron-abrasion cross section (σ_{nabr}) [?], neutron removal cross section [?], mirror nuclei ratio or isobaric ratio [?], and the isoscaling parameter (α) [?]. Parity-violating electron scattering (PVS) is the only model-independent method for determining neutron skin thickness. Reference [?] reports a theoretical investigation of PVS for ^{48}Ca and ^{208}Pb , and a theoretical investigation using a Bayesian approach is given in Ref. [?]. Determining the neutron skin thickness of ^{48}Ca and ^{208}Pb is presently of significant interest and is listed in the U.S. 2015 Long Range Plan for Nuclear Science [?]. The lead radius experiment (PREX) has been previously used to determine the neutron skin thickness of ^{208}Pb [?]. A recent PREX result indicates a much thicker neutron skin than previously predicted [?]. Determining the neutron skin thickness of nuclei near the neutron drip line remains an important research objective and one of the most interesting topics in the new era of radioactive beam facilities.

Information entropy theory was established by C.E. Shannon [?]. The theory makes it possible to transform variables in a system into an exact information quantity [?] and has been used in various applications [?, ?]. The first application of information entropy theory in heavy-ion reactions can be traced to the study of nuclear liquid-gas transition in nuclear multifragmentation [?]. Recent studies have extended it to investigate the information entropy carried by

a single fragment produced in projectile fragmentation reactions, revealing the scaling phenomenon of fragments covering a wide range of neutron excess [?]. Configurational information entropy (CIE) was developed to quantify the information entropy of a physical distribution [?], which connects the dynamical and informational contents of a physical system with localized configurations. Many applications of CIE methods can be found in Korteweg-de Vries (KdV) solitons, compact astrophysical systems, and scalar glueballs (see a brief introduction in Ref. [?]), theoretical research of new Higgs boson decay channels [?], deploying heavier eta meson states in AdS/QCD [?], confinement/deconfinement transition in QCD [?], quarkonium in a finite density plasma [?], time evolution in physical systems [?, ?], etc.

In projectile fragmentation reactions, fragment distributions show a sensitive dependence on changes in neutron density [?], which makes it possible to determine the neutron skin thickness of neutron-rich nuclei. In this study, the CIE method was adopted to quantify the CIE of nuclear density and fragment distributions in projectile fragmentation reactions. The analyzed data were generated using a modified statistical abrasion-ablation (SAA) model, which is known to be a good model for describing the fragment cross sections of projectile fragmentation reactions [?, ?].

II. THEORIES

A. Modified Statistical Abrasion-Ablation Model

The modified statistical abrasion-ablation (SAA) model [?, ?] can be used for projectile fragmentation reactions at both intermediate and high energies, improving upon the original SAA model by Brohm and Schmidt [?]. In quasi-free nucleon-nucleon collisions, the reaction is described as a two-step process. In the initial stage, the nucleons are described by a Glauber-type model as “participants” and “spectators,” where the participants interact strongly in an overlapping region between the projectile and target, while the spectators move virtually undisturbed [?]. In the second stage, the excitation energy is compared to the separation energies of protons, neutrons, and α particles to determine the type of particle the prefragment can emit according to $\min(s_p, s_n, s_\alpha)$. After the de-excitation calculation, the cross sections of final fragments comparable to the measured fragments were obtained. The description of the modified SAA model is presented in Refs. [?, ?, ?].

The colliding nuclei are composed of many parallel tubes oriented along the beam direction. Their transverse motion is neglected, and the interactions between the tube pairs are independent. For a specific pair of interacting tubes, the absorption of the projectile neutrons and protons is assumed to follow a binomial distribution. At a given impact parameter b , the transmission probabilities of neutrons (protons) of an infinitesimal tube in the projectile are calculated using

$$t_i(s - b) = \exp\{-[\mathcal{D}_T^n(s - b)\sigma_{ni} + \mathcal{D}_T^p(s - b)\sigma_{pi}]\},$$

where \mathcal{D}_T is the normalized integrated nuclear density distribution of the target along the beam direction for protons $\int d^{2s} \mathcal{D}_T^p = Z_T$ and neutrons $\int d^{2s} \mathcal{D}_T^n = N_T$ (N_T and Z_T are the neutron and proton numbers of the target, respectively). s and b are defined in a plane perpendicular to the beam direction, and $\sigma_{i'i}$ denotes the free-space nucleon-nucleon cross sections ($i' = n$ for neutrons and $i' = p$ for protons) [?].

The average absorbed mass in the infinitesimal tube limit at a given b is

$$\langle \Delta A(b) \rangle = \int d^{2s} \mathcal{D}_T^n(s) [1 - t_n(s - b)] + \int d^{2s} \mathcal{D}_P^p(s) [1 - t_p(s - b)].$$

For a specific fragment, the production cross section can be calculated using

$$\sigma(\Delta N, \Delta Z) = \int d^{2b} P(\Delta N, b) P(\Delta Z, b),$$

where $P(\Delta N, b)$ and $P(\Delta Z, b)$ are the probability distributions of the abraded neutrons and protons at a given impact parameter b , respectively. $\sigma(\Delta N, \Delta Z)$ is the residual fragment after the abrasion stage, which is called the prefragment. The excitation energy of the prefragment is calculated as $E^* = 13.3 \langle \Delta A(b) \rangle$ MeV, where $\langle \Delta A(b) \rangle$ is the number of abraded nucleons from the projectile, and 13.3 MeV is the mean excitation energy owing to an abraded nucleon [?].

Fermi-type density distributions were adopted for protons and neutrons in a nucleus, as shown in the equation below:

$$\rho_i(r) = \frac{\rho_0^i}{1 + \exp\left(\frac{r - C_i}{t_i/4.4}\right)}, \quad i = n, p$$

where ρ_0^i is the normalization constant of neutrons ($i = n$) or protons ($i = p$), t_i is the diffuseness parameter, and C_i is half the density radius of the neutron or proton density distribution.

B. Configurational Information Entropy Method

To determine the quantity of CIE incorporated in the fragment distributions, definitions of CIE were introduced. For a system with spatially localized clusters, when performing the CIE analysis, a set of functions $f(x) \in L^2(\mathbb{R})$ and their Fourier transforms $F(k)$ obey Plancherel's theorem [?]:

$$\int_{-\infty}^{\infty} |f(x)|^2 dx = \int_{-\infty}^{\infty} |F(k)|^2 dk,$$

where $f(x)$ is square-integrable and bounded. The model fraction $f(k)$ is defined as:

$$f(k) = \frac{|F(k)|^2}{\int |F(k)|^2 d^d k}$$

where the integration is over all k , $F(k)$ is defined, and d is the number of spatial dimensions.

The model fraction $f(k)$ measures the relative weight of a given mode k . The quantity of CIE $S_C[f]$ is defined as a summation of the Shannon information entropy of $f(k)$ [?]:

$$S_C[f] = - \sum f_m \ln(f_m).$$

Thus, the quantity of CIE contains information about configurations compatible with certain constraints of a given physical system. If all the modes k have the same mass, then $f_m = 1/N$. The discrete configuration entropy reaches a maximum at $S_C = \ln N$. If there is only one mode, $S_C = 0$.

Continuous CIE can also be defined for continuous distributions, such as the nuclear density distribution. For non-periodic functions in the interval (a, b) ,

$$S_C[f] = - \int \tilde{f}(k) \ln[\tilde{f}(k)] d^d k,$$

where $\tilde{f}(k) = f(k)/f(k)_{\max}$ [$f(k)_{\max}$ is the maximum fraction]. The normalized function $\tilde{f}(k)$ guarantees that $\tilde{f}(k) \leq 1$ for all k modes, and $\tilde{f}(k) \ln \tilde{f}(k)$ denotes the CIE density.

III. RESULTS AND DISCUSSION

The 350 MeV/u $^{A_p}\text{Ca} + ^9\text{Be}$ reactions were calculated using the modified SAA model (A_p refers to even mass numbers from 40 to 60). The cross sections of fragments with Z ranging from 3 to 20 were obtained. For clarity, only part of the calculated results are shown in the figures.

Figure 1 shows the Fermi-type nuclear density distributions and their fast Fourier transformation (FFT) spectra. An obvious increase in ρ_n is observed from ^{40}Ca to ^{60}Ca , whereas the opposite trend is observed for ρ_p . A two-peak structure is evident in the FFT spectra, where the second peak is lower than the first. The difference between the neutron and proton density distributions $\Delta\rho = \rho_n - \rho_p$ is also shown. For ^{40}Ca , $\Delta\rho$ is very small, while $\Delta\rho$ increases as the neutrons in the projectile increase. The peaks in the FFT spectra of ρ_n and ρ_p are not clearly shown. Based on the FFT spectra $f(k)$, the CIE of ρ_n , ρ_p , and $\Delta\rho$ can be determined using Eq. (7), which are denoted by $S_{\rho_n}^C[f]$, $S_{\rho_p}^C[f]$, and $S_{\Delta\rho}^C[f]$, respectively.

The isotopic cross section (σ_Z) distributions produced in the 350 MeV/u ${}^{A_p}\text{Ca}$ + ${}^9\text{Be}$ reactions are plotted in Fig. 2. In panels (a_i), from $Z_{fr} = 7$ to 20, the isotopic cross-section distributions in the ${}^{40-60}\text{Ca}$ reactions are similar for fragments with small Z_{fr} , while a shift to the neutron-rich side is observed for larger Z_{fr} . The symmetric Gaussian-like shape of the isotopic distribution is altered by the enhanced cross sections of neutron-rich fragments in neutron-rich reaction systems, showing the isospin effect in fragment production induced by the increased neutron density on the surface of neutron-rich nuclei [?]. The FFT spectra of the isotopic distributions are shown in Fig. 2 (b_j). In each FFT spectrum, only one peak is observed. The amplitudes of the FFT spectra of different isotopic distributions decrease as the projectile becomes more neutron-rich, except for $Z = 20$. Based on the FFT spectra of σ_Z distributions, the quantities of CIE of the isotopic distributions are determined according to Eq. (7), which is denoted by $S_{\sigma_Z}^C[f]$.

The correlation between the $S_C[f]$ of the density distribution and δ_{np} of the projectile nucleus is shown in Fig. 3(a). Both $S_{\rho_n}^C[f]$ and $S_{\rho_p}^C[f]$ decrease linearly with an increase in δ_{np} from ${}^{40}\text{Ca}$ to ${}^{60}\text{Ca}$. $S_{\Delta\rho}^C[f]$ also decreases linearly with increasing δ_{np} , except for ${}^{40}\text{Ca}$. The correlation between the $S_{\sigma_Z}^C[f]$ of different Z_{fr} and δ_{np} of the projectile nuclei are plotted in panel (b). The $S_{\sigma_Z}^C[f]$ of isotopes from $Z_{fr} = 10$ to 18 are also found to decrease with the increasing δ_{np} of the projectile nucleus. The $S_{\sigma_Z}^C[f]$ of fragments near the projectile nucleus was more sensitive to the change in δ_{np} .

The mass yield (σ_A) distributions in the 350A MeV ${}^{A_p}\text{Ca}$ + ${}^9\text{Be}$ reactions are shown in Fig. 4. In each reaction, the mass yield increases with the A_{fr} of the fragment until it is close to the projectile nucleus. In different reactions, a very similar trend of mass distribution was observed, which decreased with the increasing mass number of the projectile nucleus. The corresponding quantities of CIE were determined from σ_A distributions, which are labeled as $S_{\sigma_A}^C[f]$. The correlation between $S_{\sigma_A}^C[f]$ and δ_{np} for projectile nuclei is shown in Fig. 4(b). Except for the bend point formed at δ_{np} for ${}^{42}\text{Ca}$ owing to the transition from proton-skin to neutron skin, the $S_{\sigma_A}^C[f] \sim \delta_{np}$ correlation was found to be linear for reactions with $A_p \geq 44$.

The charge cross section is defined as the summation of the isotopic cross sections $\sigma_C = \sum_o \sigma(A_o, Z)$. The charge cross section distributions in the 350A MeV ${}^{A_p}\text{Ca}$ + ${}^9\text{Be}$ reactions are shown in Fig. 5. Similar trends of σ_C distribution were observed, comparable to those of σ_A . The determined CIE of σ_C distributions, labeled as $S_{\sigma_C}^C[f]$, were linearly correlated to the neutron skin thickness of the projectile nuclei.

IV. SUMMARY

With the vast opportunities for very asymmetric nuclei available in the new era of radioactive ion beam facilities, neutron skin thickness is one of the most

important questions in nuclear physics. In this study, CIE theory is adopted to quantify the information entropy incorporated in nuclear density distributions and fragment cross-section distributions in 350 MeV/u $^{40-60}\text{Ca} + ^9\text{Be}$ projectile fragmentation reactions calculated using the modified SAA model.

CIE quantities of nuclear density distributions ($S_{\rho_{n,p}}^C[f]$ and $S_{\Delta\rho}^C[f]$), isotopic cross-section distributions ($S_{\sigma_z}^C[f]$), mass cross-section distributions ($S_{\sigma_A}^C[f]$), and charge cross-section distributions ($S_{\sigma_C}^C[f]$) were determined. The correlations between $S_{\rho_p}^C[f] \sim \delta_{np}$, $S_{\Delta\rho}^C[f] \sim \delta_{np}$, $S_{\rho_n}^C[f] \sim \delta_{np}$, $S_{\sigma_z}^C[f] \sim \delta_{np}$, $S_{\sigma_A}^C[f] \sim \delta_{np}$, and $S_{\sigma_C}^C[f] \sim \delta_{np}$ were also investigated. For neutron-rich calcium projectiles, obvious linear dependences of $S_{\rho_n}^C[f]$ on δ_{np} were observed. The $S_{\sigma_z}^C[f]$ of fragments with different Z_{fr} is shown to linearly depend on the δ_{np} of the projectile nucleus. It is found that, if the isotopic distribution is sensitive to isospin effects in the projectiles, the extracted $S_{\sigma_z}^C[f]$ will also be sensitive to their δ_{np} . Good linear correlations between $S_{\sigma_A}^C[f]$, $S_{\sigma_C}^C[f]$, and δ_{np} of the projectile nucleus were also observed. It is suggested that, from the viewpoint of CIE, the isotopic/mass/charge distributions in the projectile fragmentation reaction may be good probes for determining the neutron-skin thickness of neutron-rich nuclei.

The CIE approach transfers the experimental distributions to quantified parameters and provides information probes for determining the properties of a system. From the $S_{\sigma_z}^C[f] \sim \delta_{np}$, $S_{\sigma_A}^C[f] \sim \delta_{np}$, and $S_{\sigma_C}^C[f] \sim \delta_{np}$ correlations, it was observed that the CIE determined from isotopic, mass, and charge distributions decrease with increasing neutron skin thickness, respectively, and they exhibit good linear correlations.

The determination of neutron skin thickness, particularly for nuclei near the neutron drip line, is limited by the unavailability of effective probes. The linear correlation between the CIE and neutron-skin thickness of neutron-rich nuclei provides new approaches to determine the neutron skin thickness of the projectile nucleus by measuring the fragment distributions in projectile fragmentation reactions.

In this work, the simple description of the nuclear density of a projectile nucleus is difficult to apply to nuclei with magic numbers, as well as those with large shape distortion. Further improvements should concentrate on the inputs of nuclear densities, such as results obtained from density functional theories and relativistic mean field theories, to better investigate the effects of nuclear density on fragment cross-section distributions and related CIE quantities.

AUTHOR CONTRIBUTIONS

All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by Yi-Pu Liu, Hui-Ling Wei and Chun-Wang Ma. The first draft of the manuscript was written by Yi-Pu

Liu and Chun-Wang Ma and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

REFERENCES

- [1] A. Ozawa, T. Moriguchi, T. Ohtsubo, et al., Charge-changing cross sections of ^{30}Ne , $^{32,33}\text{Na}$ with proton target. *Phys. Rev. C* 89, 044602 (2014). <https://doi.org/10.1103/PhysRevC.89.044602>
- [2] S. Bagchi, R. Kanungo, W. Horiuchi et al., Neutron skin and signature of the $N = 14$ shell gap found from measured proton radii of $^{17-22}\text{N}$. *Phys. Lett. B* 790, 251-256 (2019). <https://doi.org/10.1016/j.physletb.2019.01.024>
- [3] X. Roca-Maza, X. Viñas, M. Centelles et al., Neutron skin thickness from the measured electric dipole polarizability in ^{68}Ni , ^{120}Sn , and ^{208}Pb . *Phys. Rev. C* 92, 064304 (2015). <https://doi.org/10.1103/PhysRevC.92.064304>
- [4] G.F. Wei, Probing the neutron-skin thickness by photon production from reactions induced by intermediate-energy protons. *Phys. Rev. C* 92, 014614 (2015). <https://doi.org/10.1103/PhysRevC.92.014614>
- [5] B. A. Li, Probing the High Density Behavior of Nuclear Symmetry Energy with High Energy Heavy-Ion Collisions. *Phys. Rev. Lett.* 88, 192701 (2002). <https://doi.org/10.1103/PhysRevLett.88.192701>
- [6] D.C. Zhang, H.G. Cheng, Z. Q. Feng, Hyperon Dynamics in Heavy-Ion Collisions near Threshold Energy. *Chin. Phys. Lett.* 38, 092501 (2021). <https://doi.org/10.1088/0256-307X/38/9/092501>
- [7] T.Z. Yan, S. Li, Impact parameter dependence of the yield ratios of light particles as a probe of neutron skin. *Nucl. Sci. Tech.* 30, 43 (2019). <https://doi.org/10.1007/s41365-019-0572->
- [8] N. Wan, C. Xu, Z.Z. Ren, Exploring the sensitivity of α -decay half-life to neutron skin thickness for nuclei around ^{208}Pb . *Nucl. Sci. Tech.* 28, 22 (2017). <https://doi.org/10.1007/s41365-017-0022->
- [9] H. Yu, D. Q. Fang, Y.G. Ma, Investigation of the symmetry energy of nuclear matter using isospin-dependent quantum molecular dynamics. *Nucl. Sci. Tech.* 31, 61 (2020). <https://doi.org/10.1007/s41365-020-00766-x>
- [10] J. Liu, C. Gao, N. Wan, C. Xu, Basic quantities of the equation of state in isospin asymmetric nuclear matter. *Nucl. Sci. Tech.* 32, 57 (2021). <https://doi.org/10.1007/s41365-021-00955-2>
- [11] C.W. Ma, S.S. Wang, Isospin dependence of projectile fragmentation and neutron-skin thickness of neutron-rich nuclei. *Chin. Phys. C* 35, (2011). <https://doi.org/10.1088/1674-1137/35/11/007>
- [12] C.W. Ma, Y. Fu, D.Q. Fang et al., A possible experimental observable for the determination of neutron skin thickness. *Chin. Phys. B* 17, 1216-1222

(2008). <https://doi.org/10.1088/1674-1056/17/4/011>

[13] T. Aumann, C.A. Bertulani, F. Schindler et al., Peeling Off Neutron Skins from Neutron-Rich Nuclei: Constraints on the Symmetry Energy from Neutron-Removal Cross Sections. *Phys. Rev. Lett.* 119, 262501 (2017). <https://doi.org/10.1103/PhysRevLett.119.262501>

[14] C. W. Ma, H.L. Wei, Y.G. Ma, Neutron-skin effects in isobaric yield ratios for mirror nuclei in a statistical abrasion-ablation model. *Phys. Rev. C* 88, 044612 (2013). <https://doi.org/10.1103/PhysRevC.88.044612>

[15] Z.T. Dai, D.Q. Fang, Y.G. Ma et al., Effect of neutron skin thickness on projectile fragmentation. *Phys. Rev. C* 91, 034618 (2015). <https://doi.org/10.1103/PhysRevC.91.034618>

[16] J. Liu, Z. Ren, C. Xu et al., Electroweak charge density distributions in parity-violating electron scattering. *Phys. Rev. C* 88, 054321 (2013). <https://doi.org/10.1103/PhysRevC.88.054321>

[17] J. Xu, Constraining Isovector Nuclear Interactions with Giant Dipole Resonance and Neutron Skin in ^{208}Pb from a Bayesian Approach. *Chin. Phys. Lett.* 38, 042101 (2021). <https://doi.org/10.1088/0256-307X/38/4/042101>

[18] Nuclear Science Advisory Committee, Reaching for the Horizon: The 2015 U.S. Long Range Plan for Nuclear Science, (2014).

[19] S. Abrahamyan, Z. Ahmed, H. Albataineh et al., Measurement of the Neutron Radius of ^{208}Pb through Parity Violation in Electron Scattering. *Phys. Rev. Lett.* 108, 112502 (2012). <https://doi.org/10.1103/PhysRevLett.108.112502>

[20] B.T. Reed, F.J. Fattoyev, C.J. Horowitz et al., Implications of PREX-2 on the Equation of State of Neutron-Rich Matter. *Phys. Rev. Lett.* 126, 172503 (2021). <https://doi.org/10.1103/PhysRevLett.126.172503>

[21] C.E. Shannon, *Bell Syst. Tech. J.* 27, 379-429; 623-656 (1948).

[22] A.E. Bernardini, R. da Rocha, Entropic information of dynamical AdS/QCD holographic models. *Phys. Lett. B* 762, 107-115 (2016). <https://doi.org/10.1016/j.physletb.2016.09.023>

[23] C.W. Ma, Y.G. Ma, Isobaric yield ratio difference and Shannon information entropy. *Prog. Part. Nucl. Phys.* 99, 120-158 (2018). <https://doi.org/10.1016/j.ppnp.2018.01.002>

[24] J. Chen, D. Keane, Y.G. Ma et al., Antinuclei in heavy-ion collisions, *Phys. Rep.* 760, 1 (2018).

[25] Y.G. Ma, Application of Information Theory in Nuclear Liquid Gas Phase Transition. *Phys. Rev. Lett.* 83, 3617 (1999). <https://doi.org/10.1103/PhysRevLett.83.3617>

[26] C.W. Ma, H.L. Wei, S.S. Wang et al., Isobaric yield ratio difference and Shannon information entropy. *Phys. Lett. B* 742, 19-22 (2015). <https://doi.org/10.1016/j.physletb.2015.01.015>

[27] C.W. Ma, Y.D. Song, C.Y. Qiao et al., A scaling phenomenon in the difference of Shannon information uncertainty of fragments in heavy-ion collisions. *J. Phys. G: Nucl. Part. Phys.* 43, 045102 (2016). <https://doi.org/10.1088/0954-3899/43/4/045102>

[28] C.W. Ma, H.L. Wei, X.Q. Liu et al., Nuclear fragments in projectile fragmentation reactions. *Prog. Part. Nucl. Phys.* 121, 103991 (2021). <https://doi.org/10.1016/j.ppnp.2021.103911>

[29] M. Gleise, N. Stamatopoulos, Entropic measure of localized configurations: Kinks, bounces, and bubbles. *Phys. Lett. B* 713, 109-113 (2012). <https://doi.org/10.1016/j.physletb.2012.05.064>

[30] A. Alves, A.G. Dias, R. da Silva, The 7% rule: A maximum entropy prediction on new decays of the Higgs boson. *Nucl. Phys. B* 959, 115137 (2020). <https://doi.org/10.1016/j.nuclphysb.2020.115137>

[31] R. da Rocha, Deploying heavier meson states: Configurational entropy hybridizing AdS/QCD. *Phys. Lett. B* 814, 136112 (2021). <https://doi.org/10.1016/j.physletb.2021.136112>

[32] N.R.F. Braga, O.C. Junqueira, Configuration entropy holographic QCD. *Phys. Lett. B* 814, 136082 (2021). <https://doi.org/10.1016/j.physletb.2021.136082>

[33] N.R.F. Braga, R. da Mata, Configuration entropy for quarkonium in a finite density plasma. *Phys. Rev. D* 101, 105016 (2020). <https://doi.org/10.1103/PhysRevD.101.105016>

[34] F. Li, G. Chen, The evolution of information entropy components in relativistic heavy-ion collisions. *Eur. Phys. J. A* 56, 167 (2020). <https://doi.org/10.1140/epja/s10050-020-00169-x>

[35] J.C. Zhang, W. K. Ren, N.D. Jin, Rescaled Range Permutation Entropy: A Method for Quantifying the Dynamical Complexity of Extreme Volatility in Chaotic Time Series. *Chin. Phys. Lett.* 37, 090501 (2020). <https://doi.org/10.1088/0256-307X/37/9/090501>

[36] C.W. Ma, X.M. Bai, J. Yu et al., Neutron density distributions of neutron-rich nuclei studied with the isobaric yield ratio difference. *Eur. Phys. J. A* 50, 139 (2014). <https://doi.org/10.1140/epja/i2014-14139-1>

[37] C.W. Ma, S.S. Wang, H.L. Wei et al., Re-examination of Finite-Size Effects in Isobaric Yield Ratios Using a Statistical Abrasion-Ablation Model. *Chin. Phys. Lett.* 30, 052101 (2013). <https://doi.org/10.1088/0256-307X/30/5/052101>;

[38] C.W. Ma, H.L. Wei, Y.G. Ma, Neutron-skin effects in isobaric yield ratios in a statistical abrasion-ablation model. *Phys. Rev. C* 88, 044612 (2013). <https://doi.org/10.1103/PhysRevC.88.044612> for mirror nuclei

[39] D.Q. Fang, W.Q. Shen, J. Feng et al., Isospin effect of fragmentation reactions induced by intermediate energy heavy ions and its disappearance. *Phys.*

Rev. C 61, 044610 (2000). <https://doi.org/10.1103/PhysRevC.61.044610>

[40] C.W. Ma, J.Y. Wang, H.L. Wei et al., Isospin dependence of projectile-like fragment production at intermediate energies. Phys. Rev. C 79, 034606 (2009). <https://doi.org/10.1103/PhysRevC.79.034606>

[41] T. Brohm, K.-H. Schmidt, Statistical abrasion of nucleons from realistic nuclear-matter distributions. Nucl. Phys. A 569, 821-832 (1994). [https://doi.org/10.1016/0375-9474\(94\)90386-7](https://doi.org/10.1016/0375-9474(94)90386-7)

[42] Y. Eisenberg, Interaction of Heavy Primary Cosmic Rays in Lead. Phys. Rev. 96, 1378 (1954). <https://doi.org/10.1103/PhysRev.96.1378>

[43] J.J. Gaimard, K.-H. Schmidt, A reexamination of the abrasion-ablation model for the description of fragmentation reaction. Nucl. Phys. A 531, 709-745 (1991). [https://doi.org/10.1016/0375-9474\(91\)90748-U](https://doi.org/10.1016/0375-9474(91)90748-U)

[44] X. Cai, J. Feng, W. Shen et al., In-medium nucleon-nucleon cross section and its effect on total nuclear reaction cross section. Phys. Rev. C 58, 572-575 (1998). <https://doi.org/10.1103/PhysRevC.58.572>

[45] W. Rudin, Real and Complex Analysis, McGraw-Hill, 1987, Singapore

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv –Machine translation. Verify with original.