
AI translation · View original & related papers at
chinarxiv.org/items/chinaxiv-202203.00075

Effects of vegetation near-soil-surface factors on runoff and sediment reduction in typical grasslands on the Loess Plateau, China (Postprint)

Authors: LI Panpan, WANG Bing, Yanfen Yang, LIU Guobin, YANG Yanfen

Date: 2022-03-24T16:48:18+00:00

Abstract

Abstract: Vegetation near-soil-surface factors can protect topsoil from erosion, however, their contributions to the reduction of soil erosion, especially under natural rainfall events, have not been systematically recognized. This study was performed to quantify the effects of near-soil-surface factors on runoff and sediment under natural rainfall events on grasslands dominated by *Bothriochloa ischaemum* (Linn.) Keng (BI grassland) and *Artemisia gmelinii* Thunb. (AG grassland) in two typical watersheds on the Loess Plateau, China in 2018. By successive removal of the plant canopy, litter, biological soil crusts (BSCs) and plant roots, we established five treatments including plant roots, plant roots+BSCs, plant roots+BSCs+litter, intact grassland and bare land in each grassland type. In total, twenty runoff plots ($5 \text{ m} \times 3 \text{ m}$) with similar slopes and aspects were constructed in the two types of grasslands. Results showed that plant canopy, litter and roots reduced runoff, while BSCs, which swelled in the presence of water, increased runoff. In contrast, all of these factors reduced sediment yield. In addition, the reductions in runoff and sediment yield increased with I₃₀ (maximum 30-min rainfall intensity) for each vegetation near-soil-surface factor except for BSCs. Among these factors, plant canopy had the largest contribution to runoff reduction, accounting for 48.8% and 39.9% in the BI and AG grasslands, respectively. The contributions of these vegetation near-soil-surface factors to sediment yield reduction were similar (21.3%-29.9%) in the two types of grasslands except for BSCs in the AG grassland (10.3%). The total reduction in runoff in the BI grassland (70.8%) was greater than that in the AG grassland (53.1%), while the reduction in sediment yield was almost the same in both grasslands (97.4% and 96.7%). In conclusion, according to the effects of different vegetation near-soil-surface factors on runoff and sediment production, our results may provide more complete insight and scientific basis into the effects of various vegetation related factors in controlling soil erosion.

Full Text

Abstract

Vegetation near-soil-surface factors can protect topsoil from erosion; however, their contributions to reducing soil erosion, especially under natural rainfall events, have not been systematically recognized. This study quantified the effects of near-soil-surface factors on runoff and sediment yield under natural rainfall events in grasslands dominated by *Bothriochloa ischaemum* (Linn.) Keng (BI grassland) and *Artemisia gmelinii* Thunb. (AG grassland) in two typical watersheds on the Loess Plateau, China, in 2018. By successively removing plant canopy, litter, biological soil crusts (BSCs), and plant roots, we established treatments including plant roots only, plant roots + BSCs, plant roots + BSCs + litter, intact grassland, and bare land for each grassland type. In total, twenty runoff plots (5 m × 3 m) with similar slopes and aspects were constructed in the two grassland types. Results showed that plant canopy, litter, and roots reduced runoff, while BSCs, which swelled in the presence of water, increased runoff. In contrast, all these factors reduced sediment yield. Additionally, the reductions in runoff and sediment yield increased with I30 (maximum 30-min rainfall intensity) for each vegetation near-soil-surface factor except for BSCs. Among these factors, plant canopy had the largest contribution to runoff reduction, accounting for 48.8% and 39.9% in the BI and AG grasslands, respectively. The contributions of these vegetation near-soil-surface factors to sediment yield reduction were similar (21.3%–29.9%) in the two grassland types except for BSCs in the AG grassland (10.3%). The total reduction in runoff in the BI grassland (70.8%) was greater than that in the AG grassland (53.1%), while the reduction in sediment yield was almost the same in both grasslands (97.4% and 96.7%). In conclusion, based on the effects of different vegetation near-soil-surface factors on runoff and sediment production, our results provide more complete insight and a scientific basis for understanding the role of various vegetation-related factors in controlling soil erosion.

Keywords: vegetation near-soil-surface factor; grassland; arid and semi-arid; soil erosion; Loess Plateau

Citation: LI Panpan, WANG Bing, YANG Yanfen, LIU Guobin. 2022. Effects of vegetation near-soil-surface factors on runoff and sediment reduction in typical grasslands on the Loess Plateau, China. *Journal of Arid Land*, 14(3): 325–340. <https://doi.org/10.1007/s40333-022-0007-y>

1 Introduction

Soil erosion is a serious ecological problem worldwide. Many laboratory and field experiments have confirmed that planting or preserving vegetation is an effective way to prevent soil loss, especially in vulnerable areas. Due to differences in vegetation types, vegetation distribution, and community structure, the effects of vegetation on soil erosion vary [?, ?, ?]. Compared with woodlands

with low coverage, grasslands with high coverage or plant density may reduce annual runoff by 50% and annual soil loss by 92% [?]. The various effects of vegetation on soil erosion are generally related to vegetation near-soil-surface factors such as plant canopy, litter, biological soil crusts (BSCs), root system, and soil properties [?, ?, ?].

Vegetation near-soil-surface factors include both aboveground and belowground components, with the former including plant canopy, litter, and BSCs, and the latter mainly including plant roots [?]. Generally, plant canopy can intercept raindrops and reduce kinetic energy, consequently protecting the soil surface from splash erosion. Zhou et al. [?] reported that for every 1% increase in plant canopy, the amount of soil loss decreased by $456 \text{ t}/(\text{km}^2 \cdot \text{a})$ and soil erosion tended to be zero when coverage reached 52%. Plant litter improves soil surface roughness, and the combination of plant stems and litter forms a series of small dams, which decreases runoff rate and dissipates overland flow energy [?]. Furthermore, plant litter can be incorporated into the topsoil and thus improve soil structure, enhance soil infiltration ability, and promote soil stability [?]. Plant litter incorporated into soil may decrease soil rill erodibility by 24.3%-34.8% in natural grasslands or woodlands [?]. In contrast to plant canopy and litter, BSCs show both positive and negative effects on soil erosion.

On the one hand, BSCs may promote soil water infiltration by increasing soil porosity and soil aggregate stability, thereby reducing overland flow velocity and erosive forces [?]. On the other hand, BSCs may form an impermeable seal after absorbing water and thereby reduce infiltration and increase the overland flow rate [?]. Wang et al. [?] reported that BSCs can be broken by overland flow and increase soil detachment capacity when the shear stress is greater than 11 Pa (i.e., the failure point). Plant roots are distributed throughout the soil mass and form a root-soil composite that strengthens soil shear strength during root growth [?]. Plant roots also have strong direct effects on soil factors, increasing organic matter content, soil cohesion, and water-stable aggregates and decreasing rill erodibility [?]. Wang et al. [?] summarized the role of the root system in consolidating soil as involving two mechanisms: physical binding and chemical bonding. They showed that the root bonding effect accounted for a quarter of the total reducing effects of roots on soil erosion. In addition, root type greatly affects soil erosion. In general, plants with fibrous root systems are more effective in reducing soil erosion than those with tap roots [?, ?]. As reported by Wang and Zhang [?], the ability of taproots to reduce soil detachment capacity is only two-thirds that of fibrous roots. In addition, soil properties (especially physical characteristics) have strong effects on soil erosion. An increase in soil bulk density reduces soil porosity and hinders infiltration, thus increasing overland flow and soil loss. Soil organic matter can enhance the formation of soil aggregates, improving soil structure and reducing soil erosion [?].

However, most studies assessing the effects of vegetation on soil erosion have considered only one or two of the above factors. In addition, popular soil erosion prediction models ignore these parameters. For example, the cover-management

factor has been used to reveal the effects of vegetation cover and management measures on soil erosion in the Universal Soil Loss Equation (USLE) and Revised Universal Soil Loss Equation (RUSLE) [?]. However, the cover-management factor has been estimated at watershed and regional scales mainly by land-use valuation methods and vegetation coverage calculations using remote sensing images [?]. These methods do not consider the effects of some factors (such as plant roots or litter) on soil erosion, which may limit model precision. Moreover, it remains unknown to what extent soil loss can be reduced by a single vegetation near-soil-surface factor and what contribution each factor makes to the reduction of soil erosion under the participation of all the above-described vegetation near-soil-surface factors. Many studies have been carried out in this area and have achieved remarkable results. However, to our knowledge, only Wang et al. [?] and Li and Pan [?] took into consideration all vegetation near-soil-surface factors to determine the effects of vegetation on soil erosion. As reported by Wang et al. [?], plant roots, BSCs, and litter reduced the soil detachment capacity by $0.001\text{--}0.094 \text{ kg}/(\text{m}^2 \cdot \text{s})$, with contributions ranging from 17.1% to 58.7%. Li and Pan [?] showed that plant roots were the primary contributor to reduced soil erosion, with an average contribution of 84%, while for runoff reduction, the contribution of aboveground vegetation near-soil-surface factors (plant canopy and litter) was almost 86%. However, they used either scouring modeling experiments or simulated rainfall on planted grassland (for two years) rather than natural rainfall on natural grassland. This may have caused their results to be unrepresentative of field conditions. A systematic study of all vegetation near-soil-surface factors (plant canopy, plant roots, BSCs, and litter) under natural rainfall and natural grassland conditions may help us better understand their impacts on soil erosion and improve the prediction accuracy of soil erosion models.

The Loess Plateau is among the areas with the most severe soil erosion in China. The Chinese government implemented a series of ecological construction projects to solve ecological problems in the Loess Plateau, such as the Grain for Green Project in 1999. Due to the implementation of this project, the vegetation coverage of the Loess Plateau increased from 31.6% in 1999 to 63.2% in 2018, and grassland became the main land-use type on the plateau, covering an area of $2.6 \times 10^5 \text{ km}^2$ and accounting for 41.7% of the total plateau area in 2010 [?]. The near-soil-surface factor of the grassland in the Loess Plateau is worth investigating due to the 20-year restoration. Wang et al. [?] reported that plant litter accumulation in a grassland abandoned for 24 years was 0.20 kg/m^2 , which was 6.7 times that of grassland abandoned for 1 year. Plant litter reduced soil erosion by 8.1% and 13.3% in the grasslands abandoned for 1 and 24 years, respectively, compared with that of bare land [?]. In addition, BSCs developed during vegetation restoration and covered up to 60%–70% of the soil surface on grassland of the hilly Loess Plateau. Furthermore, BSCs reduced sediment yield by 92% in revegetated grassland, and soil loss could be completely inhibited by BSCs in the later successional stage [?]. Additionally, plant species composition and abundance have changed accordingly during natural succession

(1–30 years) on grassland of the Loess Plateau [?]. Different grass species with various root morphologies and structures may have diverse effects on soil erosion. For example, in a previous study, the reduction in soil detachment capacity caused by Gramineae plants was 1.6–1.9 times that by Compositae plants due to differences in their root systems [?]. Above all, previous studies concentrated on limited near-soil-surface factors and failed to understand the overall effects of vegetation on soil erosion.

In the present study, *Bothriochloa ischaemum* (Linn.) Keng and *Artemisia gmelinii* Thunb. are typical herbaceous plant species on grassland of the Loess Plateau, China. By observing natural rainfall events in the field, we studied the effects of vegetation near-soil-surface factors on soil erosion in these two types of grasslands. We hypothesized that all vegetation near-soil-surface factors reduce runoff and sediment, in which plant canopy performs better in reducing runoff whereas plant roots are most effective at decreasing sediment yield. The aims of this study were to: (1) investigate the reductions in runoff and sediment yield caused by vegetation near-soil-surface factors; (2) distinguish the relationships between runoff or sediment yield reductions and rainfall characteristics; and (3) quantify the relative contribution of each vegetation near-soil-surface factor to runoff and sediment yield reduction.

2.1 Study Area

This study was carried out on two typical grasslands in two watersheds in Ansai District, Shaanxi Province, in 2018. The grassland in the Fangta Watershed (8.4 km²; 36°47' 27" N, 109°16' 46" E) was dominated by *A. gmelinii* (abbreviated AG), and the grassland in the Xianmangou Watershed (50.6 km²; 36°44' 25" N, 109°17' 22" E) was dominated by *B. ischaemum* (abbreviated BI; Fig. 1). The altitude of the study area ranges from 470 to 1800 m a.s.l. The loess-dominated hilly and gully region has a warm temperate semi-arid climate. The annual mean temperature is 8.8°C, and the mean annual precipitation is 400–500 mm. Heavy rainfall events often occur in summer (July–September) and result in severe soil erosion. This region belongs to the forest–grassland transition zone, and Compositae and Gramineae are typical herbaceous plant taxa in this area. The original land-use type of both grasslands was sloping farmland, with a slope gradient of 60%, and both grassland areas have experienced natural succession for approximately 35 years since they were abandoned. The physical properties of the test soil are shown in Table S1.

Fig. 1 (a) Distribution of two experimental plots; (b) BI grassland is the grassland with *B. ischaemum* as the dominant species; (c) AG grassland is the grassland with *A. gmelinii* as the dominant species. A schematic diagram of T1–T4 treatments is shown in Figure S1. The treatments named T0, T1, T2, T3, T4, and T5 indicate bare land, plant roots, plant roots + BSCs, plant roots + BSCs + litter, intact grassland, and dead roots, respectively, and T5 is not analyzed in the study. BSCs, biological soil crusts.

2.2 Experimental Design

Four treatments including bare land (T0), plant roots (T1), plant roots + BSCs (T2), plant roots + BSCs + litter (T3), and intact grassland (T4) were set up to investigate and distinguish different vegetation near-soil-surface factors in controlling soil erosion. For bare land (T0), the 0–30 cm topsoil was flipped, sieved, and packed into the plots to a bulk density, which was used as the control. Two runoff plots were designed for each treatment as replicates, and a total of ten runoff plots were constructed on each type of grassland in June 2018. The runoff plots were surrounded by 40-cm-high PVC sheets buried in the soil to a depth of 20 cm. Each runoff plot had an area of 15 m² (5 m in length and 3 m in width).

2.3 Rainfall Characteristics

The rainfall amount (P) and duration (t) were recorded by a tipping bucket self-recording rainfall gauge (RG3-M, HOBO, USA). During the observation period (July–September in 2018), a total of 20 and 15 rainfall events were detected by monitoring equipment in the BI and AG grasslands, respectively (Fig. S2). Among them, 10 and 7 rainfall events produced runoff and sediment, respectively, and these events were termed effective rainfall events. To eliminate the disturbance caused by the building of runoff plots, we did not collect the first runoff and sediment samples. Therefore, a total of 15 effective rainfall events (9 rainfall events in the BI grassland and 6 in the AG grassland) were analyzed in this study (Table S2). After each rainfall event, P and t data were collected and used as the basic data for the next analysis. Other rainfall parameters, namely, average rainfall intensity (IA), maximum 30-min rainfall intensity (I₃₀), rainfall kinetic energy (E), and rainfall erosivity (R), were calculated to describe the rainfall characteristics using Rain Record software.

2.4 Measurements of Runoff and Sediment Yield

For each plot, the runoff and sediment yield were collected after each rainfall event by using a tank that was fixed to the bottom of the runoff plot. The volume of runoff-sediment mixture in the tank (VSR-P) was measured with a ruler. Next, the muddy water in each tank was fully mixed to ensure that it was uniform and sampled with a 1000-mL plastic bottle, and this procedure was repeated twice. The volume of muddy water in each plastic bottle was measured (VSR-b) and weighed (MSR-b). Then, the muddy water was moved into an aluminum box, the sediment was allowed to settle, the supernatant was drained, and the sediment was dried (105°C, 24 h) to obtain the weight of sediment yield (MS-b) in the plastic bottle. The ratio of sediment to runoff was used to calculate the amount of runoff (VR-p) and the sediment yield (MS-p) of each runoff plot.

SR-bS-bS-pSR-pSR-bSR-bMMMVVM SR-pS-pR-pwMMV R-PRDVA S-PSYMA

In this study, runoff generation was expressed by runoff depth (RD, mm), and soil loss was measured as sediment yield per unit area (SY, g/m²), calculated by the following equations: where A is the projected area of the runoff plot (15 m²).

2.5 Contributions of Near-Soil-Surface Factors to Soil Erosion Reduction

In this study, the contribution to runoff or sediment reduction was applied to represent the inhibitory effects of near-soil-surface factors on soil erosion. The calculation process was as follows: The participating near-soil-surface factors differed among the treatments, and the runoff and sediment yield varied. Therefore, the reductions in runoff and sediment yield caused by near-soil-surface factors of the plant canopy, plant roots, BSCs, and litter were calculated by Equations 5 and 6.

1ΔTiTiTiRRR 1ΔTiTiTiSSS ΔCR100%TiTiTiRR ΔCS100%TiTiTiSS %1004404 TTTTRRRRCR%10044

The actual contribution of each factor *i* (plant canopy, plant roots, BSCs, and litter) to reducing runoff (CR, %) or sediment (CS, %) was computed by Equations 7 and 8. The actual contributions of T4 (intact grassland) to runoff (CRT4, %) and sediment yield (CST4, %) reductions were expressed by Equations 9 and 10.

2.6 Statistical Analysis

Because all runoff plots on the same grassland were subjected to the same rainfall events, the reduction in runoff or sediment yield or the contribution to runoff or sediment yield reduction could be compared among different treatments within a grassland. We calculated runoff and sediment reduction amounts and the corresponding contributions of each near-soil-surface factor under each effective rainfall event according to the above formulae, and used the results as the basic data for the following analysis. Spearman correlation analysis was used to evaluate the relationships between rainfall parameters and the contributions of near-soil-surface factors to runoff and sediment reduction. A curve estimation model was used to identify the relationship between the amount of runoff or sediment reduction and I₃₀. Additionally, Kruskal-Wallis analysis was used to test for differences in the contribution to runoff or sediment yield reduction among all of the near-soil-surface factors. All analyses were performed using SPSS v19.0 software, and graphs were constructed with Origin 2018 software.

3.1 Runoff and Sediment Yield

The runoff and sediment yield varied greatly among treatments, with near-soil-surface factors successively superimposed (corresponding to T1-T4) (Table S3 and S4). Specifically, the runoff yield ranged from 0.0 to 5.3 mm in the BI grassland and from 0.0 to 7.0 mm in the AG grassland. Compared with that

in bare land, runoff was reduced by 39.7%, 43.9%, 75.1%, and 93.4% in T1, T2, T3, and T4 treatments, respectively, in the BI grassland, and the corresponding values in the AG grassland were 11.4%, 7.3%, 47.4%, and 72.2%, respectively. The ranges of sediment yield in the BI and AG grasslands were 0.0–1138.8 and 0.0–476.8 g/m², respectively, with the average values of 53.0 and 32.1 g/m², respectively. The different treatments exerted different effects on sediment yield. For example, T1, T2, T3, and T4 reduced sediment yield relative to that of bare land by 65.1%, 92.4%, 98.7%, and 99.7%, respectively, in the BI grassland and by 57.2%, 76.4%, 93.6%, and 96.9%, respectively, in the AG grassland.

3.2 Reduction of Vegetation Near-Soil-Surface Factors on Runoff and Sediment Yield

Overall, the effects on runoff and sediment reduction differed among the vegetation near-soil-surface factors—i.e., plant canopy, litter, plant roots, and BSCs—in the two types of grasslands (Fig. 2). The plant canopy, litter, and plant roots contributed to runoff reduction in both BI and AG grasslands, with litter being the greatest contributor (Fig. 2a and 2b). The amount of runoff reduction achieved by plant canopy, litter, and plant roots was 0.3, 0.6, and 0.5 mm, respectively, in the BI grassland and 0.4, 0.7, and 0.2 mm, respectively, in the AG grassland. BSCs showed an opposite effect compared with other factors, contributing to runoff yield. Similarly, all near-soil-surface factors reduced sediment yield. In the BI grassland, plant canopy, litter, BSCs, and plant roots reduced the sediment yield by 0.0–649.9 g/m² (Fig. 2d). Among the factors, plant roots achieved the highest average sediment reduction (108.0 g/m²), which was 56.0, 7.9, and 2.1 times that of plant canopy, litter, and BSCs, respectively. In the AG grassland, the near-soil-surface factors reduced sediment yield by 0.0–282.5 g/m² (Fig. 2d). The plant roots produced the greatest reduction in sediment yield (53.8 g/m²), which were 17.0, 3.3, and 2.9 times those of plant canopy, litter, and BSCs, respectively.

Fig. 2 Reduction of runoff amount and sediment yield of different near-soil-surface factors at each rainfall event in the BI grassland (a, c) and AG grassland (b, d). F1, F2, F3, and F4 refer to plant canopy, litter, BSCs, and plant roots, respectively.

3.3 Contributions of Vegetation Near-Soil-Surface Factors to Reductions of Runoff and Sediment Yield

In general, the total contribution of all near-soil-surface factors to reducing runoff in the BI grassland was 70.8% (ranging from 19.5% to 94.7%), which was 1.3 times greater than that in the AG grassland (53.1%; ranging from -19.4% to 79.4%). In addition, the factors in both BI and AG grasslands made significant contributions to controlling sediment loss, which was reduced by 97.4% (ranging from 92.2% to 99.8%) in the BI grassland and 96.7% (ranging from 92.9% to 99.1%) in the AG grassland.

In the BI grassland, the contribution of plant canopy to runoff reduction was 48.8%, which was 3.6 and 3.3 times those of litter (13.5%) and plant roots (14.8%), respectively (Table 1). In the AG grassland, the contribution of plant canopy was 39.9%, which were 1.7 and 7.5 times those of litter (23.9%) and plant roots (5.3%) (Table 1). For BSCs, their contributions to reducing runoff were negative, being -6.4% (algae) in the BI grassland and -16.1% (moss) in the AG grassland, respectively. The contributions of factors (except BSCs) to soil sediment reduction were similar, with percentages of approximately 25% (Table 1). Plant canopy, litter, BSCs, and plant roots made contributions of 21.3%, 24.3%, 24.2%, and 27.7%, respectively, in the BI grassland and 29.5%, 29.9%, 10.3%, and 27.0%, respectively, in the AG grassland. However, the contribution of BSCs to sediment yield reduction in the AG grassland was only 10.3%, which was lower than the contributions of other factors.

Table 1 Average contribution of each near-soil-surface factor to runoff and sediment yield reduction

Near-soil-surface factor	Runoff		Sediment yield	
	BI grassland	AG grassland	BI grassland	AG grassland
Plant canopy (%)	48.8	39.9	21.3	29.5
Plant litter (%)	13.5	23.9	24.3	29.9
BSCs (%)	-6.4	-16.1	24.2	10.3
Plant roots (%)	14.8	5.3	27.7	27.0

Note: Different lowercase letters within a column indicate significant differences at $P<0.05$ level. BI grassland is the grassland with *B. ischaemum* as the dominant species, and AG grassland is the grassland with *A. gmelinii* as the dominant species. BSCs, biological soil crusts.

4.1 Effects of Vegetation Near-Soil-Surface Factors on Reduction of Runoff and Sediment Yield

With the gradual improvement of vertical structure of the vegetation community in the grassland, the reductions in runoff and sediment yield gradually strengthened. Wen et al. [?] found that a complex stratified structure was more effective than a single vegetation layer in controlling soil erosion. For example, in contrast to single factors, multiple near-soil-surface factors reduced runoff generation by 51.0% and sediment yield by 72.3% due to their strong ability to absorb raindrop energy through multiple interceptions (for example, plant canopy and litter) [?]. Our results showed that runoff and sediment in both BI and AG grasslands were reduced by 7.3% to 93.4% compared with that in bare land (Tables S3 and S4), which represented simple near-soil-surface factors. Furthermore, we confirmed that different near-soil-surface factors exerted different effects on runoff and sediment yield.

In the BI and AG grasslands, plant litter was more effective in reducing runoff and sediment yield than plant canopy, plant roots, and BSCs. This result can be attributed to various water-holding capacity abilities of different factors. Generally, plant litter reduces overland flow by covering the ground and absorbing water [?]. For example, water-holding capacity of plant litter in typical plant communities was found in the range from 5.1 to 12.8 g/g in the hilly region of the Loess Plateau [?]. In addition, plant litter can be incorporated into the topsoil and hold water, thereby reducing runoff production [?]. More importantly, plant litter covering the soil surface improves soil surface roughness and reduces flow velocity and energy of overland flow, thereby inhibiting soil loss [?]. Furthermore, plant litter incorporated into the topsoil can improve soil structure, permeability, and stability, thus reducing soil detachment capacity [?]. In a previous study, the incorporation of grass litter into topsoil (with a litter rate of 0.2 kg/m² and a soil bulk density of 1.25 g/cm³, respectively) reduced soil loss compared with that on bare land by 34% and 21% under rainfall intensities of 40 and 80 mm/h, respectively [?]. Moreover, plant canopy was found to reduce runoff production by intercepting raindrops and delay the time of runoff initiation [?], and plant root systems might improve soil structure (e.g., soil porosity) and enhance soil water infiltration capability [?].

However, our study showed that BSCs increased runoff, by an average of 0.0 mm (ranging from -0.5 to 0.2 mm) in the BI grassland and an average of 0.1 mm (ranging from -19.4 to 0.3 mm) in the AG grassland (Fig. 2a and 2b). Although several studies have reported that the growth of BSCs promotes the infiltration capacity of soil by increasing soil porosity and enhancing soil aggregate stability, thereby reducing runoff [?], BSCs can form an impermeable seal by absorbing water during rainfall events, which may inhibit soil water infiltration and accelerate runoff generation [?]. In the present study, BSCs increased runoff but reduced sediment yield. The reasons may be as follows: (1) BSCs increased soil surface roughness, reduced flow velocity, and weakened flow shear stress [?]; (2) rainfall intensity was low, but rainfall duration was long, resulting in low flow velocity and energy; and (3) flow shear stress was not enough to destroy BSCs, and consequently, sediment yield did not increase with increasing runoff [?]. Overall, our results indicate that near-soil-surface factors play essential roles in reducing runoff and sediment and that different factors show diverse effects.

4.2 Contributions of Vegetation Near-Soil-Surface Factors to Runoff and Sediment Yield Reduction

Our results showed that the total contribution of all near-soil-surface factors to reducing runoff in the BI grassland was 70.8% (ranging from 19.5% to 94.7%), which was 1.3 times greater than the corresponding contribution in the AG grassland (53.1%; ranging from -19.4% to 79.4%). This difference can be attributed to the different structural and morphological traits of the two dominant herbaceous plants in the two grasslands. In the BI grassland, the canopy has a tussock-forming architecture and covers the soil surface like a carpet, which

is effective in intercepting rainfall. In the AG grassland, plant canopy features an expanding shoot architecture, and plants are spaced far apart, reducing the chance of intercepting raindrops [?]. Plant litter in the BI grassland is standing litter and is not effective in reducing runoff. However, plant litter in the AG grassland is composed of thick branches and small leaves scattered on the soil surface, which can hinder and accumulate runoff. The fibrous root systems in the BI grassland have numerous fine roots, which increase soil porosity and water infiltration more effectively than the tap root systems in the AG grassland. All these differences resulted in lower runoff reduction in the AG grassland than in the BI grassland.

Because near-soil-surface factors affected runoff in different ways and via different mechanisms, the contributions of plant canopy, litter, BSCs, and plant roots to runoff reduction or increase differed. Among all near-soil-surface factors, plant canopy had the maximum contributions to runoff reduction in both BI and AG grasslands. In the BI grassland, the contribution of plant canopy to runoff reduction was 48.8%, which was 3.6 and 3.3 times those of litter (13.5%) and plant roots (14.8%), respectively (Table 1). In the AG grassland, the contribution of plant canopy was 39.9%, which was 1.7 and 7.5 times those of litter (23.9%) and plant roots (5.3%; Table 1). These differing contributions to runoff reduction were probably related to different spatial distribution of these factors. During rainfall events, plant canopy is the first structure to intercept rainfall, and its effects on runoff are not affected by other factors (e.g., litter and plant roots). Regarding BSCs, their contributions to runoff reduction were negative, i.e., -6.4% (algae) in the BI grassland and -16.1% (moss) in the AG grassland, respectively (Table 1). BSCs increased runoff generation through the formation of an impermeable layer after swelling, and the swelling mechanisms of different components of BSCs are different. As reported by Liu et al. [?], the expansion of algae occurs through the hydration of the sheath, while that of moss occurs through the hydration of stems and leaves. Furthermore, the swelling height of moss was 6 times that of algae, which resulted in moss being more impermeable and increasing runoff generation more than algae.

The BI and AG grasslands both made significant contributions to the control of soil erosion, which was reduced by 97.4% (ranging from 92.2% to 99.8%) in the BI grassland and 96.7% (ranging from 92.9% to 99.1%) in the AG grassland (Table 1). These results indicate that long-term restored grasslands have a strong ability to control soil erosion. Differences in the reductions in sediment yield among near-soil-surface factors were not significant despite the variation in the mechanisms controlling soil erosion, as mentioned above. In other words, the contributions of these factors (except for BSCs) to sediment reduction were similar, with proportions of approximately 25% (Table 1). Plant canopy, litter, BSCs, and plant roots made contributions of 21.3%, 24.3%, 24.2%, and 27.7%, respectively, in the BI grassland and 29.5%, 29.9%, 10.3%, and 27.0%, respectively, in the AG grassland (Table 1). Therefore, near-soil-surface factors all contributed to the control of soil loss. However, the contribution of BSCs to sediment yield reduction in the AG grassland was only 10.3%, which was lower

than the contributions of other factors. As mentioned above, BSCs can reduce soil infiltration and enhance runoff generation; thus, they might be expected to exacerbate soil loss. In the AG grassland, BSCs were composed mainly of moss, which has a strong ability to promote runoff production. Wang et al. [?] reported that BSCs accelerated soil erosion after being fractured when the flow shear stress exceeded its threshold.

4.3 Effects of Rainfall Characteristics on Soil Erosion Mediated by Near-Soil-Surface Factors

Rainfall is an essential factor affecting runoff and sediment yield on slopes, and it can be described by various parameters, such as P, IA, t, I30, E, and R. In general, P, t, and IA are commonly used to reflect the effects of rainfall on soil erosion, and generally, runoff and sediment yield increase with increases in these parameters. In this study, among all 6 parameters representing rainfall characteristics, there was a strong correlation between I30 and each of the other rainfall parameters as well as the reduction in runoff or sediment yield (Fig. 3). In addition, as mentioned by Renard [?], I30 can represent the difference in rainfall intensity and R, and reflect the prolonged peak rates of detachment and runoff during a whole rainfall event. Therefore, I30 was applied in this study to reflect the effects of rainfall characteristics on soil erosion under all rainfall events.

In this study, the reduction in runoff or sediment yield of each near-soil-surface factor (with the exception of BSCs in the BI grassland) increased with I30 (Fig. 4). In the BI grassland, the reductions in runoff and sediment yield caused by plant canopy, litter, and plant roots increased with I30 as a power function ($R^2=0.74$; $*P* < 0.01$; Fig. 4). In the AG grassland, only runoff reductions by plant canopy and litter increased significantly ($P < 0.01$), and the reduction in runoff caused by plant roots and the reduction in sediment yield caused by plant canopy, litter, and plant roots tended to increase with increasing I30 (Fig. 4). In addition, when I30 was greater than 10 mm/h, the runoff and sediment reductions of plant canopy, litter, and plant roots in both BI and AG grasslands increased rapidly with increasing I30. Moreover, when taking intact grassland into consideration, power functions were detected between rainfall characteristics (represented by I30) and total contributions to runoff and sediment yield reductions in the BI and AG grasslands ($P < 0.05$; Fig. 5). As I30 increased, the reduction in runoff or sediment yield increased continuously (Fig. 5). This pattern could have been observed because the ability of all near-soil-surface factors to reduce runoff or sediment yield was not fully reflected in natural rainfall events (with I30 being less than 12 mm/h in this study); moreover, vegetation influenced the reductions in runoff and sediment yield to a greater extent than rainfall conditions. However, when I30 reaches a certain value, the ability of near-soil-surface factors to inhibit soil erosion might reach a limit. Therefore, it is necessary to further explore the rainfall conditions under which the contribution of near-soil-surface factors to the inhibition of soil erosion is the greatest and stable.

Fig. 3 Correlation analysis of rainfall characteristics with runoff and sediment yield under different near-soil-surface factors. P, precipitation; t, rainfall duration; IA, average rainfall intensity; I30, maximum 30-min rainfall intensity; E, rainfall kinetic energy; R, rainfall erosivity. RR_C and SR_C indicate runoff and sediment yield reduction caused by plant coverage; RR_L and SR_L indicate runoff and sediment yield reduction caused by litter; RR_B and SR_B indicate runoff and sediment yield reduction caused by BSCs (biological soil crusts); and RR_R and SR_R indicate runoff and sediment yield reduction caused by plant roots. **, $P<0.01$ level; *, $P^*<0.05$ level.

Fig. 4 Variations in runoff and sediment yield reduction with I30 (maximum 30-min rainfall intensity) under different near-soil-surface factors in the BI grassland and the AG grassland. (a, e), plant canopy; (b, f), litter; (c, g), BSCs (biological soil crusts); (d, h), plant roots.

Fig. 5 Contributions of grassland to runoff reduction (a) and sediment yield reduction (b)

5 Conclusions

This study was performed to investigate the effects of near-soil-surface factors (plant canopy, litter, BSCs, and plant roots) on runoff and sediment yield under natural rainfall events by successive removal of these factors in the field. Our results showed that plant canopy, litter, and plant roots reduced the runoff amount, while BSCs promoted runoff generation. All these near-soil-surface factors reduced the sediment yield. The reductions in runoff and sediment yield caused by plant canopy, litter, and plant roots either tended to increase with I30 or increased with I30 as a power function. Among all near-soil-surface factors, plant canopy had the largest contribution to runoff reduction, which was much greater than those of other factors, in both grassland types, and BSCs contributed the most to runoff production. All near-soil-surface factors contributed equally to reducing sediment yield (by approximately 25%) except for the moss crusts in the grassland dominated by *A. gmelinii* (approximately 10.3%). In general, grass is more effective at reducing soil loss than runoff. Moreover, due to the differences in structural composition and morphological characteristics between the two grassland communities, the grassland with *B. ischaemum* as the dominant species performed better at reducing runoff.

Due to the unique climate and vegetation conditions in arid and semi-arid regions, it is essential and practical to promote natural restoration of grasslands in these regions to prevent soil erosion in the long term. This study enhances our understanding of the mechanisms by which grassland communities conserve soil and water, which can improve the accuracy of many soil erosion models, especially those for arid and semi-arid regions. This study also has reference value for assessing actual soil and water loss situations and guiding the implementation of ecological projects.

Acknowledgements

This research was funded by the Chinese Academy of Sciences (CAS) “Youth Scholar of West China” Program (XAB2019A04) and the National Natural Science Foundation of China (42130717).

References

Biesemans J, Meirvenne M, Gabriels D. 2000. Extending the RUSLE with the Monte Carlo error propagation technique to predict long-term average off-site sediment accumulation. *Journal of Soil & Water Conservation*, 55(1): 35-42.

Chen H, Zhang X P, Abla M, et al. 2018. Effects of vegetation and rainfall types on surface runoff and soil erosion on steep slopes on the Loess Plateau, China. *CATENA*, 170: 141-149.

Coppola A, Basile A, Wang X, et al. 2011. Hydrological behaviour of microbiotic crusts on sand dunes: Example from NW China comparing infiltration in crusted and crust-removed soil. *Soil & Tillage Research*, 117: 34-43.

de Baets S, Poesen J, Knapen A, et al. 2007. Impact of root architecture on the erosion-reducing potential of roots during concentrated flow. *Earth Surface Processes and Landforms*, 32(9): 1323-1345.

de Baets S, Poesen J, Meersmans J, et al. 2011. Cover crops and their erosion-reducing effects during concentrated flow erosion. *CATENA*, 85(3): 237-244.

Du H D, Jiao J Y, Jia Y F, et al. 2013. Phytogenic mounds of four typical shoot architecture species at different slope gradients on the Loess Plateau of China. *Geomorphology*, 193: 57-64.

Facelli J M, Pickett S T. 1991. Plant litter-its dynamics and effects on plant community structure. *Botanical Review*, 57: 1-32.

Feng Q, Zhao W W. 2014. The study on cover-management factor in USLE and RUSLE: a review. *Acta Ecologica Sinica*, 34(16): 4461-4472. (in Chinese)

Gyssels G, Poesen J, Bochet E, et al. 2005. Impact of plant roots on the resistance of soils to erosion by water: a review. *Progress in Physical Geography-Earth and Environment*, 29(2): 189-217.

Kheirfam H, Roohi M. 2020. Accelerating the formation of biological soil crusts in the newly dried-up lakebeds using the inoculation-based technique. *Science of the Total Environment*, 706: 136036, doi: 10.1016/j.scitotenv.2019.136036.

Li C J, Pan C Z. 2018. The relative importance of different grass components in controlling runoff and erosion on a hillslope under simulated rainfall. *Journal of Hydrology*, 558: 90-103.

Lin Y M, Cui P, Ge Y G, et al. 2014. The succession characteristics of soil erosion during different vegetation succession stages in dry-hot river valley of Jinsha River, upper reaches of Yangtze River. *Ecological Engineering*, 62: 13-26.

Liu F, Zhang G H, Sun L, et al. 2016. Effects of biological soil crusts on soil detachment process by overland flow in the Loess Plateau of China. *Earth Surface Processes and Landforms*, 41(7): 875-883.

Liu J X, Liu G B, Flanagan D C, et al. 2020. Effects of soil-incorporated plant litter morphological characteristics on the soil detachment process in grassland on the Loess Plateau of China. *Science of the Total Environment*, 705: 134651, doi: 10.1016/j.scitotenv.2019.134651.

Renard K G. 1997. *Predicting soil erosion by water: a guide to conservation planning with the Revised Universal Soil Loss Equation (RUSLE)*. Washington: USDA, Agricultural Research Service, 703: 404.

Sun C L, Chai Z Z, Liu G B, et al. 2017. Changes in species diversity patterns and spatial heterogeneity during the secondary succession of grassland vegetation on the Loess Plateau, China. *Frontiers in Plant Science*, 8: 1465.

Sun L, Zhang G H, Liu F, et al. 2016. Effects of incorporated plant litter on soil resistance to flowing water erosion in the Loess Plateau of China. *Biosystems Engineering*, 147: 238-247.

Wang B, Zhang G H, Zhang X C, et al. 2014. Effects of near soil surface characteristics on soil detachment by overland flow in a natural succession grassland. *Soil Science Society of America Journal*, 78(2): 589-597.

Wang B, Zhang G H, Shi Y Y, et al. 2015. Effects of near soil surface characteristics on the soil detachment process in a chronological series of vegetation restoration. *Soil Science Society of America Journal*, 79(4): 1213-1222.

Wang B, Zhang G H. 2017. Quantifying the binding and bonding effects of plant roots on soil detachment by overland flow in 10 typical grasslands on the Loess Plateau. *Soil Science Society of America Journal*, 81(6): 1567-1576.

Wang B, Zhang G H, Yang Y F, et al. 2018. Response of soil detachment capacity to plant root and soil properties in typical grasslands on the Loess Plateau. *Agriculture Ecosystems & Environment*, 266: 68-75.

Wang J H, Li Z B, Yao W Y, et al. 2017. Influence of vegetation on runoff and sediment in wind-water erosion crisscross region in the upper Yellow River of China. *Chinese Geographical Science*, 27: 569-576.

Wang L J, Zhang G H, Zhu P Z, et al. 2020. Comparison of the effects of litter covering and incorporation on infiltration and soil erosion under simulated rainfall. *Hydrological Processes*, 34(13): 2911-2922.

Wen Z, Lees B G, Jiao F, et al. 2010. Stratified vegetation cover index: A new way to assess vegetation impact on soil erosion. *CATENA*, 83(1): 87-93.

Zhang B J, Zhang G H, Yang H Y, et al. 2019. Soil resistance to flowing water erosion of seven typical plant communities on steep gully slopes on the Loess Plateau of China. *CATENA*, 173: 375-383.

Zhang S T, Zhang J Z, Liu Y, et al. 2018. Effects of farmland vegetation row direction on overland flow hydraulic characteristics. *Hydrology Research*, 49(6): 1991-2001.

Zhao Y G, Xu M X. 2013. Runoff and soil loss from revegetated grasslands in the hilly Loess Plateau region, China: Influence of biocrust patches and plant canopies. *Journal of Hydrologic Engineering*, 18(4): 387-393.

Zheng H, Chen F, Ouyang Z Y, et al. 2008. Impacts of reforestation approaches on runoff control in the hilly red soil region of Southern China. *Journal of Hydrology*, 356(1-2): 174-184.

Zhou Z C, Shangguan Z P, Zhao D. 2006. Modeling vegetation coverage and soil erosion in the Loess Plateau Area of China. *Ecological Modelling*, 198(1-2): 263-268.

Appendix

Table S1 Basic soil information of sample plots

Sample plot	Soil layer	Soil organic carbon (g/kg)	Total nitrogen (g/kg)	Available phosphorus (mg/kg)	Available potassium (mg/kg)	Bulk density (g/cm ³)
BI grass-land	0-10 cm					
	10-20 cm					
AG grass-land	0-10 cm					
	10-20 cm					
Sample plot	Soil layer	>5.00 mm	2.00-5.00 mm	1.00-2.00 mm	0.50-1.00 mm	0.25-0.50 mm <0.25 mm
BI grass-land	0-10 cm					
	10-20 cm					

Sample plot	Soil layer	>5.00 mm	2.00-5.00 mm	1.00-2.00 mm	0.50-1.00 mm	0.25-0.50 mm	<0.25 mm
AG grass-land	0-10 cm						
	10-20 cm						

Note: BI grassland is the grassland with *B. ischaemum* as the dominant species, and AG grassland is the grassland with *A. gmelinii* as the dominant species.

Table S2 Characteristics of rainfall events

Sample site	Rainfall date	Rainfall amount (mm)	Rainfall duration (min)	IA (mm/h)	I30 (mm/h)	Rainfall kinetic energy (MJ/hm ²)	Rainfall erosivity (MJ · mm/(hm ²))
BI grass-land							
AG grass-land							

Notes: I30, maximum 30-min rainfall intensity. BI grassland is the grassland with *B. ischaemum* as the dominant species, and AG grassland is the grassland with *A. gmelinii* as the dominant species.

Table S3 Variations in runoff and sediment yield in the BI grassland under rainfall events

Treatment	Runoff (mm)	Sediment (g/m ²)	Runoff (mm)	Sediment (g/m ²)	Runoff (mm)	Sediment (g/m ²)
T0						
T1						
T2						
T3						
T4						

Notes: T0, bare land; T1, plant roots; T2, plant roots + BSCs; T3, plant roots

+ BSCs + litter; T4, intact grassland. BSCs, biological soil crusts. BI grassland refers to the grassland with *B. ischaemum* as the dominant species.

Table S4 Variations in runoff and sediment yield in the AG grassland under rainfall events

Treatment	Sediment		Sediment	
	Runoff (mm)	(g/m ²)	Runoff (mm)	(g/m ²)
T0				
T1				
T2				
T3				
T4				

Note: T0, bare land; T1, plant roots; T2, plant roots + BSCs; T3, plant roots + BSCs + litter; T4, intact grassland. BSCs, biological soil crusts. AG grassland refers to the grassland with *A. gmelinii* as the dominant species.

Fig. S1 Schematic diagram of vegetation near-soil-surface factors involved in different treatments (T1-T4). BSCs, biological soil crusts.

Fig. S2 Recorded rainfall events in the BI and AG grasslands during rainy season. BI grassland is the grassland with *B. ischaemum* as the dominant species, and AG grassland is the grassland with *A. gmelinii* as the dominant species. Non-effective rainfall events indicate that no runoff or sediment is generated. Effective rainfall events indicate that runoff and sediment are generated.

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv –Machine translation. Verify with original.