

Effects of Nitrogen Addition on Nitrogen Transformation and Related Functional Gene Abundance in *Pinus sylvestris* var. *mongolica* Plantations (Postprint)

Authors: LIU Xinjun, Wei Hongjie

Date: 2022-04-29T22:11:46Z

Abstract

Soil nitrogen (N) availability is an important factor affecting soil microbial community structure and soil nitrogen cycling; however, the effects of N addition on soil N transformation and N functional genes (NFGs) expression in Mongolian pine plantations remain poorly understood. To explore the effects of N addition on nitrogen transformation in Mongolian pine plantations and its underlying mechanisms, this study conducted a 2-year N addition experiment in a Mongolian pine plantation at Saihanba Qiancengban Forest Farm, with four N addition levels of 0, 1, 5, and 10 gN m⁻² · year⁻¹, designated as N0, N1, N5, and N10, respectively. Using the functional gene microarray GeoChip 5.0 system and laboratory soil incubation methods, the responses of soil NFGs to N addition and their effects on nitrogen transformation processes were investigated. The results showed that: (1) Compared with N0, medium and low N addition treatments (N1, N5) promoted the relative abundances of genes related to ammonification (*ureC*, *nirA*, *nrfA*), nitrification (*amoA*), and denitrification (*norB*), while high N treatment (N10) inhibited the expression of all NFGs. (2) Correlation analysis indicated that the promoting effects of N1 and N5 were significantly correlated with soil organic carbon (SOC), nitrate nitrogen (NO₃-N), and microbial biomass carbon (MBC), while N10 treatment significantly reduced the relative abundances of NFGs in all nitrogen transformation processes, and this negative effect was associated with decreased dissolved organic carbon (DOC) and MBC contents. (3) Similar to the trend observed in nitrogen transformation gene abundances, N1 and N5 treatments significantly increased net N nitrification, net N mineralization, and N₂O emission rates, whereas the promoting effect of N10 was not significant, indicating that there exists a threshold for the promoting effect of N addition on nitrogen transformation. (4) Multiple regression analysis further demonstrated that *amoA*-AOB and MBC were key

factors affecting net N nitrification, ureC, nirK, and MBC were key factors influencing net nitrogen mineralization, and narG and nirS were key factors affecting N₂O emissions. In summary, N addition can enhance nitrogen transformation and increase the relative abundances of some specific enzyme functional genes in Mongolian pine plantations; however, there exists a threshold for N addition levels. When applying 10 g N m⁻² · year⁻¹, nitrogen transformation is inhibited, and 5gN m⁻² · year⁻¹ is the optimal level for promoting soil N transformation in Mongolian pine plantations.

Full Text

Title and Authors

Effects of Nitrogen Addition on Nitrogen Transformation and Related Functional Gene Abundance in *Pinus sylvestris* var. *mongolica* Plantations

LIU Xinjun¹, WEI Hongjie² ¹ Department of Ecological Engineering, Hebei Professional College of Political Science and Law, Shijiazhuang 050061, China
² Shijiazhuang Municipal Water System Garden Center, Shijiazhuang 050061, China

Abstract: Soil nitrogen (N) availability is a critical factor influencing soil microbial community structure and N cycling, yet little is known about how N addition affects soil N transformation and N functional gene (NFG) expression in *Pinus sylvestris* var. *mongolica* plantations. To explore these effects and underlying mechanisms, we conducted a two-year N addition experiment in a *Pinus sylvestris* var. *mongolica* plantation at Saihanba Mechanical Forest Farm, applying four N addition levels: 0, 1, 5, and 10 g N m⁻² year⁻¹ (designated N0, N1, N5, and N10). Using the GeoChip 5.0 functional gene microarray system and laboratory soil incubation, we examined soil NFG responses to N addition and their influence on N transformation processes. Results showed: (1) Compared with N0, moderate N addition (N1, N5) promoted the relative abundance of genes related to ammonification (ureC, nirA, nrfA), nitrification (amoA), and denitrification (norB), while high N treatment (N10) inhibited expression of all NFGs. (2) Correlation analysis revealed that the promotional effects of N1 and N5 were significantly correlated with soil organic carbon (SOC), nitrate nitrogen (NO⁻-N), and microbial biomass carbon (MBC), whereas N10 significantly reduced relative abundance of NFGs for all N transformation processes, an effect associated with decreased dissolved organic carbon (DOC) and MBC content. (3) Consistent with NFG abundance patterns, N1 and N5 treatments significantly increased net N nitrification, net N mineralization, and N₂O emission rates, but N10 showed no significant promotion, indicating a threshold effect for N addition on N transformation. (4) Multiple regression analysis further identified amoA-AOB and MBC as key factors affecting net N nitrification, ureC, nirK, and MBC as key factors influencing net N mineralization, and narG, nirS as key factors controlling N₂O emissions. In conclusion, N addition can enhance

N transformation and increase relative abundance of specific enzyme functional genes in *Pinus sylvestris* var. *mongolica* plantations, but a threshold exists. At 10 g N m⁻² year⁻¹, N transformation was inhibited, while 5 g N m⁻² year⁻¹ represented the optimal level for promoting soil N transformation.

Keywords: nitrogen addition, nitrogen functional genes (NFGs), net nitrogen transformation, N O emissions

Introduction

Soil nitrogen (N) availability is a crucial factor affecting plant growth and soil biogeochemistry in terrestrial ecosystems (Ju, 2014). Global N deposition is altering soil N availability and influencing ecosystem N cycling (Liao et al., 2021; Wu et al., 2019), including N mineralization, nitrification, and N O emissions, which may lead to environmental problems such as N supply-demand imbalance, non-point source pollution, and greenhouse gas emissions (Li et al., 2021). Soil microorganisms are considered the primary drivers of N cycling processes, yet the response of core microbial processes in N cycling to N deposition and their relationship with N transformation remain poorly understood.

Nitrogen functional genes (NFGs) directly encode key enzymes that influence N cycling processes and serve as direct indicators of N biotransformation function (Liao et al., 2019). Previous NFG research has focused primarily on agricultural ecosystems and subtropical forest systems. In agricultural ecosystems, nitrogen fertilization has shown no significant effect on N fixation gene (*nifH*) abundance but increased relative abundance of nitrification genes (*amoA*) and denitrification genes (*nirK*, *nirS*, *nosZ*), primarily regulated by N form and soil pH (Ouyang et al., 2018). In subtropical forest soils, N addition decreased abundance of N fixation genes (*nifH*) and nitrification genes (*amoA*) while increasing denitrification gene abundance (Tian et al., 2019). Compared with subtropical forests and agricultural soils, cold-temperate forest soils in northern regions are characterized by relatively poor nutrient status, low N input, low precipitation, and large temperature fluctuations (Tian et al., 2019), which may result in different microbial community structures and differential effects on N transformation. However, few studies have reported on the effects of N addition on NFGs in cold-temperate forest soils.

In recent decades, scholars have conducted extensive research on N transformation in forest ecosystems, revealing preliminary mechanisms by which abiotic and biotic factors drive specific N transformation processes (i.e., nitrification, mineralization, and N O emissions) under N addition (Chelsea et al., 2016). Among all NFGs, the *amoA* gene affecting the rate-limiting step of nitrification, the *narG* gene affecting the first step of denitrification, and *nir* genes affecting rate-limiting steps are considered hub factors controlling N transformation (Ouyang et al., 2018; Chelsea et al., 2016). Soil available phosphorus, pH, and dissolved N have been identified as primary factors influencing NFG abundance (Wang et al., 2017), indicating that soil properties and microbial function play

important roles in N transformation. Exogenous N addition can directly affect soil properties and microbial community structure, exerting profound influences on N transformation processes (Wang et al., 2017). However, the extent to which these factors affect N transformation processes and the dominant factors influencing N transformation remain unclear. Therefore, critical questions remain: Can N addition mediate soil NFG expression to affect related N transformation processes? What are the relationships among soil N transformation processes, NFGs, and soil properties? These questions severely constrain our understanding of soil N transformation and its influencing mechanisms in *Pinus sylvestris* var. *mongolica* plantations. This study addresses these gaps by examining a *Pinus sylvestris* var. *mongolica* plantation in Saihanba, Hebei Province, using the GeoChip 5.0 functional gene microarray system and laboratory soil incubation. By analyzing effects of different N addition levels on soil properties, NFGs, and N transformation parameters, we explore the relative contributions of NFGs and soil properties to N transformation under different N addition levels and identify key factors affecting related N transformation processes, providing a theoretical basis for N fertilizer management in *Pinus sylvestris* var. *mongolica* plantations.

Materials and Methods

Study Area

The study area is located at the intersection of the Greater Khingan Mountains and the northern Hebei mountainous region, within the QianCengBan Forest Farm of Saihanba Mechanical Forest Farm in Weichang Manchu-Mongolian Autonomous County, northern Hebei Province (17°39'42" E, 42°35'45" N). Established in 1962, the forest farm encompasses 20,029.8 ha of artificial forest, representing a major artificial *Pinus sylvestris* var. *mongolica* population in China, with stand density of 1,300–1,600 trees per hectare. The plantation undergoes harvesting and replanting every 8–10 years. The region features a temperate continental semi-arid and semi-humid climate with long, cold winters and short, dry, windy spring and autumn seasons. The average background N deposition level is 3.34 g N m⁻² year⁻¹. The experimental site has an elevation of 1,432 m, mean annual precipitation of 454.2 mm, mean annual sunshine duration of 2,368.8 h, mean annual evaporation of 1,244.9 mm, mean annual relative humidity of 75.3%, and mean annual temperature of -1.4 °C. The soil type is primarily aeolian sandy soil derived from aeolian, residual, and accumulative parent materials. Understory shrubs and herbs are sparse, dominated by *Thalictrum petaloideum*, *Potentilla longifolia*, *Sanguisorba officinalis*, and *Carex lanceolata*.

Experimental Design

In March 2018, we selected 12-year-old trees and established four N treatments with four replicates each, totaling 16 plots of 6 m × 6 m, separated by 4 m buffer zones. Nitrogen was applied as ammonium nitrate (NH₄NO₃) solution at four levels: 0 (N0), 1 (N1), 5 (N5), and 10 (N10) g N m⁻² year⁻¹. Compared with the

natural N deposition rate ($3.34 \text{ g N m}^{-2} \text{ year}^{-1}$), N1, N5, and N10 represented low, moderate, and high N deposition levels, respectively. For each application, the corresponding NH_4NO_3 was dissolved in 10 L distilled water and applied every two months from June 2018 to June 2020, totaling 12 applications. Applications occurred at approximately 10:30 AM at the beginning of each month, ensuring no rainfall for 15 hours before and after application. The N0 treatment received equal volumes of distilled water with identical timing and method.

Soil Sampling and Processing

On June 20, 2020 (a sunny day), surface soil samples (0–20 cm) were collected using a stainless steel soil auger (20 cm length, 6 cm diameter) after removing surface litter. Five random sampling points were composited into one sample per plot and stored in sterile polyethylene bags. Samples were immediately placed in foam boxes with dry ice and transported to the laboratory. After removing residual roots, litter, and gravel, each sample was divided into three portions: one stored at 4°C for microbial biomass and dissolved nutrient measurements and incubation experiments; another air-dried for basic physicochemical analysis; and the final portion immediately stored at -80°C for NFG relative abundance analysis using the GeoChip 5.0 functional gene microarray system.

Soil Physicochemical Analysis

Soil physicochemical properties were determined following Bao (2000). Soil pH was measured by weighing 10.00 g air-dried soil into a 50 mL beaker, adding distilled water (water:soil = 2.5:1), intermittently stirring for 30 min, settling for 15 min, and measuring with an Ohaus ST3100/F pH meter. Soil organic carbon (SOC) was determined by weighing 0.5000 g air-dried soil (passed through 0.149 mm sieve) into a 150 mL Erlenmeyer flask, adding 0.10 g powdered AgNO_3 , then precisely adding 5 mL each of $0.8 \text{ mol} \cdot \text{L}^{-1}$ potassium dichromate ($\text{K}_2\text{Cr}_2\text{O}_7$) solution and concentrated sulfuric acid, covering with a small funnel, heating in a 230°C electric sand bath for 15 min, and titrating with $0.1 \text{ mol} \cdot \text{L}^{-1}$ ferrous sulfate solution. Total nitrogen (TN) was measured by weighing 0.5000 g air-dried soil into a Kjeldahl flask, moistening with distilled water, adding 8 mL concentrated sulfuric acid, digesting in a fume hood until light blue, and analyzing with an automatic Kjeldahl nitrogen analyzer (KDN-520, Hangzhou Lvbo Instrument Co., Ltd.). Total phosphorus (TP) was determined by weighing 0.25 g air-dried soil into a nickel crucible, adding 3 drops of anhydrous ethanol, covering with 2 g NaOH, heating in a muffle furnace at 400°C for 15 min, washing with 75°C distilled water, transferring to a 100 mL volumetric flask, taking 5 mL digested mother solution to a 50 mL volumetric flask, diluting to 30 mL with distilled water, adding 3 drops of dinitrophenol and 5 mL molybdenum-antimony anti-color reagent, and measuring at 700 nm with a UV spectrophotometer (T-6M, Shanghai Feile Instrument Co., Ltd.). Available phosphorus (AP) was measured by weighing 0.25 g air-dried soil into a 200 mL capped plastic bottle, extracting with 50 mL of $0.5 \text{ mol} \cdot \text{L}^{-1}$ NaHCO_3 at con-

stant temperature (25 °C) for 30 min, filtering with phosphorus-free filter paper, adding 5 mL ammonium molybdate solution and 1 drop of stannous chloride reagent to 5 mL filtrate, developing color for 15 min, and measuring at 680 nm with a UV spectrophotometer. Ammonium nitrogen ($\text{NH}_4\text{-N}$) and nitrate nitrogen ($\text{NO}_3\text{-N}$) were extracted by weighing 5.00 g air-dried soil into a centrifuge tube, adding 25 mL of $2 \text{ mol} \cdot \text{L}^{-1}$ KCl, shaking ($120 \text{ r} \cdot \text{min}^{-1}$) for 2 h, centrifuging at $8,000 \text{ r} \cdot \text{min}^{-1}$ for 15 min, filtering, and analyzing two aliquots of filtrate—one with HCl-EDTA buffer for $\text{NO}_3\text{-N}$ and another with phenol-EDTA buffer for $\text{NH}_4\text{-N}$ —using a continuous flow analyzer (SmartChem 200, AMS/Alliance, Italy). Dissolved organic carbon (DOC) was determined by weighing 20.0 g air-dried soil into a 150 mL capped plastic bottle, adding 60 mL of $5 \text{ g} \cdot \text{L}^{-1}$ sodium metaphosphate, reciprocally shaking at $120 \text{ r} \cdot \text{min}^{-1}$ for 18 h, drying the suspension at 60 °C for 48 h, and analyzing with a TOC auto-analyzer (TOC-LCPH, Shimadzu, Japan). Total dissolved nitrogen (TDN) was measured by weighing 5.00 g air-dried soil into a 50 mL capped plastic bottle, adding 25 mL of $1 \text{ mol} \cdot \text{L}^{-1}$ potassium chloride solution, reciprocally shaking at $180 \text{ r} \cdot \text{min}^{-1}$ for 1 h, settling for 10 min, filtering into a 50 mL volumetric flask, oxidizing 8 mL filtrate with 10 mL potassium persulfate oxidant for 15 min, and analyzing with a continuous flow auto-analyzer. Dissolved organic nitrogen (DON) was calculated as TDN minus ($\text{NH}_4\text{-N} + \text{NO}_3\text{-N}$) (Ji et al., 2020). Microbial biomass carbon (MBC) and nitrogen (MBN) were determined by fumigation-extraction: 70.0 g fresh soil at 40% field capacity was placed in an 80 mL beaker inside a vacuum desiccator with a beaker of chloroform, evacuated to -0.07 MPa until chloroform boiled vigorously for 5 min, repeated until complete fumigation, then transferred to a plastic bottle, extracted with 100 mL of $0.5 \text{ mol} \cdot \text{L}^{-1}$ potassium sulfate solution by shaking (25 °C, $300 \text{ r} \cdot \text{min}^{-1}$) for 30 min, filtered, and analyzed with a TOC auto-analyzer—one 10 mL aliquot with 10 mL sodium hexametaphosphate solution for MBC, and another digested with 5 mL concentrated sulfuric acid for 3 h for MBN.

Measurement of Net N Transformation and N O Emission Rates

Net N transformation was divided into net nitrification rate (R_n) and net mineralization rate (R_m), both determined through 14-day soil incubation. Briefly, 40.00 g soil was placed in a 500 mL polyethylene bottle (8 cm height, 6 cm diameter) at 60% water-holding capacity, pre-incubated in a 25 °C constant-temperature dark incubator for 6 days, then formally incubated for 14 days under the same conditions. Soil samples were taken on days 1, 7, and 14 to determine inorganic N concentrations. R_n and R_m ($\text{mg N kg}^{-1} \text{ d}^{-1}$) were calculated following Wang (2021): $R_n = (\text{post-incubation } \text{NO}_3\text{-N} - \text{pre-incubation } \text{NO}_3\text{-N}) / \text{incubation time}$; $R_m = (\text{post-incubation inorganic N } (\text{NH}_4\text{-N} + \text{NO}_3\text{-N}) - \text{pre-incubation inorganic N}) / \text{incubation time}$.

N O emission rates were measured in parallel incubation experiments. All polyethylene bottles were covered with sterile breathable organic film to prevent moisture loss while maintaining aeration. N O emission rates were measured on

days 1, 3, 5, 7, and 14 using a Picarro G2508 gas concentration analyzer (Picarro G2208 Environment, Picarro Inc., CA, USA) (Ma et al., 2015). Bottles were sealed for 12 h before measurement, and N₂O emission rates ($\text{g kg}^{-1} \text{d}^{-1}$) were calculated from concentration changes during the 12 h sealed period.

Soil Microbial NFG Analysis

The GeoChip 5.0 functional microarray system was used for comprehensive analysis of microbial NFG sequences. This high-throughput platform contains 57,000 oligonucleotide probes covering over 144,000 gene sequences in 373 gene families, detecting broader gene types than qPCR and enabling analysis of diverse microbial functions and biogeochemical processes (Lu et al., 2018).

For each treatment, 0.50 g soil was thawed and DNA extracted using a PowerSoil kit (MoBio, Carlsbad, USA) with three parallel samples per site. DNA purity and quantity were assessed with a spectrophotometer and microplate reader system (FLUOstar OPTIMA, BMG Labtech, Jena, Germany). Qualified DNA samples were fluorescently labeled, hybridized to chips, and scanned (Li et al., 2017), with three replicates per microarray experiment.

Detailed procedures: DNA was labeled with Cy-3 fluorescent dye using random primers, purified with a QIA purification kit (QIAGEN QUICK Purification Kit, Roche NimbleGen Inc, USA), and dried in a rotary evaporator (Savant SVC200, Thermo Savant, Holbrook, NY, USA) at 45 °C for 45 min. Labeled DNA was dissolved in 120 μL hybridization buffer (40% formamide, 0.1% SDS, 10 μg unlabeled DNA, 2 \times SSC), denatured at 90 °C for 5 min, held at 50 °C for 30 min, and hybridized on a MAUI platform at 40 °C for 16 h. Hybridized chips were scanned at 633 nm using a NimbleGen MS200 microarray scanner (Roche NimbleGen, Inc., Madison, WI, USA). Images were converted, extracted, and normalized using Imogene 6.0 software. Normalization included inter-array normalization and false-positive removal, adjusting total signal intensity to the highest sample level in each batch, removing low-quality spots with signal-to-noise ratio < 2.0, normalizing signal values, and converting to natural logarithms (Zhang et al., 2017). All NFG analyses were performed by Shanghai Personal Biotechnology Co., Ltd.

Data Analysis

One-way ANOVA and Tukey's multiple comparison tests (IBM SPSS 22.0) evaluated N addition effects on soil properties, gene relative abundance, and N transformation rates. Pearson correlation analysis examined relationships among soil properties, NFG abundance, and N transformation rates. Figures were prepared using Origin 8.5. All results are presented as means \pm standard deviation (Mean \pm SD) of four replicates.

Results

Effects of N Addition on Soil Physicochemical Properties

Two years of N addition significantly altered several soil physicochemical properties (Fig. 1). Soil NO_3^- -N and DON concentrations increased with N addition level, reaching maximum values under N10 that were significantly greater than N0 ($P < 0.05$). Soil NH_4^+ -N content showed a decreasing trend with increasing N addition, though differences among treatments were not significant. SOC, MBC, and C/N ratio exhibited initial increases followed by decreases, with maximum values under N5. DOC content also decreased with increasing N addition, with the highest value under N1. Other soil properties (MBN, TN, TP, AP, pH) showed minimal variation among treatments with no significant differences.

Effects of N Addition on NFG Relative Abundance

N addition significantly altered the relative abundance of several NFGs (Fig. 2). Among individual genes, N fixation gene (*nifH*) and denitrification gene (*narG*) showed the highest abundance values, accounting for 16.69–18.03% and 17.42–18.98% of total abundance, respectively. Compared with N0, N1 significantly increased relative abundance of *amoA*-AOA and *amoA*-AOB genes (encoding ammonia monooxygenase) involved in nitrification, *norB* gene (encoding nitrite reductase) involved in denitrification, *nirA* gene (encoding nitrate reductase) involved in assimilatory N reduction, and *nrfA* gene (encoding nitrate reductase coenzyme II) involved in dissimilatory N reduction. N5 significantly increased relative abundance of *ureC* gene (encoding urease) involved in ammonification, *amoA*-AOB, *norB*, *nirA*, and *nrfA*. In N10 treatment, all NFGs except *hzsa* gene (encoding glutamate synthase) involved in ammonia oxidation showed significantly decreased relative abundance.

Effects of N Addition on Total NFG Abundance by N Transformation Process

Total relative abundance of NFGs for each N transformation process revealed that denitrification, N fixation, and ammonification had the greatest microbial influence, collectively accounting for over 73.7% of total NFG abundance (Fig. 3). Across the seven N transformation processes (denitrification, N fixation, ammonification, anammox, nitrification, assimilatory N reduction, and dissimilatory N reduction), all processes except anammox showed significantly lower abundance under N10 compared with other treatments, following the trend $\text{N5} > \text{N1} > \text{N0} > \text{N10}$, with no significant differences between N0 and N1.

Effects of N Addition on Net N Nitrification, Net N Mineralization, and N O Emission Rates

Two years of N addition significantly increased net N nitrification, net N mineralization, and N O emission rates under N1 and N5 compared with N0, reaching

maximum values under N5 and declining under N10, which showed no significant difference from N0 (Table 1). Compared with N0, N addition increased net N nitrification, mineralization, and N O emission rates by 18.62–57.41%, 10.95–49.88%, and 1.69–28.55%, respectively.

Correlation Analysis Between NFG Abundance and Soil Factors

Correlation analysis among 12 soil physicochemical indicators and NFG abundance for seven N transformation processes showed that TN, TP, NO⁻N, and DOC were significantly correlated with NFG abundance for certain processes, while other soil factors showed no clear relationships (Table 2). TN was significantly positively correlated with nitrification, assimilatory N reduction, and dissimilatory N reduction processes. TP showed significant positive correlation with dissimilatory N reduction NFG abundance. DOC was significantly positively correlated with N fixation, ammonification, denitrification, assimilatory N reduction, and dissimilatory N reduction, while NO⁻N was significantly positively correlated with NFG abundance for all processes except anammox.

Correlation Analysis of Net N Transformation Rates with NFGs and Soil Properties

Net nitrification rate (R_n), net mineralization rate (R_m), and N O emission rate (R_e) were closely correlated with relative abundance of specific NFGs and soil properties (Table 3). R_n showed significant positive correlations with ureC, amoA-AOB, nirA, nrfA, SOC, NO⁻N, and MBC, with particularly strong correlations with MBC and SOC. R_m was significantly positively correlated with NFGs involved in N cycling (ureC, gdh, nifH, amoA-AOB, narG, nirK, norB, nirA, nrfA) and SOC and MBC, with stronger correlations with MBC and nrfA. R_e showed highly significant correlations with all nitrification and denitrification genes except amoA-AOA and hao, and highly significant positive correlations with SOC and MBC, particularly with amoA-AOB and narG.

Regression Analysis of Net N Transformation and N O Emission Rates

Stepwise multiple regression analysis identified amoA-AOB relative abundance and soil MBC content as dominant factors affecting net N nitrification rate (R_n), with $R^2 = 0.64$ ($P < 0.001$) (Table 4). UreC and nirK relative abundance and soil MBC content were dominant factors affecting net N mineralization rate (R_m), with $R^2 = 0.75$ ($P < 0.001$). NarG and nirS relative abundance were dominant factors affecting N O emission rate (R_e), with $R^2 = 0.69$ ($P < 0.001$).

Discussion

Appropriate N addition can alleviate N limitation in ecosystems, providing substrates and energy for microbial growth and basal metabolism, thereby stimulating soil nutrient availability and enhancing microbial functional activity (Li

et al., 2017). Our study site in Saihanba has been confirmed to have low N reserves (Ren, 2012), suggesting N addition may influence substrate availability. Different N levels had differential effects: SOC, MBC, and DOC showed initial increases followed by decreases with increasing N addition, suggesting that excessive N addition may cause soil N saturation, affecting nutrient allocation and indicating a threshold for N addition effects. During the two-year experiment, soil pH ranged from 7.62 to 7.76 with no significant differences among treatments and no significant correlation with NFG relative abundance (Table 2), suggesting weak pH effects on NFGs in this study. This may be due to high soil buffering capacity (Jiang et al., 2019) or co-effects with other microorganisms such as arbuscular mycorrhizal hyphae (Cao et al., 2021). Additionally, no significant differences were observed among treatments for TN, TP, AP, and MBN, possibly because the two-year N addition period was too short to cause comprehensive changes in soil fertility (Ouyang et al., 2016).

Using the GeoChip 5.0 microarray system, we explored relative abundance of key genes involved in N transformation processes. Moderate N addition (N1, N5) showed promotional effects on all NFGs, particularly those related to ammonification, nitrification, and denitrification, while high N addition (N10) significantly reduced relative abundance of all NFGs.

Among ammonification-related genes, ureC relative abundance significantly increased under N1 and N5, peaking under N5 and declining significantly under N10, while glutamine synthesis gene (gdh) and anammox gene (hzsa) abundance remained unchanged, reflecting a trend toward mineralization as N approached saturation threshold (Zhang et al., 2019). Additionally, nifH showed no significant increase under N1 and N5, consistent with previous findings that N fertilization does not affect nifH relative abundance (Berthrong et al., 2014). This may be because available N (NH_4^+-N) obtained from the environment eliminates the need for additional energy metabolism investment (Zheng et al., 2017).

For nitrification-related genes, N addition (N1, N5) significantly increased relative abundance of amoA-AOA and amoA-AOB, consistent with findings in temperate forests by Szukics (2012), suggesting regional climate is not the primary driver of nitrification. Nitrification gene abundance was significantly positively correlated with soil TN and NO_3^--N (Table 2), indicating that N addition-induced increases in soil N content can enhance nitrification gene abundance. Notably, amoA-AOA abundance was consistently higher than amoA-AOB across all treatments. Previous studies indicate amoA-AOB is mediated by functional microorganisms with ribosomes, showing higher relative abundance in nutrient-rich environments or under moderate N addition, while amoA-AOA has higher affinity for ammonium and better tolerance for N-limited or N-saturated conditions (Ouyang et al., 2019). Therefore, amoA-AOA dominance in abundance but greater sensitivity of amoA-AOB to N addition in our study region mirrors observations in agricultural soils (Song et al., 2020).

For denitrification functional genes, N1 and N5 significantly increased norB relative abundance while other denitrification genes remained largely unchanged,

differing from previous forest system studies showing that *narG*, *nirK*, and *nirS* abundance increased substantially with N addition while *norB* and *nosZ* were insensitive (Tang et al., 2016). This discrepancy may relate to the sequential steps of denitrification: nitrate is the initial substrate, and its concentration significantly affects *narG* expression; as denitrification proceeds, nitrate concentration decreases due to plant uptake or gaseous N loss (Liu et al., 2020), potentially weakening N addition effects on *norB* and *nosZ* (Chen et al., 2012). Additionally, denitrification gene relative abundance was significantly positively correlated with NO_3^- -N and DOC (Table 2), consistent with previous findings that *narG* and *napA* relative abundance positively correlate with SOC and NO_3^- -N content (Wang et al., 2017), likely because denitrifying microorganisms are mostly heterotrophic and thus more dependent on organic matter.

Soil N transformation processes, primarily nitrification, mineralization, and N₂O emission, are important indicators of N supply and loss (Lin et al., 2020). Our study found that moderate and low N addition significantly increased net nitrification rate (*R_n*), net mineralization rate (*R_m*), and N₂O emission rate (*R_e*), but these metrics declined under high N treatment (N10). *R_n* and *R_m* are direct indicators of net changes in soil inorganic N pools, reflecting interactions among organic matter decomposition, mineralization, nitrification, and immobilization (Chen et al., 2012). *R_n* was significantly correlated with core genes of related N transformation processes (Table 3). Stepwise regression analysis identified *amoA*-AOB relative abundance and MBC content as key factors affecting *R_n*, with correlation analysis showing significant relationships between *amoA*-AOB and *R_n*, suggesting *amoA*-AOB may be the primary gene driving soil nitrification in semi-arid plantation ecosystems (Zhang et al., 2019; Tang et al., 2019). Similarly, stepwise regression identified *ureC*, *nirK* abundance, and MBC content as important parameters affecting *R_m*, indicating *R_m* is closely related to organic N mineralization and denitrification processes. *R_e* was significantly correlated with NFGs involved in nitrification and denitrification (Table 3), with denitrification pathway genes *narG* and *nirS* closely related to *R_e*, indicating denitrification is the primary pathway for N₂O release in temperate forest soils. The significant positive correlation between *narG*, *nirS* and *R_e* suggests *nirS* abundance can serve as an effective parameter for estimating *R_e* emissions (Flaa et al., 2019).

Conclusion

This study demonstrates that moderate and low N addition levels (1, 5 g N m⁻² year⁻¹) had no significant effect on total relative abundance of N functional genes (NFGs) but increased relative abundance of specific genes involved in ammonification, nitrification, and denitrification. High N addition treatment (N10) significantly reduced relative abundance of NFGs for all N transformation processes. These effects were primarily associated with changes in soil C and N nutrient indicators (SOC, NO_3^- -N, DON) rather than pH. The promotional effects were significantly correlated with soil SOC, NO_3^- -N, and MBC,

while the negative effects of N10 were associated with reduced DOC and MBC content. Consistent with NFG abundance patterns, N1 and N5 treatments significantly increased net N nitrification, net N mineralization, and N O emission rates, but N10 showed no significant promotion, indicating a threshold effect. Stepwise multiple regression analysis identified amoA-AOB relative abundance and MBC content as key predictors of net N nitrification rate; ureC, nirK relative abundance and MBC content as key predictors of net N mineralization rate; and narG, nirS relative abundance as key predictors of N O emission rate. These results provide a theoretical basis for N fertilizer management in *Pinus sylvestris* var. *mongolica* plantations.

References

- BAO SD, 2000. Soil Agrochemical Analysis[M]. Beijing: China Agriculture Press.
- CAO BF, JIANG HX, LIU L, et al., 2021. Research progress on mechanism of arbuscular common mycorrhizal networks in plant-plant interactions[J]. J Appl Ecol, 32(9):3385-3396.
- CHEN Z, LIU JB, WU MN, et al., 2012. Differentiated response of denitrifying communities to fertilization regime in paddy soil[J]. Microbiol Ecol, 63(2):446-459.
- CHELSEA JC, NICHOLAS CD, MICHAEL BJ, et al., 2016. Meta-analysis reveals ammonia-oxidizing bacteria respond more strongly nitrogen addition ammonia-oxidizing archaea[J]. Soil Biol Biochem, 99:158-166.
- FLAA B, AAMC A, AF A, et al., 2019. Different groups of nitrite-reducers and N O-reducers have distinct ecological niches and functional roles in West African cultivated soils - Science Direct[J]. Soil Biol Biochem, 129:39-47.
- JU XT, 2014. The concept and significance of nitrogenous fertilizer efficiency –concurrently discussing the misunderstanding of traditional nitrogenous fertilizer use efficiency[J]. J Soil, 51(5):921-933.
- JIANG J, CAO NN, YU MX, et al., 2019. A review of the key mechanisms of acidic forest soils to buffer acid deposition[J]. J Trop Subtrop Bot, 27(5):491-499.
- JI HK, WU YY, FU PJ, et al., 2020. Composition and size distribution characteristics of soil dissolved organic nitrogen under different land use types in tropical coastal areas[J]. J Ecol Environ, 29(3):525-535.
- JING H, ZHANG P, LI J, et al., 2019. Effect of nitrogen addition on the decomposition and release of compounds from fine roots with different diameters: the importance of initial substrate chemistry[J]. Plant Soil, 438(1/2):281-296.
- LIN W, LI YZ, LI YJ, et al., 2020. Advances in the mechanism of microbe-driven nitrogen cycling[J]. J Plant Nutr Fert, 26(6):1146-1155.

- LIAO K, SHEN FF, LIU WF, et al., 2020. C, N and P stoichiometric characteristics of litterfall and soil in a Chinese fir plantation under long-term nitrogen deposition[J]. *Ggiihaia*, 40(11):1551-1562.
- LIAO LR, WANG J, ZHANG C, et al., 2019. Effects of grazing exclusion on the abundance of functional genes involved in soil nitrogen cycling and nitrogen storage in semiarid grassland[J]. *J Appl Ecol*, 30(10):3473-3481.
- LIU D, WANG L, CAO ZB, et al., 2020. Effects of mycorrhizal fungi on nitrification and denitrification in the rhizospheric soil of aquatic plants and its microbial mechanism[J]. *Environ Sci*, 41(2):932-940.
- LI HX, XIN YJ, LI ZH, et al., 2021. Effects of different nitrate form on the growth and physiological characteristics for quercus mongolica seedlings[J]. *For Eng*, 37(2):35-40.
- LI H, YANG S, XU ZW, et al., 2017. Responses of soil microbial functional genes to global changes are indirectly influenced by aboveground plant biomass variation[J]. *Soil Biol Biochem*, 104: 18-29.
- LU H, ZHAO Y, SHENG YY, et al., 2018. Research on soil microbial functional genes in alpine meadow based on GeoChip technique[J]. *J Ecol*, 37(10):3031-3037.
- MA F, MA HL, QIU H, et al., 2015. Effects of water levels and the additions of different nitrogen forms on soil net nitrogen transformation rate and N O emission in subtropical forest soils[J]. *J Appl Ecol*, 26(2):379-387.
- OUYANG Y, EVANS SE, FRIESEN ML, et al., 2018. Effect of nitrogen fertilization on the abundance of nitrogen cycling genes in agricultural soils: a meta-analysis of field studies[J]. *Soil Biol Biochem*, 127:71-78.
- OUYANG Y, NORTON J M, STARK J M, et al., 2016. Ammonia-oxidizing bacteria are more responsive than archaea to nitrogen source in an agricultural soil[J]. *Soil Biol Biochem*, 96:4-15.
- REN YL, 2012. Effects of Precipitation Change on Inorganic Nitrogen and Net Nitrogen Mineralization Rate at a Plantation of Mongolian Pine[J]. *J Peiking Univ(Nat Sci Ed)*, 48(6):925-932.
- SZUKICS U, HACKL E, ZECHMERISTER BS, et al., 2012. Rapid and dissimilar response of ammonia oxidizing archaea and bacteria to nitrogen and water amendment in two temperate forest soils[J]. *Microbiol Res*, 167(2):103-109.
- TIAN J, DUNGAIT JAJ, LU XK, et al., 2019. Long-term nitrogen addition modifies microbial composition and functions for slow carbon cycling and increased sequestration in tropical forest soil[J]. *Glob Chage Biol*, 25(10):3267-3281.
- TANG YQ, YU GR, ZHANG XY, et al., 2019. Environmental variables better explain changes in potential nitrification and denitrification activities than microbial properties in fertilized forest soils[J]. *Sci Total Envir*, 647: 653-662.

TANG YQ, ZHANG XY, LI DD, et al., 2016. Impacts of nitrogen and phosphorus additions on the abundance and community structure of ammonia oxidizers and denitrifying bacteria in Chinese fir plantations[J]. *Soil Biol Biochem*, 103: 284-293.

SONG YJ, MA L, LI M, et al., 2020. Spatial Distribution of ammonia oxidizers in rhizosphere soil of *Phragmites australis* in the Yellow River Delta[J]. *J Shandong Agr Sci*, 52(10):144-150.

WU YF, GAO XP, GUI DW, et al., 2019. Research progress on the monitoring methods of atmospheric nitrogen deposition[J]. *J Appl Ecol*, 30(10):3605-3614.

WANG ZL, LIU CX, JIANG QX, et al., 2021. Effects of climate warming on the key process and index of black soil carbon and nitrogen cycle during freezing period[J]. *Environ Sci*, 42(4):1967-1978.

WANG H, NA D, WU D, et al., 2017. Long-term net transformation and quantitative molecular mechanisms of soil nitrogen during natural vegetation recovery of abandoned farmland on the Loess Plateau of China[J]. *Sci Total Envir*, 607/608(31):152-159.

ZHANG YG, LIU X, CONG J, et al., 2017. The microbially mediated soil organic carbon loss under degenerative succession in an alpine meadow[J]. *Mol Ecol*, 26(14): 3676-3686.

ZHANG C, SONG ZL, ZHUANG DH, et al., 2019 Urea fertilization decreases soil bacterial diversity, but improves microbial biomass, respiration, and N-cycling potential in a semiarid grassland[J]. *Biol Fert Soils*, 55(3): 229-242.

ZHANG M, WANG W, WANG D, et al., 2018. Short-term responses of soil nitrogen mineralization, nitrification and denitrification to prescribed burning in a suburban forest ecosystem of subtropical Australia[J]. *Sci Total Environ*, 642(15):879-886.

ZHENG M, WEI Z, LUO Y, et al., 2017. Different responses of asymbiotic nitrogen fixation to nitrogen addition between disturbed and rehabilitated subtropical forests[J]. *Sci Total Environ*, 601/602:1505-1512.

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv –Machine translation. Verify with original.