

Changes in diversity, composition and assembly processes of soil microbial communities during *Robinia pseudoacacia* L. restoration on the Loess Plateau, China postprint

Authors: WANG Kun, WANG Xiaoxia, FEI Hongyan, WAN Chuanyu, HAN Fengpeng

Date: 2022-06-13T00:00:00+00:00

Abstract

Abstract: *Robinia pseudoacacia* L. (RP) restoration has increased vegetation cover in semi-arid regions on the Loess Plateau of China, but ecological problems have also occurred due to RP restoration, such as reduced soil moisture. Further, it is still uncertain how microbial diversity, composition and assembly processes change with RP restoration in semi-arid regions. Therefore, amplicon sequencing of small subunit ribosomal ribonucleic acid (16S rRNA) and internal transcribed spacer (ITS) genes was performed to study soil bacterial and fungal diversity, composition and assembly processes at four study sites with different stand ages of RP plantations (Y10, RP plantation with stand ages less than 10 a; Y15, RP plantation with stand ages approximately 15 a; Y25, RP plantation with stand ages approximately 25 a; and Y40, RP plantation with stand ages approximately 40 a) along a 40-a chronosequence on the Loess Plateau. The diversity of soil bacteria and fungi increased significantly during the restoration period from 10 to 15 a ($P < 0.05$). However, compared with Y15, bacterial diversity was lower at Y25 and Y40, and fungal diversity remained stable during the restoration period between 25 and 40 a. The relative abundances of Proteobacteria and Ascomycota increased during the restoration period from 10 to 15 a. Conversely, after 15 a of restoration, they both decreased, whereas the relative abundances of Actinomycetes, Acidobacteria and Basidiomycota gradually increased. The variations in soil bacterial communities were mainly related to changes in soil total nitrogen, nitrate nitrogen and moisture contents, while soil fungal communities were mainly shaped by soil organic carbon and nitrate nitrogen contents. Bacterial communities were structured by the heterogeneous selection and stochastic process, while fungal communities were structured primarily by the stochastic process. The RP restoration induced an increase in

the relative importance of heterogeneous selection on bacterial communities. Overall, this study reveals the changes in microbial diversity, community composition and assembly processes with RP restoration on the Loess Plateau and provides a new perspective on the effects of vegetation restoration on soil microbial communities in semi-arid regions.

Full Text

Preamble

Changes in Diversity, Composition and Assembly Processes of Soil Microbial Communities During *Robinia pseudoacacia* L. Restoration on the Loess Plateau, China

Kun Wang¹, Xiaoxia Wang¹, Hongyan Fei¹, Chuanyu Wan¹, Fengpeng Han^{1, 2*}

¹State Key Laboratory of Soil Erosion and Dryland Farming on the Loess Plateau, College of Natural Resources and Environment, Northwest A & F University, Yangling 712100, China

²Research Center on Soil & Water Conservation, Institute of Soil and Water Conservation, Chinese Academy of Sciences and Ministry of Water Resources, Yangling 712100, China

Abstract: *Robinia pseudoacacia* L. (RP) restoration has increased vegetation cover in semi-arid regions on the Loess Plateau of China, but ecological problems have also occurred due to RP restoration, such as reduced soil moisture. Further, it is still uncertain how microbial diversity, composition and assembly processes change with RP restoration in semi-arid regions. Therefore, amplicon sequencing of small subunit ribosomal ribonucleic acid (16S rRNA) and internal transcribed spacer (ITS) genes was performed to study soil bacterial and fungal diversity, composition and assembly processes at four study sites with different stand ages of RP plantations (Y10, RP plantation with stand ages less than 10 a; Y15, RP plantation with stand ages approximately 15 a; Y25, RP plantation with stand ages approximately 25 a; and Y40, RP plantation with stand ages approximately 40 a) along a 40-a chronosequence on the Loess Plateau. The diversity of soil bacteria and fungi increased significantly during the restoration period from 10 to 15 a ($P < 0.05$). However, compared with Y15, bacterial diversity was lower at Y25 and Y40, and fungal diversity remained stable during the restoration period between 25 and 40 a. The relative abundances of Proteobacteria and Ascomycota increased during the restoration period from 10 to 15 a. Conversely, after 15 a of restoration, they both decreased, whereas the relative abundances of Actinomycetes, Acidobacteria and Basidiomycota gradually increased. The variations in soil bacterial communities were mainly related to changes in soil total nitrogen, nitrate nitrogen and moisture contents, while soil fungal communities were mainly shaped by soil organic carbon and nitrate nitrogen contents. Bacterial communities were structured by heterogeneous selection

and stochastic process, while fungal communities were structured primarily by stochastic process. The RP restoration induced an increase in the relative importance of heterogeneous selection on bacterial communities. Overall, this study reveals the changes in microbial diversity, community composition and assembly processes with RP restoration on the Loess Plateau and provides a new perspective on the effects of vegetation restoration on soil microbial communities in semi-arid regions.

Keywords: *Robinia pseudoacacia*; microbial diversity; community structure; community assembly; restoration; semi-arid regions

Citation: Wang K, Wang X, Fei H, Wan C, Han F. 2022. Changes in diversity, composition and assembly processes of soil microbial communities during *Robinia pseudoacacia* L. restoration on the Loess Plateau, China. *Journal of Arid Land*, 14(5): 561–575. <https://doi.org/10.1007/s40333-022-0064-2>

Corresponding author: Fengpeng Han (E-mail: hanxiangzi007@163.com)

Received 2021-12-27; revised 2022-04-01; accepted 2022-04-06

© Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Science Press and Springer-Verlag GmbH Germany, part of Springer Nature 2022

Introduction

Vegetation restoration of degraded farmlands has been widely implemented to improve soil quality and enhance the ecological and economic values of land in arid and semi-arid regions with fragile ecosystems [?, ?]. *Robinia pseudoacacia* L. (RP), a nitrogen (N)-fixing tree species, has been selected as an important pioneer species for the conversion of farmland to forests. RP is planted worldwide due to its tolerance to diverse soil conditions, surviving in soil with pH ranging from extremely acidic to strongly alkaline, as well as in soil with base-saturation levels ranging from medium to high [?, ?]. During RP restoration, originally barren land undergoes a series of soil processes, including carbon (C) and N input and transformation, and nutrient accumulation [?, ?]. Soil microbes play important roles in mediating these processes. For example, soil fungi complete the decomposition of plant litter, and bacteria primarily participate in the biogeochemical cycle of soil C and N [?, ?]. In addition, metrics related to microbial communities (i.e., diversity, microbial biomass and presence of beneficial microbes) can be used as important indicators to evaluate the success of the restoration of degraded ecosystems. For example, soil bacterial diversity can be recovered to pre-farmland levels after approximately 20 a of secondary succession on abandoned farmland and this process is highly dependent on the availability of soil nutrients [?]. Soil microbial biomass, one of the main nutrient sources of plants in nutrient-restricted areas, accumulates during restoration, and it is thus considered as an important indicator of soil fertil-

ity [?, ?]. Increases in the diversity of N-fixing microbes associated with RP restoration were significantly positively correlated with the amount of available N in soils on the Loess Plateau, China [?]. Therefore, studying soil microbial communities can increase our understanding of the succession of ecosystems and enable restoration.

RP restoration significantly affects the diversity and composition of soil microbial communities [?, ?]. On the one hand, during RP restoration, the increase in plant litter provides more substrates for decomposition by microbes [?, ?]. On the other hand, RP restoration changes soil properties, such as reducing the pH and bulk density of soil, as well as increasing the C and N contents, which have significant impacts on the metabolism and activity of soil microbes. For example, Liu et al. (2018) reported that during a 35-a RP restoration, Acidobacteria eventually replaced other species to become the dominant phylum in soil, and phosphorus (P) content was the main factor shaping the composition of microbial communities. Xu et al. (2019) found that compared with farmland, 42 a of RP restoration significantly improved the diversity index and species richness of soil bacteria and fungi. Recently, however, the problem of soil moisture (SM) decline following RP restoration has become particularly serious. Several studies have found that SM showed a linear decreasing trend with increasing recovery age in different precipitation zones (400.0–600.0 mm), which led to increased RP plantation mortality and land degradation [?, ?, ?]. These findings suggest that the response of soil microbes to vegetation restoration may be more complex in semi-arid regions. Therefore, it is necessary to reassess the response of soil microbial communities to RP restoration in semi-arid regions.

Microbial community assembly is an important research topic in ecology [?]. In contrast to the body of knowledge regarding large organisms, understanding of the mechanisms of microbial community assembly is still very limited. Currently, it is generally believed that microbial community assembly is driven by both deterministic process (environmental selection) and stochastic process (dispersal, ecological drift and diversification). Moreover, the importance of these two types of processes has been quantified by using many models [?, ?]. Liu et al. (2021) found that the assembly of prokaryotic microbial communities was controlled by deterministic process in the early succession of subtropical forests. In contrast, the relative influence of stochastic process increased in the later stage. Osburn et al. (2021) revealed that the assembly of soil bacterial and fungal communities was primarily affected by deterministic process in historically disturbed forests (e.g., logging and agricultural conversion). In pristine forests that have remained undisturbed for approximately 100 a, stochastic process is more important. Despite preliminary studies of soil microbial assembly processes in forests, the assembly processes along a chronosequence (i.e., RP restoration) are still unclear, especially in semi-arid regions. This is crucial for understanding the drivers of soil microbial community succession.

The Loess Plateau is a semi-arid region in China with severe soil erosion and ecological fragility [?]. RP has been widely planted in this region and has been

a pioneer species for afforestation of farmland since the 1970s [?]. Although RP restoration increased the coverage of local vegetation, it also caused a series of environmental problems, e.g., declines in SM [?]. To date, the variation in microbial communities and their ecological processes during RP restoration on the Loess Plateau have not been fully explored. Here, based on the spatial (as opposed to temporal) method, we studied soil bacterial and fungal communities at RP plantations with different stages on the Loess Plateau. Our aims were to (1) determine the variation in the diversity and composition of soil bacterial and fungal communities at RP plantations with different stages; (2) illustrate important soil parameters related to this variation; and (3) determine the ecological process that drives the assembly of soil bacterial and fungal communities.

2.1 Study Area

The study area is located in a loess hilly and gully region on the Loess Plateau, China. The loess thickness in this area is 30–180 m and the soil type is mainly loessial soil developed from loess parent material. The area is characterized by a temperate and semi-arid climate, with the annual mean temperature of 8.8°C and the mean annual precipitation of 505.3 mm. The average altitude in the study area is 1371.6 m, the average annual sunshine duration is 2395.6 h, and the annual frost-free period is 157 d.

2.2 Soil Sampling

In April 2019, we identified four RP plantations at four study sites along a chronosequence as follows: Y10, RP plantation with stand ages less than 10 a; Y15, RP plantation with stand ages approximately 15 a; Y25, RP plantation with stand ages approximately 25 a; and Y40, RP plantation with stand ages approximately 40 a. Apart from differences in stand ages, the four plantations were similar in terms of geographical characteristics and soil types, and they were all agricultural lands with maize and wheat as the main crops before RP restoration. The different plantations are located more than 5 km apart. After RP restoration, these plantations were rarely managed by humans; thus, their development can be considered a natural restoration process. Previous study has shown that on the Loess Plateau, the mature period of RP plantations is 20 a [?]. Therefore, Y25 and Y40 represent the later restoration stage. More information on the study sites is shown in Table S1.

To avoid the problem of ecological pseudoreplication as much as possible, we established five 10 m × 10 m plots with distance no more than 50 m from each other at each site, and the sampling area at each site was approximately 10 ha (100 m × 100 m). In each plot, five locations for soil cores were randomly selected. First, the litter on the surface of soil was carefully removed. Then, bulk soil samples were collected from the top soil with depth of 0–20 cm using a 5 cm diameter stainless steel corer. Finally, the litter and roots in each

soil sample were removed, and the five individual soil samples collected from the same plot were sieved using a 2 mm mesh and fully mixed into a single soil sample. In this way, a total of 20 soil samples were obtained. Each soil sample was divided into two subsamples. The first subsample was transported to the laboratory on ice and stored at -80°C until deoxyribonucleic acid (DNA) extraction was performed. The second subsample was used for soil physical and chemical analyses.

2.3 Analyses of Soil Physical and Chemical Properties

Soil pH was measured (1.0:2.5 soil/water) using a pH metre (IS126, Insmark Inc., Shanghai, China). SM was measured by a FieldScout time domain reflectometry (TDR) moisture metre sensor (TDR 300, Spectrum Technologies Inc., Plainfield, USA). Soil organic carbon (SOC) was determined by the $\text{K}_2\text{Cr}_2\text{O}_7\text{-H}_2\text{SO}_4$ oxidation method [?]. Total nitrogen (TN) was determined using the Kjeldahl method [?]. Nitrate nitrogen (NO_3^- -N) and ammonium nitrogen (NH_4^+ -N) were extracted with 1 mol/L KCl and determined by a continuous flow analytical system. Available phosphorus (AP) was determined by the molybdenum antimony colorimetric method after extraction with 0.5 mol/L NaHCO_3 [?].

2.7 Statistical Analyses

Significant differences ($P < 0.05$) in soil properties and alpha diversity indices between RP plantations with different stand ages were tested by one-way analysis of variance (ANOVA). The data were log transformed if necessary. Pearson correlation analysis was used to summarize the relationship between microbial diversity indices and soil properties. Principal component analysis (PCA) was used to describe the differences between soil properties among different RP plantations. Principal coordinate analysis (PCoA) based on the Bray-Curtis distance was conducted to visualize the variations in community structure, while the Adonis test was used to examine for significant differences in community structure between RP plantations with different stand ages (based on 999 iterations). Redundancy analysis (RDA) was conducted to evaluate the effects of soil properties on bacterial and fungal community structures using CANOCO software v5. Linear regression was further used to examine the relationship between the temporal distance among samples and similarity in microbial composition (Bray-Curtis distance). A forward selection procedure was performed to reveal the significant explanatory variables by Monte Carlo permutations. All the statistical analyses mentioned above were performed in R 3.6.3 software without further modification.

3.1 Soil Properties

The contents of SOC, TN, NH_4^+ -N and AP increased gradually from Y10 to Y40, with significant increases of 1.60-, 1.50-, 1.00- and 0.92-fold, respectively

($P < 0.05$; Table 1). The content of NO_3^- -N decreased gradually from Y10 to Y40 ($P < 0.05$). The soil pH continuously decreased from Y10 to Y40. SM was the lowest at Y25 and Y40 and the highest at Y15 ($P < 0.05$). Based on the measured soil characteristics, the four stand ages of RP plantations were distinctly separated along the first principal component (PC) coordinate axis (Fig. S1), explaining more than 78.0% of the total variation. Overall, these results indicated that the restoration of RP significantly altered soil properties.

3.2 Diversity of Microbial Communities

After quality trimming and chimera removal, the data set was found to contain a total of 794,499 high-quality bacterial sequences and 1,011,433 high-quality fungal sequences for the soil samples. The average numbers of bacterial OTUs at Y10, Y15, Y25 and Y40 were 2452 (± 18), 2493 (± 29), 2277 (± 86) and 2471 (± 183), respectively. A Venn diagram showed that 3183, 2975 and 2988 bacterial OTUs at Y15, Y25 and Y40 were shared with Y10, respectively, and 639, 590 and 559 fungal OTUs at Y15, Y25 and Y40 were shared with Y10, respectively (Fig. S2), suggesting that compared with Y10, the similarity of microbial communities generally decreased with restoration time. The rarefaction curves appeared to flatten, and Good's coverage reached 99.98%, collectively indicating that the amount of sequencing data was reasonable.

For bacteria, the Shannon and Chao indices both reached their peak values at Y15 and then significantly decreased to their lowest values at Y25 and Y40 ($P < 0.05$; Fig. 1a [Figure 1: see original paper] and b). In contrast, the Shannon and Chao index values for fungi showed a sharp increasing tendency from Y10 to Y25, with significant increases of 19.8% and 29.7% ($P < 0.05$), respectively, and both of them exhibited a slightly increasing trend from Y25 to Y40 ($P > 0.05$; Fig. 1c and d). Pearson correlation analysis (Table S2) revealed that SM was significantly positively related to the bacterial Shannon and Chao indices, while the increases in SOM, TN, NH_4^+ -N and AP were significantly negatively related to the bacterial Shannon index ($P < 0.05$). For fungi, the increases in SOM, TN, NH_4^+ -N and AP were all significantly positively related to the Shannon and Chao indices, and the decrease in pH was significantly negatively related to the Shannon and Chao indices ($P < 0.05$).

3.3 Relative Abundance of Dominant Phyla

At the bacterial phylum level (Fig. 2a [Figure 2: see original paper]), Proteobacteria (19.1%–34.5%), Actinobacteria (21.9%–32.8%) and Acidobacteria (14.3%–20.4%) were the dominant phyla across different RP plantations. The relative abundance of Proteobacteria was lowest at Y25 and Y40 and highest at Y15 ($P < 0.05$). The relative abundance of Actinobacteria was highest at

Y25 and Y40 and lowest at Y10 ($P < 0.05$), and the relative abundance of Acidobacteria reached the highest value at Y40 ($P < 0.05$). At the fungal phylum level (Fig. 2b), Ascomycota (66.2%–75.0%) was the dominant phylum across different RP plantations, followed by Zygomycota (15.0%–20.8%) and Basidiomycota (3.9%–12.3%). The relative abundance of Ascomycota significantly decreased from Y15 to Y25 and that of Basidiomycota increased gradually from Y10 to Y40 ($P < 0.05$).

3.4 Microbial Community Variation and Determinants

Based on the Bray-Curtis distance (Fig. 3a [Figure 3: see original paper] and b), PCoA showed that bacterial and fungal community structures were clearly separated along the first and second axes among the four RP plantations. Accordingly, the Adonis test showed that bacterial and fungal community structures across years were extremely significantly different ($P < 0.05$; Table S3). Linear regression analyses described the changes in microbial beta diversity over time. The results revealed that bacteria showed a larger slope (temporal turnover) than fungi based on the Bray-Curtis distance (Fig. S3).

According to RDA (Fig. 3c and d), the variations in bacterial and fungal community compositions were mostly affected by the variations in soil factors, which explained 67.1% and 56.7% of the bacterial and fungal community variance, respectively. According to forward selection (Table S4), the dissimilarities in TN, NO_3^- -N and SM were closely related to the dissimilarity of bacterial community composition across the RP plantations ($P < 0.01$), and those of SOC and NO_3^- -N were closely related to the dissimilarity of fungal community composition across the RP plantations.

3.5 Assembly Processes of Microbial Communities

We compared the relative importance of the two types of assembly processes (deterministic and stochastic processes) of bacterial and fungal communities at different RP plantations based on the betaNTI. The assembly of bacterial communities was driven by both deterministic and stochastic processes (Fig. 4 [Figure 4: see original paper]), and the deterministic process gradually became the main force (50.0% to 70.0% from Y10 to Y40) with restoration time. Surprisingly, all betaNTI values of fungal communities were between -2 and 2 during the restoration time period from 10 to 25 a, indicating that the stochastic process dominated the assembly of fungal communities. The deterministic process became one of the forces (20.0%) for the assembly of fungal communities at Y40.

4 Discussion

During the early RP restoration period (stand ages from 10 to 15 a), the alpha diversity index of soil bacteria and fungi showed an upwards trend (Fig. 1). This is consistent with the results of previous studies [?, ?, ?]. Similarly, the restoration of other types of vegetation, such as *Platycladus orientalis* and Masson pine, also increases soil microbial alpha diversity [?, ?]. This increase might occur because vegetation restoration in the early period promotes more diverse ecological niches and high availability of resources for microbes [?]. However, in this study, we found that the variation patterns of soil microbial alpha diversity during the later RP restoration period (stand ages from 15 to 40 a) were not consistent with those during the early RP restoration period (Fig. 1).

Liang et al. (2018) proposed that large-scale (an area of approximately 6.4×10^5 km²) and long-term (more than 30 a) vegetation restoration leads to a decrease in SM on the Loess Plateau, and our study also found a decline in SM during the later restoration period (Table 1). SM was an important resource for the survival of microbes. The reduction in SM directly affected the metabolic activities of microbes and decreased the utilization of microbial C and N resources [?]. Moreover, we found that there was a significant positive correlation between bacterial diversity index and SM during the RP restoration period, while there were significant negative correlations of bacterial diversity index with SOC and soil nutrients (TN, NH₄⁺-N and AP; Table S2). Compared to bacteria, soil fungi were more resistant to the declines in SM [?], which may explain why the fungal alpha diversity only exhibited a slow trend during the RP restoration period (Fig. 1). Furthermore, Pearson correlation analysis showed that the fungal alpha diversity was significantly positively related to SOC and soil nutrients but negatively related to soil pH (Table S2), suggesting that the soil fungal alpha diversity indices were more dependent on SOC, soil nutrients and pH during the RP restoration period. Notably, the fungal alpha diversity indices increased slowly during the later restoration period (Fig. 1). Similar to the dynamics of the secondary plant succession, the increased lignin in litter indicated the decreased resource availability of soil fungi, which may limit the number of some fungi [?, ?]. In addition, due to the increased dominance of a particular tree species, the narrowing of the range of organic substrates in soil during the late succession may limit the number of niches for saprophytic fungi [?]. Thus, we speculate that variations in litter quality and vegetation composition might limit the fungal diversity despite improvements in soil nutrients and pH.

Results of RDA showed that soil N and SM were more closely related to the variations in bacterial community compositions (Fig. 3; Table S4). This is because soil nutrients and water are important resources for microbes. Changes in resource availability directly affect the metabolism and compositions of microbes [?, ?, ?]. These results have been extensively studied in terrestrial ecosystems. During the early RP restoration period, soil N showed an overall upwards trend. Sufficient labile substrates are conducive to the growth of Proteobacteria, a type of microbe that have faster access to substrates than other microbes [?].

Therefore, after 15 a of restoration, the reduction in soil NO_3^- -N might hinder the growth of eutrophic microbes. In addition, the decline in SM stimulated the growth of some drought-tolerant bacteria, such as Actinobacteria and Acidobacteria, which were considered as the dominant phyla in harsh and stressful environments [?, ?]. For fungi, SOC content was closely related to the community compositions. This result is supported by the study of Liu et al. (2021), who found that during vegetation restoration, the types of SOC become more complex and are difficult to degrade. However, only a small number of fungi can secrete enzymes that catalyse the degradation of large organic C molecules, such as lignin. To date, the microbes that can secrete these enzymes mainly exist in the higher Basidiomycota [?, ?]. Therefore, in this study, we found that during the later RP restoration period, the relative abundance of Ascomycota decreased, while the relative abundance of Basidiomycota began to increase. As was the case for fungi, soil NO_3^- -N was also an important factor (Table S4). In terrestrial ecosystems, reduced available N content often leads to intense competition between microorganisms and plants [?]. In this study, as the amount of available N increased, the classification and functional characteristics of soil fungi changed; for example, the relative abundance of mycorrhizal fungi was reduced [?].

Based on our calculated betaNTI values, we found greater effects of the deterministic process on bacterial communities than on fungal communities. It has also been previously reported that bacteria are more susceptible to environmental factors than fungi [?, ?]. Many environmental factors can shape bacterial communities on spatial or temporal scales, such as climate [?], soil pH [?], soil nutrient availability [?], SM [?], etc. In this study, we found that SM was an important nonbiological factor that drove bacterial diversity and community compositions but did not have the same effect on fungi (Tables S2 and S4). Environmental factors may have less impact on fungal communities. This may be because fungi promote the acquisition of SM and soil nutrients through complex hyphal networks, which in return make them more resilient to environmental changes than bacteria [?]. Consistent with this conclusion, we found that during the 40-a restoration of RP plantations, the turnover rate of soil fungal communities was lower than that of soil bacterial communities (Fig. S3). In addition, some species of fungi can produce tens of billions of spores per square kilometer each year, promoting diffusion in ecosystems [?]. In short, the differences in the community assembly of bacteria and fungi may reflect the differences in the life strategies (e.g., nutrient acquisition) of bacteria and fungi during the RP restoration period.

In addition, we found that the relative importance of the deterministic effect on bacterial communities gradually increased with increasing restoration time. During the initial RP restoration period (5 a), the historical manual farmland management, such as ploughing, fertilization and irrigation, may provide an ideal condition for the diffusion of bacteria [?]. Previous study has suggested that the importance of the stochastic process generally increases as the supply of SOC increases [?]. However, this study suggested that the relative importance

of the deterministic process was negatively correlated with SOC content. This may be because RP restoration had a stronger selection effect on bacteria than the supply of SOC by reducing soil NO_3^- -N and SM. Further work is needed to study the thresholds of these environmental factors that trigger a decrease in the importance of the stochastic process in bacterial communities. In addition, we observed that the heterogeneous selection effect of bacteria was strengthened during the RP restoration period (Fig. 4), which may be related to the topography of the study sites. Generally, there are large differences in SM and soil nutrients between the top and bottom of a slope, especially in the erosion-prone Loess Plateau region [?]. Therefore, the environmental heterogeneity in the region may gradually increase with increasing restoration time.

5 Conclusions

The diversity of soil bacteria and fungi showed an increasing trend during the early RP restoration period (from 10 to 15 a) on the Loess Plateau, and Proteobacteria and Ascomycota were the dominant phyla for bacteria and fungi, respectively. However, during the later RP restoration period (from 15 to 40 a), soil bacterial diversity was significantly lower than that during the early RP restoration period, and the rate of increase in fungal diversity decreased. During the later RP restoration period, the relative abundances of Proteobacteria and Ascomycota were lower than those during the early RP restoration period, while the relative abundances of Actinobacteria, Acidobacteria and Basidiomycota were greater during this period. The variations in soil bacterial communities were mainly related to the changes in soil total N, nitrate N and SM, while soil fungal communities were mainly shaped by SOC and soil nitrate N. Bacterial communities were structured by heterogeneous selection and stochastic process, while fungal communities were structured primarily by stochastic process. The restoration of RP plantations induced an increase in the relative importance of heterogeneous selection on bacterial communities. Overall, this study reveals the changes in microbial diversity, community compositions and assembly processes during the RP restoration period (40-a chronosequence) on the Loess Plateau and provides a new perspective for studying the effects of vegetation restoration on soil microbial communities in semi-arid regions.

Acknowledgements

This study was supported by the National Natural Science Foundation of China (41471437), the National Key R&D Program of China (2016YFA0600801, 2017YFC0504504), the West Light Foundation of the Chinese Academy of Science (XAB2016A04), and the Key R&D Program of Ningxia Hui Autonomous Region, China (2022BBF02033). We thank all helpful comments for this manuscript.

References

- Adams R I, Miletto M, Taylor J W, et al. 2013. Dispersal in microbes: fungi in indoor air are dominated by outdoor air and show dispersal limitation at short distances. *The ISME Journal*, 7(7): 1262–1273.
- Arenas F, Navarro-Ródenas A, Marqués-Gálvez J E, et al. 2021. Different patterns in root and soil fungal diversity drive plant productivity of the desert truffle *Terfezia claveryi* in plantation. *Environmental Microbiology*, 23(10): 5917–5933.
- Banerjee S, Misra A, Sar A, et al. 2020. Poor nutrient availability in open-cast coalmine influences microbial community composition and diversity in exposed and underground soil profiles. *Applied Soil Ecology*, 152: 103544, doi: 10.1016/j.apsoil.2020.103544.
- Barka E A, Vatsa P, Sanchez L, et al. 2016. Taxonomy, physiology, and natural products of Actinobacteria. *Microbiology and Molecular Biology Reviews*, 80(1): 1–43.
- Biddle J F, Fitz-Gibbon S, Schuster S C, et al. 2008. Metagenomic signatures of the Peru Margin seafloor biosphere show a genetically distinct environment. *Proceedings of the National Academy of Sciences of the United States of America*, 105(30): 10583–10588.
- Cline L C, Zak D R. 2015. Soil microbial communities are shaped by plant-driven changes in resource availability during secondary succession. *Ecology*, 96(12): 3374–3385.
- Crits-Christoph A, Robinson C K, Barnum T, et al. 2013. Colonization patterns of soil microbial communities in the Atacama Desert. *Microbiome*, 1: 28, doi: 10.1186/2049-2618-1-28.
- de Menezes A B, Richardson A E, Thrall P H. 2017. Linking fungal-bacterial co-occurrences to soil ecosystem function. *Current Opinion in Microbiology*, 37: 135–141.
- de Vries F T, Griffiths R I, Bailey M, et al. 2018. Soil bacterial networks are less stable under drought than fungal networks. *Nature Communications*, 9(1): 1–12.
- Delgado-Baquerizo M, Oliverio A M, Brewer T E, et al. 2018. A global atlas of the dominant bacteria found in soil. *Science*, 359(6373): 320–325.
- Dini-Andreote F, Stegen J C, Van Elsas J D, et al. 2015. Disentangling mechanisms that mediate the balance between stochastic and deterministic processes in microbial succession. *Proceedings of the National Academy of Sciences of the United States of America*, 112(11): e1326–e1332.

- Dong X D, Gao P, Zhou R, et al. 2021. Changing characteristics and influencing factors of the soil microbial community during litter decomposition in a mixed *Quercus acutissima* Carruth. and *Robinia pseudoacacia* L. forest in Northern China. *CATENA*, 196: 104811, doi: 10.1016/j.catena.2020.104811.
- Dumbrell A J, Nelson M, Helgason T, et al. 2010. Relative roles of niche and neutral processes in structuring a soil microbial community. *The ISME Journal*, 4(3): 337–345.
- Fierer N, Bradford M A, Jackson R B. 2007. Toward an ecological classification of soil bacteria. *Ecology*, 88(6): 1354–1364.
- Guo X, Feng J J, Shi Z, et al. 2018. Climate warming leads to divergent succession of grassland microbial communities. *Nature Climate Change*, 8(9): 813–818.
- Hu P L, Xiao J, Zhang W, et al. 2020. Response of soil microbial communities to natural and managed vegetation restoration in a subtropical karst region. *CATENA*, 195: 104849, doi: 10.1016/j.catena.2020.104849.
- Jin Z, Li X R, Wang Y Q, et al. 2016. Comparing watershed black locust afforestation and natural revegetation impacts on soil nitrogen on the Loess Plateau of China. *Scientific Reports*, 6: 25048, doi: 10.1038/srep25048.
- Knoke T, Bendix J, Pohle P, et al. 2014. Afforestation or intense pasturing improve the ecological and economic value of abandoned tropical farmlands. *Nature Communications*, 5(1): 1–12.
- Knops J M H, Tilman D. 2000. Dynamics of soil nitrogen and carbon accumulation for 61 years after agricultural abandonment. *Ecology*, 81(1): 88–98.
- Leff J W, Jones S E, Prober S M, et al. 2015. Consistent responses of soil microbial communities to elevated nutrient inputs in grasslands across the globe. *Proceedings of the National Academy of Sciences of the United States of America*, 112(35): 10967–10972.
- Li G Y, Zhang M X, Wu C F. 2021. Agricultural land consolidation impacted edaphic microbial community dynamics and assembly in China: a case study from Zhejiang Province. *CATENA*, 205: 105424, doi: 10.1016/j.catena.2021.105424.
- Liang H B, Xue Y Y, Li Z S, et al. 2018. Soil moisture decline following the plantation of *Robinia pseudoacacia* forests: Evidence from the Loess Plateau. *Forest Ecology and Management*, 412: 62–69.
- Liu H F, Xu H W, Wu Y, et al. 2021. Effects of natural vegetation restoration on dissolved organic matter (DOM) biodegradability and its temperature sensitivity. *Water Research*, 191: 116792, doi: 10.1016/j.watres.2020.116792.
- Liu J L, Yang Z L, Dang P, et al. 2018. Response of soil microbial community dynamics to *Robinia pseudoacacia* L. afforestation in the loess plateau: a chronosequence approach. *Plant and Soil*, 423(1): 327–338.

- Liu L, Zhu K, Krause S, et al. 2021. Changes in assembly processes of soil microbial communities during secondary succession in two subtropical forests. *Soil Biology and Biochemistry*, 154: 108144, doi: 10.1016/j.soilbio.2021.108144.
- Ma L, Guo C, Lv X, et al. 2015. Soil moisture and land use are major determinants of soil microbial community composition and biomass at a regional scale in northeastern China. *Biogeosciences*, 12(8): 2585–2596.
- Nelson D W, Sommers L E. 1983. Total carbon, organic carbon, and organic matter. In: Page A. *Methods of Soil Analysis*. Madison: Agronomy Society of America, 539–579.
- Olsen S R, Sommers L E. 1982. Phosphorous. In: Page A. *Methods of Soil Analysis*. Madison: Agronomy Society of America, 403–430.
- Osburn E D, Aylward F O, Barrett J E. 2021. Historical land use has long-term effects on microbial community assembly processes in forest soils. *ISME Communications*, 1: 48, doi: 10.1038/s43705-021-00051-x.
- Papaioannou A, Chatzistathis T, Papaioannou E, et al. 2016. *Robinia pseud-acacia* as a valuable invasive species for the restoration of degraded croplands. *CATENA*, 137: 310–317.
- Peay K G, Schubert M G, Nguyen N H, et al. 2012. Measuring ectomycorrhizal fungal dispersal: macroecological patterns driven by microscopic propagules. *Molecular Ecology*, 21(16): 4122–4136.
- Powell J R, Karunaratne S, Campbell C D, et al. 2015. Deterministic processes vary during community assembly for ecologically dissimilar taxa. *Nature Communications*, 6(1): 1–10.
- Purahong W, Wubet T, Lentendu G, et al. 2016. Life in leaf litter: novel insights into community dynamics of bacteria and fungi during litter decomposition. *Molecular Ecology*, 25(16): 4059–4074.
- Qiu L P, Zhang Q, Zhu H S, et al. 2021. Erosion reduces soil microbial diversity, network complexity and multifunctionality. *The ISME Journal*, 15: 2474–2489.
- Rédei K, Csiha I, Keserú Z, et al. 2014. Growth and yield of black locust (*Robinia pseudoacacia* L.) stands in Nyírség growing region (North-East Hungary). *South-East European Forestry*, 5(1): 13–22.
- Romaní A M, Fischer H, Mille-Lindblom C, et al. 2006. Interactions of bacteria and fungi on decomposing litter: differential extracellular enzyme activities. *Ecology*, 87(10): 2559–2569.
- Rong G H, Li W R, Zhu H S, et al. 2020. Dynamics of new- and old- organic carbon and nitrogen in bulk soils and aggregates following afforestation on farmland. *CATENA*, 195: 104838, doi: 10.1016/j.catena.2020.104838.
- Schimel J P. 2018. Life in dry soils: effects of drought on soil microbial communities and processes. *Annual Review of Ecology, Evolution, and Systematics*,

49(1): 409–432.

Shao P S, Liang C, Lynch L, et al. 2019. Reforestation accelerates soil organic carbon accumulation: Evidence from microbial biomarkers. *Soil Biology and Biochemistry*, 131: 182–190.

Singh J S, Singh D, Kashyap A K. 2009. A comparative account of the microbial biomass-N and N-mineralization of soils under natural forest, grassland and crop field from dry tropical region, India. *Plant, Soil and Environment*, 55(6): 223–230.

Stegen J C, Lin X J, Fredrickson J K, et al. 2013. Quantifying community assembly processes and identifying features that impose them. *The ISME Journal*, 7(11): 2069–2079.

Su B Q, Shangguan Z P. 2019. Decline in soil moisture due to vegetation restoration on the Loess Plateau of China. *Land Degradation & Development*, 30(3): 290–299.

Sun S Y, Sun H, Zhang D H, et al. 2019. Response of soil microbes to vegetation restoration in coal mining subsidence areas at Huaibei coal mine, China. *International Journal of Environmental Research and Public Health*, 16(10): 1757, doi: 10.3390/ijerph16101757.

Tiemann L K, Billings S A. 2011. Changes in variability of soil moisture alter microbial community C and N resource use. *Soil Biology and Biochemistry*, 43(9): 1837–1847.

Vitkova M, Tonika J, Muellerova J. 2015. Black locust—Successful invader of a wide range of soil conditions. *Science of the Total Environment*, 505: 315–328.

Wang P D, Li S P, Yang X, et al. 2020. Mechanisms of soil bacterial and fungal community assembly differ among and within islands. *Environmental Microbiology*, 22(4): 1559–1571.

Wang Y Y, Chen Y L, Xue Q Y, et al. 2021. The abundance of the *nifH* gene became higher and the *nifH*-containing diazotrophic bacterial communities changed during primary succession in the Hailuoguo Glacier chronosequence, China. *Frontiers in Microbiology*, 12: 672656, doi: 10.3389/fmicb.2021.672656.

Wei J S, Li Z S, Feng X Y, et al. 2018. Ecological and physiological mechanisms of growth decline of *Robinia pseudoacacia* plantations in the Loess Plateau of China: A review. *The Journal of Applied Ecology*, 29(7): 2433–2444. (in Chinese)

Wen X, Théau J. 2020. Spatiotemporal analysis of water-related ecosystem services under ecological restoration scenarios: a case study in northern Shaanxi, China. *Science of the Total Environment*, 720: 137477, doi: 10.1016/j.scitotenv.2020.137477.

Williamson, D F, Parker R A, Kendrick J S. 1989. The box plot: a simple visual method to interpret data. *Annals of Internal Medicine*, 110: 916–921.

Xu M P, Gao D X, Fu S Y, et al. 2020a. Long-term effects of vegetation and soil on the microbial communities following farmland afforestation with *Robinia pseudoacacia* plantations. *Geoderma*, 114263, doi: 10.1016/j.geoderma.2020.114263.

Xu M P, Lu X Q, Xu Y D, et al. 2020b. Dynamics of bacterial community in litter and soil along a chronosequence of *Robinia pseudoacacia* plantations. *Science of the Total Environment*, 703: 135613, doi: 10.1016/j.scitotenv.2019.135613.

Xu Y D, Wang T, Li H, et al. 2019. Variations of soil nitrogen-fixing microorganism communities and nitrogen fractions in a *Robinia pseudoacacia* chronosequence on the Loess Plateau of China. *CATENA*, 174: 316–323.

Yang Y, Dou Y X, Huang Y M, et al. 2017. Links between soil fungal diversity and plant and soil properties on the Loess Plateau. *Frontiers in Microbiology*, 8: 2198, doi: 10.3389/fmicb.2017.02198.

Yuste J C, Peñuelas J, Estiarte M, et al. 2011. Drought-resistant fungi control soil organic matter decomposition and its response to temperature. *Global Change Biology*, 17(3): 1475–1486.

Zak D R, Holmes W E, White D C, et al. 2003. Plant diversity, soil microbial communities, and ecosystem function: are there any links? *Ecology*, 84(8): 2042–2050.

Zhang C, Liu G B, Xue S, et al. 2016. Soil bacterial community dynamics reflect changes in plant community and soil properties during the secondary succession of abandoned farmland in the Loess Plateau. *Soil Biology and Biochemistry*, 97: 40–49.

Zhang W, Liu W C, Xu M P, et al. 2019. Response of forest growth to C: N: P stoichiometry in plants and soils during *Robinia pseudoacacia* afforestation on the Loess Plateau, China. *Geoderma*, 337(6): 280–289.

Zhao H F, He H M, Wang J J, et al. 2018. Vegetation restoration and its environmental effects on the Loess Plateau. *Sustainability*, 10(12): 4676, doi: 10.3390/su10124676.

Zheng, F L, Wang, B. 2014. Soil erosion in the Loess Plateau region of China. In: Tsunekawa A, Liu G, Yamanaka N, et al. *Restoration and Development of the Degraded Loess Plateau, China*. Tokyo: Ecological Research Monographs, 77–92.

Zhou H, Zhang D G, Jiang Z H, et al. 2019. Changes in the soil microbial communities of alpine steppe at Qinghai-Tibetan Plateau under different degradation levels. *Science of the Total Environment*, 651: 2281–2291.

Zhou J Z, Ning D L. 2017. Stochastic community assembly: does it matter in microbial ecology? *Microbiology and Molecular Biology Reviews*, 81(4): e00002–e00017.

Zuo X A, Wang S K, Lv P, et al. 2016. Plant functional diversity enhances associations of soil fungal diversity with vegetation and soil in the restoration of semiarid sandy grassland. *Ecology and Evolution*, 6(1): 318–328.

Appendix

Table S1 Geographical characteristics of *Robinia pseudoacacia* L. (RP) plantations with different stand ages

plantation	Location	Altitude (m)	ATH (m)	Slope gradient (°)	Trees density (numbers/hm ²)	Litter biomass (g/m ²)
Y10	109°28 E, 36°27 N	1692±0.24	6.40±0.42	7.00±0.34	190.05±7.92	418.39±11.58
Y15	109°33' E, 36°27' N					7.33±0.
Y40	110°22 E, 38°47 N	-	-	-	-	-

Note: ATH, average tree height; DBH, diameter at breast height. Values of ATH and DBH represent the mean values of randomly selected 20 RP trees with DBH greater than 2.00 cm. The leaves of RP were collected using five collectors (1 m × 1 m) and dried to constant weight at 65°C to obtain the litter biomass. Y10, Y15, Y25 and Y40 indicate RP plantations with stand ages of 10, 15, 25 and 40 a, respectively. Mean±SD.

Table S2 Relationships between soil properties and microbial alpha diversity indices

Soil property	Bacterial alpha diversity	Fungal alpha diversity
	Shannon	Chao
Moisture	0.728**	0.556*
SOC	-0.754**	-0.269
TN	-0.760**	-0.255
NO ₃ ⁻ -N	-0.640**	-0.084
NH ₄ ⁺ -N	-0.658**	-0.265
AP	-0.421	-0.382
pH	0.016	-0.378

*Note: SM, soil moisture; SOC, soil organic carbon; TN, total nitrogen; NO₃⁻-N, nitrate nitrogen; NH₄⁺-N, ammonium nitrogen; AP, available phosphorus. **, $P < 0.01$ level, *, $P < 0.05$ level.

Table S3 Significance tests of the variations of fungal communities between Y10 and Y15 using the Adonis test

Group factor	Sums of square	Mean of square
Residual	-	-
Total	-	-

Note: Significant level is $P < 0.05$.

Table S4 Forward selection results of the redundancy analysis (RDA) for microbial communities and soil properties

Microbe	Soil property	Explanation (%)	Contribution (%)
Bacteria	TN	-	-
Bacteria	NO ₃ ⁻ -N	-	-
Bacteria	SM	-	-
Fungi	SOC	-	-
Fungi	NO ₃ ⁻ -N	-	-

Note: #, soil property was significantly affected by microbial communities. Significant level is $P < 0.05$.

Fig. S1 Principal component analysis (PCA) of soil properties at four *Robinia pseudoacacia* (RP) plantations with different stand ages. PCA and visualization were performed using CANOCO v5.0. SM, soil moisture; SOC, soil organic carbon; TN, total nitrogen; NO₃⁻-N, nitrate nitrogen; NH₄⁺-N, ammonium nitrogen; AP, available phosphorous.

Fig. S2 Venn analyses of the operational taxonomic units (OTUs) of soil bacterial (a) and fungal (b) communities at four RP plantations with different stand ages. Venn analyses and visualization were performed using the “VennDiagram” package in R 3.4.1.

Fig. S3 Time-decay relationships of fungal and bacterial communities at RP plantations with 40 a of restoration. The slope means the temporal turnover rate of microbial communities across restoration time.

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv — Machine translation. Verify with original.