

GPC3 and PEG10 polypeptides associated with placental gp96 elicit specific T cell immunity against hepatocellular carcinoma

Authors: Qin, Lijuan, Wang, Jiuru, Cheng, Fang, Zhang, Han, Zheng, Huaguo, Liu, Yongai, Liang, Zhentao, Wang, Baifeng, Wang, Haoyu, Ju, Ying, (Please provide the Chinese academic text with tags for translation.), Meng, Songdong, Ju, Ying, Tian, Huaqin, Meng, Songdong

Date: 2023-05-22T00:00:00+00:00

Abstract

Songdong MENG conceived of the project. Huaqin TIAN, Ying JU supervised the project. Lijuan QIN, Fang CHENG, and Jiuru WANG performed the experiments and analyzed the data. Lijuan QIN edited the original draft. Yongai LIU, Han ZHANG, Huaguo ZHENG, Baifeng WANG, Zhentao LIANG, and Haoyu WANG assisted with performing the experiments. Songdong MENG, Lijuan QIN, and Ying JU wrote the manuscript.

Full Text

Preamble

GPC3 and PEG10 Polypeptides Associated with Placental gp96 Elicit Specific T Cell Immunity Against Hepatocellular Carcinoma

Lijuan QIN^{1,2,#}, Jiuru WANG^{1,2,#}, Fang CHENG^{1,2,#}, Han ZHANG^{1,2}, Huaguo ZHENG^{1,2}, Yongai LIU^{1,2}, Zhentao LIANG^{1,2}, Baifeng WANG^{1,2}, Haoyu WANG^{1,2}, Ying JU¹,, *Huaqin TIAN*³,, & Songdong MENG^{1,2,*}

¹ CAS Key Laboratory of Pathogen Microbiology and Immunology, Institute of Microbiology, Chinese Academy of Sciences (CAS), Beijing, China

² University of Chinese Academy of Sciences, Beijing, China

³ Foshan Hospital of Traditional Chinese Medicine, Guangdong, China

These authors contributed equally to this work.

- Corresponding authors:
Songdong Meng (E-mail: mengsd@im.ac.cn)
Ying Ju (E-mail: juying7527@163.com)
Huaqin Tian (E-mail: 13929969262@139.com)

Abstract

The placenta and tumors can exhibit shared expression profiles of proto-oncogenes. The mechanism by which placenta-derived heat shock protein gp96 induces prophylactic and therapeutic T cell responses against cancers, including hepatocellular carcinoma (HCC), remains unknown. Here, we identified associated polypeptides from human placental gp96 using matrix-assisted laser desorption/ionization-time-of-flight mass spectrometry and analyzed the resulting proteins through disease enrichment analysis. We found that placental gp96 binds numerous peptides derived from 73 proteins that are enriched in multiple cancer types. Epitope-harboring peptides from GPC3 and PEG10 were the major antigens mediating anti-HCC T cell immunity. Molecular docking analysis revealed that GPC3- and PEG10-derived peptides, primarily obtained from the cytotrophoblast layer of mature placenta, bind to the luminal channel and client-bound domain of the gp96 dimer. Immunization with bone marrow-derived dendritic cells pulsed with recombinant gp96-GPC3 or recombinant gp96-PEG10 peptide complexes induced specific T cell responses, and T cell transfusion led to pronounced growth inhibition of HCC tumors in nude mice. We demonstrated that the chaperone gp96 can capture antigenic peptides as an efficient approach for defining tumor rejection oncoantigens in the placenta and provide a basis for developing GPC3- and PEG10 polypeptide-based vaccines against HCC. This study provides insight into the underlying mechanism of the antitumor response mediated by embryonic antigens from fetal tissues and will incite further studies to identify potential tumor rejection antigens from the placenta.

Keywords: Placental gp96, hepatocellular carcinoma, epitopes, GPC3, PEG10

1. Introduction

Aberrant expression, mutations, and somatic alterations in certain oncogenes and/or tumor suppressor genes have been detected in various human malignancies. The placenta is a heterogeneous, invasive tumor-like organ that invades the uterus to enable exchange of gases, nutrients, hormones, and other molecules between maternal and fetal blood while providing a barrier that protects the fetus from maternal immune attack. Placenta-specific trophoblast cells exhibit various similarities with cancer cells, including rapid proliferation capacity, epithelial-mesenchymal transition and invasion, cell-cell fusion, induction of angiogenesis, and immune escape. Moreover, the mechanisms regulating trophoblastic and malignant cell functions are similar. Ferretti et al. reported that

activation of the phosphatidylinositol 3-kinase/AKT axis represents a central feature of signaling pathways governing proliferative, migratory, and invasive processes in both trophoblasts and cancer cells. A recent study revealed that human placental tissues contain extensive somatic mutations; the mutation rate per trophoblast cluster is similar to that in childhood cancers, primarily induced by mutations in the uterus. Furthermore, the decidual microenvironment that regulates trophoblasts shows many similarities with the tumor microenvironment, such as supporting tumor cell survival through immune cells, oxygen levels, and energy metabolism.

The heat shock protein (HSP) gp96, the most abundant chaperone in the endoplasmic reticulum (ER), can modulate innate and adaptive antitumor immune responses. As an essential master chaperone, gp96 is involved in trafficking of Toll-like receptors and alpha integrins to regulate innate immunity as well as T- and B-cell development. Regarding antigen-specific immunity, tumor-derived gp96-peptide complexes or cell-based gp96-Ig-secreting vaccines elicited specific T cell immunity against parent tumors in both rodent models and clinical trials. gp96 has been proposed to exert multiple roles in T-cell activation, with its unique ability to bind a variety of tumor-associated antigens (TAAs) and disease-associated antigens (DAAs) peptides for antigen presentation to major histocompatibility complex (MHC) molecules playing a key role in this process. We previously identified a gp96-associated peptide in human liver tumor tissues; this peptide homolog was found to bind to human leukocyte antigen (HLA)-A11 molecules. Additionally, we and others showed that cellular gp96 interacts with MHC-I and the transporter associated with antigen processing, and that gp96 and calreticulin in the ER constitute a relay line for transferring associated peptides to MHC-I molecules for T-cell recognition. These studies demonstrated the ability of gp96 to capture and deliver antigens to MHC molecules to generate antigen-specific T-cell responses.

Because of the similarity in antigen expression between cancer and embryonic tissues, gp96 extracted from the placenta may bind carcinoembryonic antigens and/or proto-oncogene antigens. We previously showed that placenta-derived gp96 (Pgp96) can serve as a prophylactic and therapeutic vaccine against several cancers, including transplantable melanoma and breast tumors in mice. Moreover, placenta-derived peptides bound to gp96 pulsed dendritic cells (DCs) and generated specific CD8⁺ T cell responses against melanoma and Lewis lung cancer in mouse models. A clinical trial was recently initiated to test the safety and therapeutic efficacy of placental gp96-peptide complexes against non-small cell lung cancer and hepatocellular carcinoma (HCC) (No. ChiCTR2100052023). In this study, we isolated gp96-associated peptides from the human placenta, identified T cell epitopes through peptidome analysis using mass spectrometry (MS), and explored the mechanism of placental antigen-mediated immunotherapeutic effects against HCC. These results may facilitate the design of vaccines and T cell immunotherapies against cancer.

2. Materials and Methods

Detection and Identification of Peptides Bound to Human Placental gp96 by Mass Spectrometry

Bulk polypeptide fractions were isolated from placental gp96 samples using an acid treatment release method. Prior to mass spectrometry (MS) analysis, polypeptide samples were freeze-dried with a centrifuge concentrator, redissolved in 100 μ L water, sonicated for 5 min, and centrifuged at $14,000 \times g$ to obtain the supernatant. The supernatant was enriched using a polypeptide chip (L121 mesoporous silicon wafer) and detected using matrix-assisted laser desorption/ionization (MALDI).

To identify peptide sequences, the eluents of all samples were combined, lyophilized, and redissolved in 15 μ L of 0.1% formic acid. Peptide sequences were identified using a nanoLC-Q EXACTIVE (Thermo Fisher Scientific). Analytical separation was performed using gradients of H₂O/formic acid 100%/0.1% (solvent A) and CH₃CN/formic acid 100%/0.1% (solvent B). The gradient was run as follows: 0 min 4% B, then to 8% B at 8 min, 22% B at 58 min, 32% B at 70 min, 90% B at 71 min, 90% B at 78 min at a flow rate of 300 nL/min. The tandem MS scan range was 300-1,600 m/z and dynamic exclusion time was 40 s.

The peptide mixture was analyzed using liquid chromatography-tandem mass spectrometry (LC-MS/MS). The SEQUEST HT search engine of Thermo Proteome Discoverer (1.4.0.288) was used to search for and identify proteins in the Uniprot-proteome-human (update-20160226) database. The search parameters were as follows: no enzyme was required. The initial maximum allowed mass deviation of the precursor ion was set to 10 ppm, and the maximum fragment ion mass deviation was set to 20 mDa. Methionine oxidation was used as a variable modification. The filter parameter of the results was that the percolator filtered the spectrogram. The delta Cn was <0.1 , and the false discovery rate was set to 1%.

Additional Methods

Details regarding mice and cell culture, preparation of placental gp96 and recombinant gp96 proteins, peptide synthesis and epitope prediction, preparation and immunization strategy of pulsed-BMDCs vaccines, animal tumor models and adoptive therapy, enzyme-linked immunosorbent spot (ELISPOT) analysis, in vitro tumor cell killing assays, T2 binding assays, tetramer staining and flow cytometry, molecular docking analysis, quantitative real-time PCR, Western blot assays, and statistical analysis are provided in Supplementary Materials and Methods.

3. Results

Identification of Placental gp96-Bound Peptides and Cytotoxicity Testing

Placenta-derived gp96 (Pgp96) binds intracellular antigenic peptides, including those from proto-oncogenes, that may be shared between the placenta and HCC tumors. Therefore, gp96-peptide complexes were isolated from human placenta. The Pgp96-associated peptides were extracted, freeze-dried, enriched using L121 mesoporous silicon wafers, and subjected to matrix-assisted laser desorption/ionization-time-of-flight (MALDI-TOF) analysis. Peptide sequences were detected using Q-Exactive mass spectrometry (MS). We obtained 1,130 and 1,107 peptides with high confidence levels in two individual experiments. The resulting tandem MS data of the common peptides were searched against the Uniprot-proteome-human (update-20160226) database using the SEQUEST HT search engine of Thermo Proteome Discoverer (1.4.0.288). We identified 73 common proteins in both experiments. Disease enrichment analysis using data from the Disease Ontology database demonstrated that these protein genes were enriched in multiple cancer types, including liver, breast, and pancreatic cancers [Figure 1: see original paper], suggesting that the peptide repertoire from placental gp96 covers a large quantity of tumor antigens.

We then examined whether human placental gp96-bound peptides induce tumor-specific T cell responses. HLA-A2.1/Kb transgenic mice immunized with Pgp96-pulsed bone marrow-derived dendritic cells (BMDCs) exhibited a significant increase in HepG2-specific T cells, an HLA-A2-positive HCC cell line [Figure 1: see original paper], and an enhanced cytotoxic effect compared to mice immunized with recombinant gp96-pulsed or unpulsed BMDCs [Figure 1: see original paper]. Similar results were observed with the human pancreatic adenocarcinoma cell line PANC-1 and lung adenocarcinoma cell line A549. In contrast, tumor antigen-free recombinant gp96 expressed in a baculovirus system did not induce a tumor-specific T cell response, indicating that the antitumor T cell response induced by Pgp96 was attributed to the associated tumor antigens.

We further dissected the HCC antigens from the Pgp96-bound peptides. Through Venn analysis using the Gene Expression Omnibus public transcriptome dataset of HCC (GSE101675) and placenta (GSE1133), we found that two proteins, GPC3 and PEG10, were present in all three datasets [Figure 1: see original paper]. Another 25 common genes were detected in three other different datasets for HCC (GSE101685, GSE105130, and GSE121248), which also included GPC3 and PEG10. We analyzed the sources of GPC3 and PEG10 in the placenta by tracing the expression of these two genes in single-cell sequencing data of sorted placental cells from first- and second-trimester human placentas. The 1,567 cells were divided into 13 groups, with each group containing specific high-expression genes (Supplementary Figure S1A and S1B), suggesting that the overall cell clustering was reliable. Next, we selected marker genes of different cells based on results reported by Tsang et al. and

re-annotated the data on the above early placental single cells. As shown in Supplemental Figure S1C, re-annotated cells in the 13 groups included nine cell types. By searching for cell populations highly expressing GPC3 and PEG10 using the FeaturePlot() function of Seurat, GPC3 was found to be highly expressed in stromal and syncytiotrophoblast cells, whereas PEG10 was highly expressed in stromal and cytotrophoblast cells (Figure S1D). The GSE173193 dataset provides 10× Genomics single-cell transcriptome sequencing data for normally delivered placentas. As shown in [Figure 1: see original paper] (left panel), the cells were divided into 19 groups after t-stochastic neighborhood embedding analysis. Analysis of different placental marker genes showed that COL1A1, a marker gene of stromal cells, was not significantly expressed in normal parturient placentas, whereas the GPC3 and PEG10 genes were highly expressed in villous cytotrophoblast cells annotated by the PARP1 gene (Figure S1B-D, S2, and [Figure 1: see original paper]). The same expression pattern was obtained for GPC3 and PEG10 through annotation of the villous cytotrophoblast layer using HLA-G- and KRT7+ double genes in first- and second-trimester human placentas (Supplementary Figure S1B). These results indicate that GPC3- and PEG10-derived peptides bound to gp96 in the mature placenta were mainly derived from the cytotrophoblast layer [Figure 1: see original paper], right panel.

Furthermore, we analyzed single-cell RNA sequencing data from HCC tumor specimens based on a study by Sun et al. and found that GFP, a marker gene of HCC, was co-expressed with GPC3 and PEG10 in the same cell [Figure 1: see original paper]. These results indicate that the expression of GPC3 and PEG10 is HCC-specific.

Analysis of Immunogenicity of Placental gp96-Bound GPC3 and PEG10 Peptides

Four polypeptides from GPC3 and six polypeptides from PEG10 bound to placental gp96 were identified. Bioinformatics analysis revealed that these peptides contained abundant HLA-A-restricted epitopes. Twelve HLA-A0201-restricted epitopes of PEG10 and eight epitopes of GPC3 were synthesized. As shown in a T2 binding assay [Figure 2: see original paper], three epitopes of GPC3 and four epitopes of PEG10 showed high affinity for HLA-A0201 molecules. Subsequently, HLA-A2.1/Kb transgenic mice were subcutaneously vaccinated twice with BMDCs pulsed with a mixture of these three GPC3 peptides or a mixture of the four PEG10 peptides incorporated with recombinant gp96, and peptide-specific CD8+ T cells were quantified by analyzing IFN- γ + CD8+ T cells [Figure 2: see original paper]. As shown in [Figure 2: see original paper], immunized mice showed an activated cytotoxic T lymphocyte (CTL) response against epitopes GPC3^{152–160} and PEG10^{229–237}. The immunogenicity of epitope peptides GPC3^{152–160} and PEG10^{229–237} was confirmed in tetramer [Figure 2: see original paper] and enzyme-linked immunosorbent spot (ELISPOT) [Figure 2: see original paper] assays. Importantly, mice

immunized with Pgp96-pulsed BMDCs also exhibited specific CTL responses to GPC3^{152–160} and PEG10^{229–237}, validating that placental gp96 was associated with epitope peptides from GPC3 and PEG10.

We further modeled the structures of the two defined epitopes from GPC3 and PEG10 bound to HLA-A2 using molecular docking analysis. The results showed that both epitopes bind to the active site of the α -subunit of HLA-A2 based on visible hydrogen bonds and strong electrostatic interactions. Epitope GPC3^{152–160} (KVFGNFPKL) interacts with E63, T73, Y99, Q155, Y159, T163, and W167 via hydrogen bonding in the active cavity [Figure 3: see original paper]. Epitope PEG10^{229–237} (ALIGQCIHI) interacts with D77, T80, Y84, R97, W147, and Y159 via hydrogen bonding in the cavity [Figure 3: see original paper]. The two epitopes showed low binding energies of -8.0 and -8.8 kcal/mol with HLA-A2, indicating highly stable binding.

We also performed docking modeling of the gp96-dimer with the GPC3 polypeptide containing epitope GPC3^{152–160} and PEG10 polypeptide containing epitope PEG10^{229–237} using ZDOCK. As shown in [Figure 3: see original paper] and [Figure 3: see original paper], the red region represents a lumen formed by two loops of the M-domain (residues 394–407) and a client protein-binding domain (residues 652–678) of the gp96 dimer. The bound polypeptides adopted α -helices and/or β -turns as motifs for gp96 dimer recognition. GPC3 and PEG10 polypeptides bind near the lumen in different manners. The GPC3 polypeptide was inserted into the lumen in an α -helical form [Figure 3: see original paper], whereas the PEG10 polypeptide bound to the pre-luminal client protein-binding domain [Figure 3: see original paper]. As shown in [Figure 3: see original paper], [Figure 3: see original paper], and Supplementary Table S1, the GPC3 polypeptide can form hydrogen bond interactions with the R395, Y678, and D672 residues of gp96, and the PEG10 polypeptide can form hydrogen bond interactions with the D672, T675, and S674 residues to achieve stable binding. The binding, hydrolysis, and release of ATP can lead to transitions between different conformations of the gp96-dimer (Figure S3). Conformational changes in the polypeptide-binding cavity formed by the two loops of the M domain and client protein-binding domain, binding state, and binding energy of the gp96-dimer with GPC3 or PEG10 polypeptide may also vary [Figure 3: see original paper] and [Figure 3: see original paper]. Therefore, the process of ATP hydrolysis by the gp96-dimer may affect the binding and release of peptides through conformational changes.

GPC3 and PEG10 Epitopes Within Pgp96-Bound Peptides Induce Anti-HCC Immune Response

Using the Cancer Cell Line Encyclopedia (CCLE) database, we found that the expression levels of GPC3 and PEG10 were high in HepG2 cells and low in SK-Hep-1 cells (Figure S4). Transcriptome analysis, quantitative reverse transcription-polymerase chain reaction (PCR), and Western blot assay showed that the expression levels of GPC3 and PEG10 were much higher in HepG2

cells than in SK-Hep-1 cells [Figure 4: see original paper], [Figure 4: see original paper], and [Figure 4: see original paper]. Twenty-one overlapping genes were detected in intersection analysis between protein genes related to placental gp96-bound peptides and protein genes that were highly expressed in HepG2 cells compared to SK-Hep-1 cells, including GPC3 and PEG10 [Figure 4: see original paper]. As shown in [Figure 4: see original paper], T cells induced by Pgp96, GPC3^{152–160}, or PEG10^{229–237} epitope exhibited much higher cytotoxicity against GPC3- and PEG10-high-expressing HepG2 cells than against GPC3- and PEG10-low-expressing SK-Hep-1 cells.

We further investigated the ability of the GPC3 and PEG10 epitopes to inhibit HCC in vivo [Figure 4: see original paper]. Transfer of epitope-induced CD8+ T cells resulted in decreased HCC tumor growth and weight in BALB/c nu mice compared to the effects of control CD8+ T cells [Figure 4: see original paper] and [Figure 4: see original paper]. Similar tumor inhibitory effects were observed in mice treated with placental gp96-induced CD8+ T cells.

Polypeptides from GPC3 and PEG10 Complexed with Recombinant gp96 Exhibit Anti-HCC Activity

As shown in [Figure 5: see original paper], similar to the above epitopes, these two GPC3 and PEG10 precursor peptides complexed with recombinant gp96 exhibited specific cytotoxicity for HepG2 cells but not for SK-Hep-1 cells. The ELISPOT assay revealed that GPC3 or PEG10 polypeptide-pulsed BMDC immunization resulted in an approximately 6- or 7-fold increase in GPC3- and PEG10-high-expressing HepG2 cell-specific T cells compared to control (both $P < 0.01$). In contrast, no obvious specific T cell response to GPC3- and PEG10-low-expressing SK-Hep-1 cells was detected, indicating that the peptide-induced anti-HCC response was GPC3- and PEG10-specific [Figure 5: see original paper].

We further assessed the in vivo therapeutic efficacy of GPC3 and PEG10 polypeptides in HepG2-bearing mice. As shown in [Figure 5: see original paper]-[Figure 5: see original paper], treatment with CD8+ T cells induced by GPC3 or PEG10 polypeptide showed a significant therapeutic effect against HCC compared to that in the control (all $P < 0.05$ or 0.01).

4. Discussion

In the current study, using MS-based techniques, we found that abundant tumor-associated antigens that were preferentially expressed in multiple tumor tissues bound to placental gp96, and a large portion of these antigens were primarily associated with HCC. We further established an MS-based immunopeptidomic approach for identifying tumor-specific antigens from the placenta and identified GPC3 and PEG10 as HCC antigens with high immunogenicity recognized by T cells [Figure 6: see original paper]. These results provide insight into the mechanism of the antitumor response mediated by embryonic antigens from

fetal tissues or stem cells.

The structure of gp96 contains four domains: N-terminal, charged linker, middle, and C-terminal. Numerous studies have shown that exogenous gp96 mediates the uptake and presentation of its chaperoned peptides by antigen-presenting cells to MHC-I molecules and activates specific CTLs in vivo. However, the structural interaction between gp96 and the associated peptides remains unclear. A recent study of the gp96:polypeptide interaction revealed that polypeptides associate with the luminal channel formed between two loops in the middle domain and client-binding domain of the gp96 dimer. In the present study, we showed that placental gp96 associates with epitope-harboring polypeptides derived from GPC3 and PEG10. The GPC3 and PEG10 polypeptides bind to the luminal channel and client-bound domain. Moreover, intramolecular conformational changes induced by ATP hydrolysis may regulate the association and release of polypeptides from gp96 [Figure 3: see original paper]. Thus, we further uncovered the structural basis of gp96 binding of cellular antigenic peptides and validated that the immunopeptidomic analysis of gp96-bound peptides is an effective approach for antigen capture and identification.

An association between placenta and tumor antigens has been widely reported. In this study, we found that placental gp96 was associated with epitope-harboring polypeptides from GPC3 and PEG10 that elicited anti-HCC T cell responses. We further analyzed the source of these antigens in the placenta and found that the GPC3 and PEG10 genes were significantly highly expressed in villous cytotrophoblast cells of normally delivered placentas [Figure 1: see original paper]. GPC3 is an oncofetal glycoprotein that binds to the cell membrane via a glycosylphosphatidylinositol (GPI) anchor. GPC3 is expressed in the placenta, numerous embryonic tissues, and various tumors such as HCC but not in healthy adult liver, ovarian carcinoma, and melanomas. GPC3 promotes the growth of HCC cells through Wnt/ β -catenin signaling and is emerging as a potential therapeutic target. PEG10, a paternally expressed imprinted gene that encodes a cytosolic protein, is primarily expressed in the placenta, and its expression levels are elevated in various cancers, including HCC. PEG10 enhances the proliferation and invasion of cancer cells. In HCC, PEG10 overexpression can improve cell invasion ability and decrease cell apoptosis mediated by SIAH1. Our current data, along with previous studies, clearly demonstrate the similarity of antigen expression patterns between the placenta and tumors and support the design of a new generation of proto-oncogene antigen-based cancer vaccines. Further studies are needed to define new placental antigens and determine their potential functions against cancers, including pancreatic cancer, breast carcinoma, and lung cancer.

In summary, we showed that using chaperone gp96 to capture antigenic peptides is an efficient approach for identifying tumor rejection antigens generated in the placenta. GPC3 and PEG10 antigen-pulsed DCs exhibit therapeutic effects against HCC. Large-scale, systematic studies of inherent oncofetal antigens from the human placenta may help reveal novel proto-oncogenes with therapeutic

potential against cancer.

Acknowledgments

We would like to thank Tong Zhao for FACS support.

Disclosure Statement: Funding Information

This work was supported by a grant from the National Key R&D Program of China (2022YFC2304203); the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB29040000); the National Natural Science Foundation of China (32070163); the Foshan “Summit Plan” for building high-level hospitals; and the Industrial Innovation Team Grant from the Foshan Industrial Technology Research Institute, Chinese Academy of Sciences.

Conflict of Interest

The authors declare no conflict of interest.

Ethics Statement

Approval of the research protocol by an Institutional Review Board: N/A.

Informed Consent: N/A.

Registry and the Registration No. of the study/trial: N/A.

Animal Studies

The treatment of animals complied with legal and ethical requirements. All animal studies were approved by the Research Ethics Committee of the Institute of Microbiology, Chinese Academy of Sciences (approval number PZIM-CAS2011001) and followed the ARRIVE guidelines.

Availability of Data

All data relevant to the study are available upon reasonable request.

References

1. Burrell RA, McGranahan N, Bartek J, Swanton C. The causes and consequences of genetic heterogeneity in cancer evolution. *Nature*. 2013;501(7467):338-45. doi:10.1038/nature12625
2. Lala PK, Nandi P, Hadi A, Halari C. A crossroad between placental and tumor biology: What have we learnt? *Placenta*. 2021;116:12-30. doi:10.1016/j.placenta.2021.03.003
3. Costanzo V, Bardelli A, Siena S, Abrignani S. Exploring the links between cancer and placenta development. *Open Biol*. 2018;8(6). doi:10.1098/rsob.180081
4. Pang H, Lei D, Guo Y, Yu Y, Liu T, Liu Y, et al. Three categories of similarities between the placenta and cancer that can aid cancer treatment: Cells, the microenvironment, and metabolites. *Front Oncol*. 2022;12:977618. doi:10.3389/fonc.2022.977618

5. Ferretti C, Bruni L, Dangles-Marie V, Pecking AP, Bellet D. Molecular circuits shared by placental and cancer cells, and their implications in the proliferative, invasive and migratory capacities of trophoblasts. *Hum Reprod Update*. 2007;13(2):121-41. doi:10.1093/humupd/dml048
6. Coorens THH, Oliver TRW, Sanghvi R, Sovio U, Cook E, Vento-Tormo R, et al. Inherent mosaicism and extensive mutation of human placentas. *Nature*. 2021;592(7852):80-5. doi:10.1038/s41586-021-03345-1
7. Krstic J, Deutsch A, Fuchs J, Gauster M, Gorsek Sparovec T, Hiden U, et al. (Dis)similarities between Decidual Tumor Microenvironment. *Biomedicines*. 2022;10(5). doi:10.3390/biomedicines10051065
8. Pugh KW, Alnaed M, Brackett CM, Blagg BSJ. The biology and inhibition of glucose-regulated protein 94/gp96. *Med Res Rev*. 2022;42(6):2007-24. doi:10.1002/med.21915
9. Staron M, Yang Y, Liu B, Li J, Shen Y, Zuniga-Pflucker JC, et al. gp96, an endoplasmic reticulum master chaperone for integrins and Toll-like receptors, selectively regulates early T and B lymphopoiesis. *Blood*. 2010;115(12):2380-90. doi:10.1182/blood-2009-07-233031
10. Bolhassani A, Rafati S. Heat-shock proteins as powerful weapons in vaccine development. *Expert Rev Vaccines*. 2008;7(8):1185-99. doi:10.1586/14760584.7.8.1185
11. Randazzo M, Terness P, Opelz G, Kleist C. Active-specific immunotherapy of human cancers with the heat shock protein Gp96-revisited. *J Cancer*. 2012;130(10):2219-31. doi:10.1002/jjc.27332
12. Oizumi S, Strbo N, Pahwa S, Deyev V, Podack ER. Molecular and cellular requirements for enhanced antigen cross-presentation to CD8 cytotoxic T lymphocytes. *Immunol*. 2007;179(4):2310-7. doi:10.4049/jimmunol.179.4.2310
13. Binder RJ, Blachere NE, Srivastava PK. Heat shock protein-chaperoned peptides but not free peptides introduced into the cytosol are presented efficiently by major histocompatibility complex I molecules. *J Biol Chem*. 2001;276(20):17163-71. doi:10.1074/jbc.M011547200
14. Udono H, Levey DL, Srivastava PK. Cellular requirements for tumor-specific immunity elicited by heat shock proteins: tumor rejection antigen gp96 primes CD8+ T cells in vivo. *Proc Natl Acad Sci U S A*. 1994;91(8):3077-81
15. Jacqueline C, Dracz M, Xue J, Binder RJ, Minden J, Finn O. LCVM infection generates tumor antigen-specific immunity and inhibits growth of nonviral tumors. *Oncoimmunology*. 2022;11(1):2029083. doi:10.1080/2162402X.2022.2029083

16. Meng SD, Gao T, Gao GF, Tien P. HBV-specific peptide associated with heat-shock protein gp96. *Lancet*. 2001;357(9255):528-9
17. Meng S-D, Song J, Rao Z, Tien P, Gao GF. Three-step purification of gp96 from human liver tumor tissues suitable for isolation of gp96-bound peptides. *J Immunol Methods*. 2002;264(1-2):29-35
18. Kropp LE, Garg M, Binder RJ. Ovalbumin-derived precursor peptides are transferred sequentially from gp96 and calreticulin to MHC class I in the endoplasmic reticulum. *J Immunol*. 2010;184(10):5619-27. doi:10.4049/jimmunol.0902368
19. Qin L, Liu Y, Xu Y, Li Y, Hu J, Ju Y, et al. MHC Class I Assembly Function and Intracellular Transport Routes for Hepatitis B Virus Antigen Cross-presentation by Heat Shock Protein gp96. *Infectious Diseases & Immunity*. 2022;2(3):183-92. doi:10.1097/id9.0000000000000058
20. Zhao B, Wang Y, Wu B, Liu S, Wu E, Fan H, et al. Placenta-derived gp96 as a multivalent prophylactic cancer vaccine. *Sci Rep*. 2013;3:1947. doi:10.1038/srep01947
21. Zheng H, Liu L, Zhang H, Kan F, Wang S, Li Y, et al. Dendritic cells pulsed with placental gp96 promote tumor-reactive immune responses. 2019;14(1):e0211490. doi:10.1371/journal.pone.0211490
22. Liu Y, Fan X, Wang R, Lu X, Dang YL, Wang H, et al. Single-cell RNA-seq reveals the diversity of trophoblast subtypes and patterns of differentiation in the human placenta. *Cell Res*. 2018;28(8):819-32. doi:10.1038/s41422-018-0066-y
23. Tsang JCH, Vong JSL, Ji L, Poon LCY, Jiang P, Lui KO, et al. Integrative single-cell and cell-free plasma RNA transcriptomics elucidates placental cellular dynamics. *Proc Natl Acad Sci U S A*. 2017;114(37):E7786-E95. doi:10.1073/pnas.1710470114
24. Sun Y, Wu L, Zhong Y, Zhou K, Hou Y, Wang Z, et al. Single-cell landscape of the ecosystem in early-relapse hepatocellular carcinoma. *Cell*. 2021;184(2):404-21.e16. doi:10.1016/j.cell.2020.11.041
25. Suto R, Srivastava PK. A mechanism for the specific immunogenicity of heat shock protein-chaperoned peptides. *Science*. 1995;269(5230):1585-8
26. Huck JD, Que NL, Hong F, Li Z, Gewirth DT. Structural and Functional Analysis of GRP94 in the Closed State Reveals an Essential Role for the Pre-N Domain and a Potential Client-Binding Site. *Cell Rep*. 2017;20(12):2800-9. doi:10.1016/j.celrep.2017.08.079
27. Dong W, Du J, Shen H, Gao D, Li Z, Wang G, et al. Administration of embryonic stem cells generates effective antitumor immunity in mice with minor and heavy tumor load. *Cancer Immunol Immunother*. 2010;59(11):1697-705. doi:10.1007/s00262-010-0899-9

28. Li W, Liu D, Chang W, Lu X, Wang YL, Wang H, et al. Role of IGF2BP3 in trophoblast cell invasion and migration. *Cell Death Dis.* 2014;5:e1025. doi:10.1038/cddis.2013.545
29. Haruyama Y, Kataoka H. Glypican-3 is a prognostic factor and an immunotherapeutic target in hepatocellular carcinoma. *World J Gastroenterol.* 2016;22(1):275-83. doi:10.3748/wjg.v22.i1.275
30. Maeda D, Ota S, Takazawa Y, Aburatani H, Nakagawa S, Yano T, et al. Glypican-3 expression in clear cell adenocarcinoma of the ovary. *Mod Pathol.* 2009;22(6):824-32. doi:10.1038/modpathol.2009.40
31. Nakatsura T, Nishimura Y. Usefulness of the novel oncofetal antigen glypican-3 for diagnosis of hepatocellular carcinoma and melanoma. *Bio-Drugs.* 2005;19(2):71-7
32. Capurro MI, Xiang Y-Y, Lobe C, Filmus J. Glypican-3 promotes the growth of hepatocellular carcinoma by stimulating canonical Wnt signaling. *Cancer Res.* 2005;65(14):6245-54
33. Ip WK, Lai PB, Wong NL, Sy SM, Beheshti B, Squire JA, et al. Identification of PEG10 as a progression-related biomarker for hepatocellular carcinoma. *Cancer Lett.* 2007;250(2):284-91. doi:10.1016/j.canlet.2006.10.012
34. Okabe H, Satoh S, Furukawa Y, Kato T, Hasegawa S, Nakajima Y, et al. Involvement of PEG10 in human hepatocarcinogenesis through interaction with SIAH1. *Cancer Res.* 2003;63(12):3043-8
35. Sun L, Zhang Y, Zhao B, Deng M, Liu J, Li X, et al. A new unconventional HLA-A2-restricted epitope from HBV core protein elicits antiviral cytotoxic T lymphocytes. *Protein Cell.* 2014;5(4):317-26
36. Liu W, Chen M, Li X, Zhao B, Hou J, Zheng H, et al. Interaction of Toll-Like Receptors with the Molecular Chaperone Gp96 Is Essential for Its Activation of Cytotoxic T Lymphocyte Response. *PLoS One.* 2016;11(5):e0155202. doi:10.1371/journal.pone.0155202
37. Inaba K, Inaba M, Romani N, Aya H, Deguchi M, Ikehara S, et al. Generation of large numbers of dendritic cells from mouse bone marrow cultures supplemented with granulocyte/macrophage colony-stimulating factor. *J Exp Med.* 1992;176(6):1693-702
38. Zhang M, Tang H, Guo Z, An H, Zhu X, Song W, et al. Splenic stroma drives mature dendritic cells to differentiate into regulatory dendritic cells. *Nat Immunol.* 2004;5(11):1124-33. doi:10.1038/ni1130
39. Zhou M, Xu D, Li X, Li H, Shan M, Tang J, et al. Screening and identification of severe acute respiratory syndrome-associated coronavirus-specific epitopes. *Immunol.* 2006;177(4):2138-45. doi:10.4049/jimmunol.177.4.2138
40. Morris GM, Huey R, Olson AJ. Using AutoDock for ligand-receptor docking. *Curr Protoc Bioinformatics.* 2008;Chapter 8:Unit 8.14.

doi:10.1002/0471250953.bi0814s24

Supporting Information

Supplementary Figure 1: Analysis and Annotation of Single-Cell Transcriptome Data of Human Early Placentas

Related to [Figure 1: see original paper]. (A) Heatmap showing characteristic genes of different cell populations. The t-stochastic neighborhood embedding (t-SNE) algorithm in the R package Seurat was used to group cells and draw heatmaps of cell-specific genes from different populations to test the reliability of grouping. (B) Expression of different marker genes. (C) Cell population mapping and annotation after t-SNE dimensionality reduction of 1,567 single cells. (D) GPC3 and PEG10 gene expression in different cell populations in single-cell transcriptome data. Darker colors indicate higher expression levels in cells. The cell population circled by the red frame expressed high levels of GPC3 and PEG10.

Supplementary Figure 2: t-Stochastic Neighborhood Embedding (t-SNE) Dimensionality Reduction and Gene Expression of Single-Cell Sequencing Data of Normal Delivery Placenta

Related to [Figure 1: see original paper]. (A) Expression map of different genes visualized using t-SNE dimensionality reduction; the red dashed cell group represents villous cytotrophoblast cells. (B) Expression of different marker genes visualized using the FeaturePlot() function in the Seurat package. Darker colors indicate higher expression levels in cells.

Supplementary Figure 3: Schematic Diagram of the Mechanism by Which ATP Hydrolysis by gp96 Leads to Conformational Change

Related to [Figure 3: see original paper]. In the “twisted V” dimer state, gp96's N-terminal domain can bind ATP/AMP-PNP (non-hydrolytic), leading to a conformational change in the N-terminal domain and 90° rotation around the N-terminal/middle domain interface, thus promoting N-terminal dimerization. At this point, a closed dimer binding ATP/AMP-PNP is formed to prepare for ATP hydrolysis. After ATP hydrolysis and ADP release, the transient N-terminal dimer decomposes and re-rotates around the N-terminal/middle domain interface to form a mobile N-terminal domain.

Supplementary Figure 4: Heatmap of Expression Levels of GPC3 and PEG10

Related to [Figure 4: see original paper]. The expression profile data of different hepatocellular carcinoma cell lines were downloaded from the Cancer Cell Line Encyclopedia cancer database and analyzed.

Supplementary Table 1: Results of Molecular Docking ZDOCK, Discovery Studio 2019

Receptor	Ligand	ZDOCK Score	Hydrogen Bond Interaction	Electrostatic Interaction	Hydrophobic Interaction
gp96-dimer	GPC3 ¹⁴⁹⁻¹⁹⁵	-	A:ARG395:NH1 C:GLU238:OE1A:ARG587:NH2 - C:ASP204:OD1 - C:ILE215:OA:ARG395:NH2 - C:GLU238:OE1A:ARG587:NH1 - C:ASP204:OD1		
gp96-dimer	PEG10 ²⁰¹⁻²⁴⁷	-	A:TYR678:OH C:VAL219A:TRP654 C:PHE208:OC:PRO212:CD - C:ILE215A:MET662 A:TYR678:OHA:TYR652:OH - C:ILE235A:ILE673 C:LEU202:OA:THR675:N - C:ILE237A:TYR652 C:ILE235:OA:THR675:OG1 - C:LEU227A:TRP654 C:ILE235:OA:TYR677:OH - C:ILE235A:TYR677 C:GLY209:OA:TYR678:OH - C:LEU210A:TYR678 C:ILE231:OC:SER234:OG - A:GLN578:OA:SER674:CB - C:GLN233:O		

Supplementary Materials and Methods**Mice**

HLA-A2.1/Kb (HLA-A2) transgenic mice were maintained in the laboratory. BALB/c-nu mice were purchased from SPF (Beijing, China). All animals were maintained in a specific pathogen-free animal facility and studied at 6–10 weeks of age. Animal experiments were approved by the Research Ethics Committee of the Institute of Microbiology, Chinese Academy of Sciences (approval number PZIMCAS2011001).

Cell Culture

HepG2 cells were kindly provided by Prof. X Ye (Institute of Microbiology, Center for Biosafety Mega-Science, Chinese Academy of Sciences, Beijing, China). PANC-1 cells were cultured in Dulbecco's modified Eagle's medium (HyClone, Logan, UT, USA) supplemented with 10% fetal bovine serum (FBS). SK-Hep-1, A549 cells, and HepG2 cells were cultured in high-glucose Dulbecco's modified Eagle's medium supplemented with 10% FBS. All culture media contained 100 U/mL penicillin and 100 g/mL streptomycin.

Preparation of Placental gp96 and Recombinant gp96 Proteins

Placental gp96 proteins (Pgp96) of human origin were extracted as previously described. Briefly, after grinding placental tissue, the supernatant was collected and subjected to ammonium sulfate precipitation to obtain the sediment. The dissolved precipitate was purified using ConA-Sepharose affinity chromatography, followed by anion exchange chromatography. Endotoxin levels were determined using the Limulus Amebocyte Lysate assay (<1 EU/mg) (BioWhittaker, Walkersville, MD, USA).

Soluble recombinant human gp96 (Rgp96) was isolated as previously described. Briefly, recombinant human heat shock protein gp96 constructs were subcloned into the pFastBac1 vector and expressed using the Bac-to-Bac Baculovirus expression system. After filtration and concentration, the supernatant was collected and purified using a HiTrap Q column (GE Healthcare, Little Chalfont, UK). After desalting and concentration, protein purity was greater than 95% as determined by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE).

Peptide Synthesis and Epitope Prediction

The 8- to 10-mer epitopes were predicted using long peptides from two websites: SYFPEITHI (<http://www.syfpeithi.de/>) and the Immune Epitope Database Analysis Resource (<http://tools.iedb.org/main/tcell/>). Peptides with scores >20 (for SYFPEITHI) or percentile rank <0.5 typically have high MHC affinity. All peptides were synthesized by GenScript (Nanjing, China) and were >95% pure.

Preparation and Immunization Strategy of Pulsed-BMDCs Vaccines

Mouse bone marrow-derived DCs (BMDCs) were obtained as previously described. Briefly, bone marrow cells were harvested from the femur and tibia of 5-7-week-old mice, and red blood cells were depleted using red blood cell lysis buffer. Bone marrow cell suspensions were cultured in complete RPMI medium containing 10% heat-inactivated FBS, 20 ng/mL granulocyte-macrophage colony-stimulating factor (GM-CSF), 10 ng/mL interleukin-4 (IL-4), and 50 M 2-mercaptoethanol (Sigma-Aldrich, St. Louis, MO, USA). All cytokines were purchased from PeproTech (Rocky Hill, NJ, USA). After 7 days of culture, nonadherent BMDCs were generated and collected, then randomly divided into different groups according to experimental design: Pgp96 group (BMDCs incubated with 100 g/mL placental gp96); Rgp96 group (BMDCs incubated with 100 g/mL recombinant gp96); GPC3¹⁵²⁻¹⁶⁰ group (BMDCs treated

with 100 g/mL recombinant gp96 and 10 g/mL GPC3^{152–160}); PEG10^{229–237} group (BMDCs treated with 100 g/mL recombinant gp96 and 10 g/mL PEG10^{229–237}); epitope group (BMDCs treated with 100 g/mL recombinant gp96 plus 5 g/mL GPC3^{152–160} and 5 g/mL PEG10^{229–237}); positive control (PC) group (BMDCs loaded with 50 g/mL tumor cell lysate); DC group (BMDCs alone). To prepare gp96-peptide complexes, gp96 and peptide were mixed and incubated for 10 min at 50°C, followed by incubation for 30 min at room temperature. All BMDCs were then cultured in the presence of 1 g/mL lipopolysaccharide (LPS) for 24 h at 37°C. After incubation, non-adherent mature BMDCs were collected and washed three times with PBS. BMDCs were administered subcutaneously into HLA-A2 female mice on days 0 and 7 (5×10^5 cells in 100 μ L PBS/mouse). On day 5 after the last immunization, mice were euthanized and splenocytes were isolated for analysis by tetramer staining, ELISPOT assay, and in vitro tumor killing assay. The same protocol was used in tumor inhibition experiments with polypeptide-induced T cells.

Animal Tumor Model and Adoptive Therapy

BALB/c nude mice, approximately 4–6 weeks of age, were subcutaneously inoculated with 2×10^6 HepG2 tumor cells in 100 μ L PBS in the right flank. Tumor growth was monitored with calipers every other day, and tumor volume was calculated using the formula: volume = length \times width² \times 0.5. At four days after tumor cell inoculation, tumor-bearing mice were randomly divided into five groups and immunized intravenously in the tail with $3\text{--}4 \times 10^6$ T cells isolated from the spleens of Pgp96 group mice, Rgp96 group mice, epitope or peptide group mice, and PC group mice four times at 3- or 4-day intervals. Adoptive T cells were enriched using a CD8+ T cell Isolation Kit (Miltenyi Biotec, Gladbach Bergisch, Germany). After treatment, mice were euthanized if tumor depth exceeded 15 mm.

Enzyme-Linked Immunosorbent Spot (ELISPOT) Analysis

IFN- γ ELISPOT assays were performed according to the manufacturer's instructions (Mabtech, Mariemont, OH). Briefly, isolated mouse splenocytes were added to wells of 96-well polyvinylidene difluoride plates (BD-Pharmingen, San Diego, CA) in 4–6 replicates and subjected to different stimuli. Typical groups included cells with no stimulant, cells with antigens (tumor cell lysate or peptides), and cells treated with anti-CD3 (2 g/mL) and anti-CD28 (1 g/mL) (PC) and incubated at 37°C for indicated time periods. Spot-forming cells (SFCs) were counted and analyzed using an Immunospot S5 Versa Analyzer (Cellular Technology Limited, Shaker Heights, OH, USA).

In Vitro Tumor Cell Killing Assay

Splenocytes were isolated from immunized mice on day 5 after immunization, then co-cultured with BMDCs (at a 1:20 ratio) that had been incubated with the same treatments as immunization. Each well also contained 20 IU/mL recombinant murine IL-2 (PeproTech). After 5 days of culture, splenic effector cells were collected as effector cells. Target cells (HepG2 or SK-Hep-1) were labeled with 2.5 μ M 5-(6)-carboxy-fluorescein succinimidyl ester (CFSE) dye

and washed with PBS. CFSE-labeled target cells were then seeded in 100 μ L complete RPMI 1640 medium into 96-well round-bottom plates, and effector cells in 100 μ L complete RPMI 1640 medium were added to CFSE-labeled target cells at varying ratios for 4–6 h at 37°C. Dead cells were labeled with 7-AAD, and cytotoxicity of splenocytes to tumor cells was detected using flow cytometry.

T2 Binding Assay

T2 cells were incubated with 1 μ M human β 2m protein (Sigma–Aldrich) and 50 μ M polypeptide and cultured overnight in an incubator. Cells were collected and stained with FITC-labeled HLA-A2 antibody and detected using flow cytometry. The fluorescence index (FI) was calculated as previously described; a fluorescence index ≥ 1 was considered to indicate a high-affinity candidate epitope.

Tetramer Staining and Flow Cytometry

Tetramer antibody was prepared according to the instructions of the QuickSwitch™ Quant HLA-A*02:01 Tetramer Kit APC (MBL, Nagoya, Japan). Specific operations were as follows: First, lyophilized peptide was dissolved in dimethyl sulfoxide to a final concentration of 10 mg/mL. Second, 50 μ L of QuickSwitch™ Tetramer was mixed with 1 μ L peptide in a microtube, followed by addition of 1 μ L Peptide Exchange Factor. These steps were repeated for other peptides. Finally, the mixture was incubated for at least 4 h at room temperature in the dark. After reaction, tetramer antibody was stored at 4°C protected from light until use. Peptide exchange was quantified in a Flow Cytometric Sandwich immunoassay according to the manufacturer's instructions and analyzed using a Calibur flow cytometer (BD Biosciences, USA) and FlowJo Software (Tree Star Inc.). The exchange efficiency of the peptide was $91.9 \pm 1.8\%$, indicating successful tetramer antibody preparation.

Splenocytes obtained from immunized mice were stimulated with 2 μ g/mL peptide for 7 days at 37°C and 5% CO₂. Cells were harvested and stained with percp-conjugated anti-mouse CD8, FITC-conjugated anti-mouse CD3, and APC-conjugated peptide tetramer antibody. Data were acquired using a Fortessa flow cytometer (BD Biosciences, USA).

Molecular Docking Analysis

We used AutoDockVina 1.2.2, a computerized protein-ligand docking software, to assess binding energy and interaction patterns between candidate epitopes and HLA-A2 protein. The three-dimensional structures of proteins GPC3 (PDB ID: AF-P51654-F1), PEG10 (PDB ID: 7LGA, resolution 1.9 Å), and HLA-A2 (PDB ID: 3HLA, resolution 2.6 Å) were downloaded from the PDB (<http://www.rcsb.org/>). Molecular structures of the GPC3^{152–160} and PEG10^{229–237} epitopes were extracted from them. We first converted all protein and molecular files to PDBQT format, removed all water molecules, and added polar hydrogen atoms. The grid box was centered to cover the domain of each protein and accommodate free molecular motion. The docking pocket was set as a 30 \times 30 \times 30 Å square pocket with a grid distance of 0.05 nm for docking. Molecular docking results were analyzed and models were visualized using PyMol 2.5 and LigPlot+ v.2.2.

AlphaFold2 2.2.4 was used to predict the structure of the GPC3 polypeptide and PEG10 polypeptide bound to gp96 identified using MS. The amino acid sequence of human gp96 was downloaded from NCBI, and the dimer structure was predicted using AlphaFold2-multimer 2.2.4. The ranked_0 structure file was selected as the final simulation structure of the gp96-dimer protein by comparing the five result files. ZDOCK version 3.0.2 was used to predict the complex model of receptor and ligand via rigid docking. Interactions between gp96-dimer with GPC3 polypeptide or with PEG10 polypeptide were analyzed using Discovery Studio 2019. Final results were visualized and analyzed using PyMol 2.5 and LigPlot+ v.2.2.

We downloaded the gp96 dimer structure in different conformations from the RCSB PDB database to analyze its interaction with GPC3 or PEG10 polypeptides under different torsion dimer states. The Gp96-NMC/AMP-PNP-bound closed dimer coordinates were obtained from PDB ID 5ULS, GP96-NMC/AMP-PNP-bound “twisted V” dimer coordinates were obtained from PDB ID 2O1U, and GP96-NMC/ADP-bound “twisted V” dimer coordinates were obtained from PDB ID 2O1V. ZDOCK version 3.0.2 was used to predict the complex model of receptor and ligand via rigid docking. The top 10 ZDOCK scores were analyzed.

Quantitative Real-Time PCR

Total RNA was extracted from HepG2 or SK-Hep-1 cells using TRIzol reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. RNA was reverse-transcribed using PrimeScript™ RT Master Mix to generate cDNA (TaKaRa, Shiga, Japan). cDNA was analyzed by real-time PCR using a Rotor-Gene Q (Qiagen, Hilden, Germany). Primers used for amplification were as follows: GPC3 (5'-TGGAGAACGTAAGTCTGGTC-3' forward, 5'-TCTTCTCAGTTTCAGTGGTGG-3' reverse); PEG10 (5'-GAGAACAGCGGAGAAGGTCC-3' forward; 5'-CAAAACCCGCTTATTTACGC-3' reverse); GAPDH (5'-GGAGCGAGATCCCTCCAAAAT-3' forward; 5'-GGCTGTTGTCATACTTCTCATGG-3' reverse). These primers were used with TB Green Premix Ex Taq (TaKaRa). Quantitative PCR conditions were as follows: 95°C for 30 s, followed by 40 cycles of 95°C for 5 s and 60°C for 45 s. Amplification of specific transcripts was confirmed from melting curve profiles generated at the end of the PCR program. Expression levels of target genes were normalized to GAPDH and calculated using the comparative cycle threshold (CT) method ($2^{-\Delta\Delta CT}$).

Western Blot Assay

Western blot analysis was performed as previously described. Antibodies against GPC3 (Bioworld Technology, catalog no. BS7410) and PEG10 (Bioworld Technology, catalog no. MB0036) were used.

Transcriptome Analysis

Total RNA was prepared from HepG2 or SK-Hep-1 cells dissolved in TRIzol reagent according to the manufacturer's instructions. Transcriptome analysis was performed by Novogene Tianjin (Tianjin, China).

Common Database Analysis

All public databases used in this study have been published and can be obtained from the Gene Expression Omnibus database of NCBI, including the human tissue-related transcriptome database GSE1133; HCC-related transcriptome databases GSE101685, GSE105130, and GSE121248; and placental single-cell transcriptome database GSE89497.

Statistical Analysis

All data are expressed as mean \pm SD. Statistical variance between groups was analyzed using unpaired two-tailed t-test and one-way ANOVA using GraphPad Prism 8 software (GraphPad Software, USA). Variance between Kaplan-Meier curves was compared using log-rank test. Differences were considered statistically significant at $P < 0.05$. * $P < 0.05$, ** $P < 0.01$.

Figure Legends

Figure 1. Immunoepitidome Analysis of Mass Spectrometry-Identified Peptides from Human Placental gp96. (A) Disease enrichment analysis of proteins from placental gp96-bound peptides identified using mass spectrometry. (B-C) Female HLA-A2.1/Kb transgenic mice were immunized with BMDCs pulsed with Pgp96, recombinant gp96 (Rgp96), or tumor lysate as a positive control (PC), or with unpulsed BMDCs (DC) or PBS alone as a negative control. Splenocytes from immunized mice were stimulated with HepG2, PANC1, or A549 whole-cell lysates and assayed using an IFN- γ ELISPOT (B). Splenocytes from immunized mice were stimulated and analyzed for cytotoxic activity with CFSE-labeled HepG2, PANC1, or A549 cells as target cells (C). All experiments were conducted three times. Results are presented as mean \pm SD of three mice/group. * $P < 0.05$, ** $P < 0.01$. (D) Venn diagram showing distribution of shared genes of proteins from placental gp96-bound peptides and genes in the Gene Expression Omnibus public transcriptome dataset for hepatocellular carcinoma (HCC) (GSE101675) and placenta (GSE1133). (E) Venn diagram showing common genes of highly expressed genes in three different datasets for HCC. (F) t-Stochastic neighborhood embedding dimensionality reduction and gene expression of single-cell sequencing data of normal delivery placenta (GSE173193 database). (G) t-Stochastic neighborhood embedding dimensionality reduction and gene expression of single-cell sequencing data of HCC.

****Figure 2. Analysis of HLA-A*0201-Restricted Epitopes Within Placental gp96-Bound GPC3 or PEG10 Peptides.**** (A) Binding affinity of predicted epitopes was quantified using MHC stabilization assays with T2 cells using flow cytometry. HBC¹⁸⁻²⁷ and HBC⁸²⁻⁹⁰ peptides served as positive (PC) and negative controls (NC). (B) Schema of animal experiments for vaccination strategy. Female HLA-A2.1/Kb mice were subcutaneously immunized with BMDCs pulsed with a mix of three GPC3 epitopes or four PEG10 epitopes using Rgp96 as an adjuvant at weeks 1 and 2. (C) Splenocytes were isolated and stimulated with a single epitope (10 μ g/mL). Epitope-specific IFN- γ ⁺CD8⁺ T cells were ana-

lyzed using flow cytometry. (D-E) Female HLA-A2.1/Kb transgenic mice were subcutaneously immunized with BMDCs pulsed with Rgp96, Pgp96, Rgp96 complexed with GPC3¹⁵²⁻¹⁶⁰ or PEG10²²⁹⁻²³⁷ epitope, or control BMDCs (DC) at weeks 1 and 2. Isolated splenocytes were stimulated and stained with GPC3¹⁵²⁻¹⁶⁰ or PEG10²²⁹⁻²³⁷ tetramer. Staining with HBc¹⁸⁻²⁷ tetramer was used as negative tetramer control. The percentage of GPC3¹⁵²⁻¹⁶⁰ or PEG10²²⁹⁻²³⁷ epitope-specific tetramer⁺CD8⁺ T cells was analyzed using flow cytometry (D). Splenocytes were detected in an IFN- γ ELISPOT assay. The control peptide HBcAg¹⁸⁻²⁷ was used for background evaluation (E). Data are presented as mean \pm SD of five mice. Ns, not significant; *P < 0.05, **P < 0.01.

Figure 3. Binding Modes of GPC3- and PEG10-Derived Epitope/HLA-A2 Complexes and Epitope-Harboring Polypeptide/gp96-Dimer Complexes. (A-B) Models of GPC3¹⁵²⁻¹⁶⁰ epitope/HLA-A2 complexes (A) and PEG10²²⁹⁻²³⁷ epitope/HLA-A2 complexes (B) were constructed through molecular docking analysis. Three-dimensional structure and interaction sites of the binding pocket were visualized using PyMol software (left panel), and two-dimensional interactions of epitopes and their targets were exhibited, with hydrogen bonding interactions indicated in green (right panel). (C-D) Models of GPC3 polypeptide (aa 149-185)/gp96-dimer complexes (C) and PEG10 polypeptide (aa 201-240)/gp96-dimer complexes (D) using ZDOCK. Structures of gp96 and GPC3 and PEG10 polypeptides were modeled using AlphaFold2. Left panel shows three-dimensional structure and interaction sites of the binding pocket. Right panel represents binding mode of peptides in binding pocket formed by two loops of the middle-domain and client protein domain of gp96 protein. (E-F) Important binding sites for interaction between gp96-dimer and GPC3 (E) or PEG10 (F) polypeptide analyzed using Discovery Studio. (G-H) Interaction of gp96-dimer with GPC3 (G) or PEG10 (H) polypeptide under different torsion dimer states. Left panel of G or H shows interaction between GPC3 or PEG10 polypeptide with gp96-NMC/AMP-PNP-bound closed dimer, gp96-NMC/AMP-PNP-bound “twisted V” dimer, or gp96-NMC/ADP-bound “twisted V” dimer. Right panel of G or H represents ZDOCK score analysis of the top 10 complex conformations. Ns, not significant; $P < 0.05$, $P < 0.01$, $P < 0.001$, ****P < 0.0001.

Figure 4. GPC3¹⁵²⁻¹⁶⁰ or PEG10²²⁹⁻²³⁷-Specific T Cells Possess Anti-Hepatocellular Carcinoma (HCC) Activity. (A) Volcano plots of differentially expressed genes (DEGs) from HepG2 cells compared to SK-Hep-1 cells. DEGs were selected by $P < 0.05$ and $|\log_2(\text{fold-change})| > 2$. X-axis shows fold-change in gene expression between HepG2 and SK-Hep-1 cells, and y-axis shows significance of differences. Colors represent different genes: black indicates genes without significantly different expression and red indicates significantly differentially expressed genes. (B) Relative expression of GPC3 or PEG10 was analyzed using quantitative real-time PCR. Data are mean \pm SD of three replicates. (C) Western blot analysis of GPC3 and PEG10 expression. (D) Venn diagrams illustrating number of common genes between placental gp96-binding protein

genes and genes highly expressed in HepG2 compared to SK-Hep-1 cells. (E) Female HLA-A2.1/Kb transgenic mice were subcutaneously immunized with BMDCs pulsed with Rgp96, Pgp96, Rgp96-GPC3¹⁵²⁻¹⁶⁰, Rgp96-PEG10²²⁹⁻²³⁷, or Rgp96-GPC3¹⁵²⁻¹⁶⁰ and PEG10²²⁹⁻²³⁷ complexes, or unpulsed control DCs at weeks 1 and 2. Five days after immunization, splenocytes were harvested for cytotoxic activity with CFSE-labeled HepG2 or SK-Hep-1 cells as target cells. (F) Schedule of animal experiments for T cell transfer strategies in HepG2-bearing mice. Mice treated with PBS were used as negative control. (G) Representative images and weights of tumors isolated from HepG2-bearing mice. (H) Size of HepG2 tumors was measured at 2-day intervals. n = 5 mice/group. PC, positive control. Ns, not significant; *P < 0.05, **P < 0.01.

Figure 5. GPC3 and PEG10 Peptides Associated with Placental gp96 Have Therapeutic Effects Against Hepatocellular Carcinoma (HCC).

Female HLA-A2.1/Kb transgenic mice were subcutaneously immunized with BMDCs pulsed with Rgp96-GPC3 polypeptide (aa 149-185), Rgp96-PEG10 polypeptide (aa 201-240) complex, or Rgp96 alone as control at weeks 1 and 2. At 5 days after the last immunization, splenocytes were harvested and stimulated in vitro for 5 days. (A) Splenocytes were harvested for cytotoxic activity with CFSE-labeled HepG2 or SK-Hep-1 cells as target cells. (B) Splenocytes from immunized mice were stimulated with HepG2 or SK-Hep-1 whole cell lysate antigens and assayed in IFN- γ ELISPOT assays. (C-D) HepG2 tumor-bearing mice (n = 6/group) were intravenously injected with GPC3 or PEG10 polypeptide-induced T cells ($3-4 \times 10^6$ /mice) at indicated times. Representative images and weights of tumors from mice with indicated treatment are shown (C). Tumor volumes were measured at 2-day intervals (D). Results are mean \pm SD. Ns, not significant. Experiments were performed twice with similar results. *P < 0.05, **P < 0.01.

Figure 6. Schematic Illustration of Using Chaperone gp96 to Capture and Identify Antigenic Peptides from Placenta Using Mass Spectrometry Followed by Immunopeptidome Analysis.

Placenta-derived gp96 binds abundant tumor-associated antigen peptides associated with multiple cancers. Antigenic peptides derived from GPC3 and PEG10 were identified from among gp96-bound peptides, which elicit anti-HCC activity.

Table 1. Prediction of HLA-A-Restricted Epitopes of Placental gp96-Bound Peptides Identified by Mass Spectrometry (Related to [Figure 2: see original paper])

Peptide Sequences (Position)	HLA Type	Epitopes	Position	Score	Percentile Rank
GPC3 RDLKVFNGNFPKLIMTQMSKASLQVRFKLNQAKNIIIGV- (149-185)	A*02:01				-

Peptide Sequences (Position)	HLA Type	Epitopes	Position	Score	Percentile Rank
	HLA- A*11:01	KLIMTQVSK		-	-
NVLLGLFSTIHDSIQYVQIKNAGKIMLLTICFSDTAHSQQRQYRSAYYPEDLFIDKKVLKV (264-321)	A*02:01	TIHDSIQYV FIDKKVLKV		- -	- -
	HLA- A*11:01	STIHDSIQY		-	-
HSPLKLLTSMAISVVCFFHLA/H (505-526)	A*24:02	AYYPEDLFI		-	-
YTNAMFKNNYPSLTPQAEAFVGBELDSSLAISV (71-100)	A*02:01	AMFKNNYPSL		-	-
	HLA- A*24:02				
PEG10					
AHLATYTEFVPQIPGYQILXPTYAAYAPITYPVGF (620-652)	A*02:01	YAAPTYPV		-	-
	HLA- A*11:01	QTYPTYAAY		-	-
	HLA- A*24:02				
ALIDQYHEGLSDHIQEESHLEVAKSESAMTCQCIHIERR (201-240)	A*02:01	TYTEFVPQI AAYPTYVGF ALIDQYHEGL ALIGQCIEE GLSDHIQEEL	19-237	- - - - -	- - - - -
QVQKLTEENTTLREQVHPTPEDHODDEENTTL (33-59)	A*11:01			-	-
DHRLVDPHIEMIPGAHSHPSGHVYSLSEPEMAAILR (524-558)	A*02:01			-	-
NPDMLAPFMAQCQIFMEKSTRDRSVDPHIEMI (89-114)	A*02:01	FMAQCQIFM MLAPFMAQC		- -	- -
	HLA- A*11:01	AQCQIFMEK		-	-
	HLA- A*24:02				
VRWLSTHDPNITWSTRSIVADSEYCRYSSTRSLSPIPPSLPP (445-482)	A*02:01			-	-
	HLA- A*11:01	WLSTHDPNI		-	-

Peptide Sequences (Position)	HLA Type	Epitopes	Position	Score	Percentile Rank
	HLA- A*24:02	IVFDSEYCR		-	-
		MYSPIPPSL		-	-

The position of epitopes according to Protein Group Accessions (P51654-2 for GPC3, A0A087WXX2 for PEG10). The 8- to 10-mer epitopes were predicted to bind to HLA-A02 (A0201), HLA-A11 (A1101), and HLA-A24 (A2402) using different algorithms: the Immune Epitope Database (IEDB) and SYFPEITHI.

SYFPEITHI (<http://www.syfpeithi.de/bin/MHCServer.dll/EpitopePrediction.htm>) score for each peptide. A high score indicates better MHC affinity.

NetMHCpan EL 4.1 (<http://tools.iedb.org/main/tcell/>) score for each peptide. A low percentile rank indicates better MHC affinity.

Peptide and epitope were used for anti-HCC animal assay.

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv –Machine translation. Verify with original.