

Improved preparation and chemical kinetics on fully automated synthesis of [18F]-THK523, a PET imaging probe for Tau pathologies postprint

Authors: KONG Yan-Yan, SI Zhan, CAO Guo-Xian, ZHANG Zheng-Wei, WU Ping, XUE Fang-Ping, DU Fu-Qiang, ZHU Jian-Hua, Cong Li, CHEN Jian, GUAN Yi-Hui

Date: 2023-06-18T00:00:00+00:00

Abstract

Extensive accumulation of neurofibrillary tangles (NFTs) consistently correlate with the degree of cognitive impairment and neuronal circuitry deterioration associated with Alzheimer's disease. However, no PET probe is currently available for selective detection of NFTs in the living human brain. [18F]-THK523 was developed as a potential in vivo imaging probe for tau pathology. In this paper, we report a new protected precursor, 2-((2-(4-((tert-butoxycarbonyl)amino)phenyl)quinolin-6-yl)oxy)ethyl 4-methylbenzenesulfonate (THK-7), instead of 2-((2-(4-aminophenyl)quinolin-6-yl)oxy)ethyl 4-methylbenzenesulfonate (BF241), and an improved automated radiosynthesis of [18F]-THK523 and the study on chemical kinetics of the labeling reaction of [18F]-THK523, with high-yield ($70\pm 5\%$, $n=6$, decay-corrected to end of bombardment), and high radiochemical purity ($>90\%$) and specific activity (2.5 ± 0.5 Ci/mol) from protected precursor on fully automated module at the end of radiosynthesis (45–55 min). The chemical kinetics for [18F]-THK523 demonstrates that nucleophilic substitution can be carried out easily with protected precursor.

Full Text

Preamble

NUCLEAR SCIENCE AND TECHNIQUES 25, 040302 (2014)

Improved Preparation and Chemical Kinetics on Fully Automated Synthesis of [18F]-THK523, a PET Imaging Probe for Tau Pathologies

Authors: Kong Yan-Yan¹, Si Zhan¹, Cao Guo-Xian², Zhang Zheng-Wei¹, Wu Ping¹, Xue Fang-Ping¹, Du Fu-Qiang¹, Zhu Jian-Hua³, Li Cong³, Chen Jian³, and Guan Yi-Hui¹,†

Affiliations: ¹PET Center, Huashan Hospital, Fudan University, Shanghai 200235, China

²Key Laboratory of Nuclear Medicine, Ministry of Health, Jiangsu Key Laboratory of Molecular Nuclear Medicine, Jiangsu Institute of Nuclear Medicine, Wuxi 214063, China

³Key Laboratory of Smart Drug Delivery, Ministry of Education & PLA, School of Pharmacy, Fudan University, Shanghai 200032, China

Received: October 22, 2013; **Accepted in revised form:** February 13, 2014;

Published online: August 6, 2014

Abstract: Extensive accumulation of neurofibrillary tangles (NFTs) consistently correlates with the degree of cognitive impairment and neuronal circuitry deterioration associated with Alzheimer's disease. However, no PET probe is currently available for selective detection of NFTs in the living human brain. [18F]-THK523 was developed as a potential in vivo imaging probe for tau pathology. In this paper, we report a new protected precursor, 2-((2-(4-((tert-butoxycarbonyl)amino)phenyl)quinolin-6-yl)oxy)ethyl 4-methylbenzenesulfonate (THK-7), instead of 2-((2-(4-aminophenyl)quinolin-6-yl)oxy)ethyl 4-methylbenzenesulfonate (BF241), and an improved automated radiosynthesis of [18F]-THK523 and the study on chemical kinetics of the labeling reaction of [18F]-THK523, with high-yield ($70 \pm 5\%$, $n = 6$, decay-corrected to end of bombardment), and high radiochemical purity ($> 90\%$) and specific activity (2.5 ± 0.5 Ci/ μmol) from protected precursor on fully automated module at the end of radiosynthesis (45–55 min). The chemical kinetics for [18F]-THK523 demonstrates that nucleophilic substitution can be carried out easily with protected precursor.

Keywords: Fluorine-18 radiolabeling, Neurofibrillary tangles, Alzheimer's disease, Automated radiosynthesis, Chemical kinetics

DOI: 10.13538/j.1001-8042/nst.25.040302

INTRODUCTION

Alzheimer's disease (AD) is characterized by progressive impairment of cognitive abilities, such as memory, learning, and social skills. The effects of AD are devastating for both patients and their families, affecting everyday life. AD mostly afflicts the elderly, resulting in an economic challenge for healthcare systems as the elderly population grows [1].

A clinical diagnosis of AD is assessed by several tools, such as medical history and neuropsychological criteria (e.g., National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's disease and Related Disorders Association) [2]. However, a definitive diagnosis of AD can only be

made by confirming the presence of cerebral extracellular senile plaques (SPs) and intracellular neurofibrillary tangles (NFTs) from a postmortem assessment. SPs and NFTs consist of amyloid- β peptides ($A\beta$) and paired helical filaments (PHFs) of hyperphosphorylated tau protein, which are neuropathological hallmarks in AD.

Tau protein is a microtubule-associated protein present in axons, with roles that include stabilization of microtubules, modulation of cytoskeleton plasticity, and promotion of neurite outgrowths. When hyperphosphorylated, tau proteins aggregate into PHFs and NFTs, resulting in destabilization of microtubules and disruption of axonal transport [3–5]. Eventually, these stresses cause neuronal deterioration and ultimately neuronal death. The severity of NFT accumulation correlates with the degree of cognitive impairment and neuronal deterioration associated with AD [6–9]. This correlation makes NFTs potential biomarkers that can be targeted to study how AD pathology progresses and its association with cognitive deterioration [3, 10, 11].

The detection of NFTs in the early stages of AD might be of great value for diagnostic and treatment purposes. Positron emission tomography (PET) is a noninvasive diagnostic imaging modality that utilizes radioisotope-labeled target molecular probes and is considered a diagnostic tool that enables early detection of pathologies. Synthesizing molecular probes with high specificity and affinity to NFTs for PET imaging can allow earlier diagnosis and monitoring of AD progression in vivo [12].

Reports on PET imaging agents selectively targeting NFTs include: radioiodinated styrylbenzimidazole (SBIM) derivative [13]; radioiodinated phenyldiazenylbenzothiazole (PDB) derivative [14]; quinoline and benzimidazole derivatives, [11C]BF158 [15] and [18F]-THK523 [16]; rhodanine and thiohydantoin derivative [17]; phenyldiazenyl benzothiazole derivatives, [18F]T807/808 [3, 18] and [18F]-(E)-4-((6-(2-(2-(2-fluoroethoxy)ethoxy)ethoxy)benzo[d]thiazol-2-yl)-diazenyl)-N,N-dimethylaniline ([18F]FPPDB) [19], and 2-(1-{6-[(2-[F]fluoroethyl)(methyl)amino]-2-naphthyl}ethylidene) malononitrile {[18F]FDDNP} [20]. One of these imaging agents, [18F]-THK523, displayed high affinity and selectivity for tau pathology both in vitro and in vivo [16, 21]. In this paper, we elevate the automated radiosynthesis and reaction kinetics of [18F]-THK523 from a new precursor, 2-((2-(4-((tert-butoxycarbonyl)amino)phenyl)quinolin-6-yl)oxy)ethyl 4-methylbenzenesulfonate (THK-7). This differs from the previous synthesis of [18F]-THK523 by Fodero-Tavoletti et al. [Figure 1: see original paper], which utilized 2-((2-(4-aminophenyl)quinolin-6-yl)oxy)ethyl 4-methylbenzenesulfonate (BF241) as a precursor [17].

II. EXPERIMENTAL

A. Reagents and Instrumentation

KryptofixTM2.2.2, sodium bicarbonate, sodium hydroxide, and hydrochloric acid of analytical grade were purchased from Sigma-Aldrich Corporation (St. Louis, MO, USA). Acetonitrile and ethanol of HPLC grade were obtained from Shanghai Lingfeng Chemical Reagent Co., Ltd. (Shanghai, China). Sep-Pak tC18 solid phase extraction (SPE) cartridge (78.4 μm particle size) and sterile filters (0.22 μm) were purchased from Waters Corporation (Milford, Massachusetts, USA).

Semi-preparative high-performance liquid chromatography was conducted using a Waters pump (Waters Corporation, Milford, Massachusetts, USA) with a Bioscan radioactivity detector. Analytical radio-HPLC (Waters Corporation) was equipped with a dual λ absorbance detector (Waters 2487), binary HPLC pump (Waters 2487), and a Bioscan radioactivity detector. TLC plate radioactivity was measured on a Wizard 1470 automatic gamma counter (U.S. Perkin Elmer Company) equipped with a multi-channel analyzer. The [18F]-THK523 synthesis module (PET-IT-I Reactor Module) was purchased from PET Science & Technology Co. Ltd. (Beijing, China). NMR and LC-MS were purchased from Bruker Corporation (Germany).

B. Precursor Synthesis

1. Preparation of 6-methoxy-2-(4-nitrophenyl)quinoline: THK-1 A mixture of 4-nitro-cinnamaldehyde (20 g, 112.9 mmol) and methoxyphenethylamine (25 g, 203 mmol) was combined in 37% HCl (70 mL) under nitrogen and heated to 140 °C reflux for 3 h. The solution was poured into ice water and adjusted to pH 8 with ammonia. The aqueous layer was extracted with three portions of ethyl acetate. The combined organic layer was washed with saturated sodium chloride, dried, filtered, and concentrated. The residue was purified by flash column chromatography (N-hexane : Dichloromethane = 1 : 1) to afford THK-1 (8.8 g, 31.4 mmol) in 28% yield. ¹H NMR (300 MHz, CDCl₃) δ 8.30–8.40 (m, 4H), 8.18 (d, 1H, J = 9.3 Hz), 8.08 (d, 1H, J = 9.3 Hz), 7.89 (d, 1H, J = 8.6 Hz), 7.43 (dd, 1H, J = 8.3, 2.8 Hz), 7.12 (d, 1H, J = 2.8 Hz); LC-MS: calculated for C₁₆H₁₂N₂O₃, 280.08; found [M+H] 281.0.

2. Preparation of 2-(4-nitrophenyl)quinolin-6-ol: THK-2 THK-1 (3 g, 10.7 mmol) was added to 30% HBr (260 mL) and heated to 125–130 °C reflux for 3 h. The solution was alkalinized by NaHCO₃ and extracted with ethyl acetate. The combined organic layer was washed with saturated sodium chloride, dried, filtered, and concentrated. The residue was purified by flash column chromatography (Dichloromethane : Ethyl acetate = 10 : 1) to afford THK-2 (2.35 g, 8.8 mmol) in 82.5% yield. ¹H NMR (300 MHz, d-DMSO) δ 10.2 (s, 1H), 8.51 (d, 2H, J = 9.2 Hz), 8.37 (d, 2H, J = 9.0 Hz), 8.33 (d, 1H, J = 8.5 Hz), 8.16 (d, 1H, J = 8.6 Hz), 7.98 (d, 1H, J = 9.1 Hz), 7.39 (dd, 1H, J = 9.1,

2.7 Hz), 7.21 (d, 1H, J = 2.7 Hz); LC-MS: calculated for C₁₆H₁₂N₂O₃, 266.07; found [M+H] 267.0.

3. Preparation of 6-(2-((tert-butyldiphenylsilyl)oxy)ethoxy)-2-(4-nitrophenyl)quinoline: THK-3 THK-2 (2.35 g, 8.8 mmol) was dissolved in 80 mL acetonitrile. To the solution were added K₂CO₃ (14.6 g, 105.6 mmol) and TBDPSOCH₂CH₂Br (4.8 g, 13.2 mmol). The mixture was heated to 90 °C reflux for 16 h under nitrogen. The solution was filtered, washed with dichloromethane, and evaporated. The residue was purified by flash column chromatography (Dichloromethane : N-hexane = 1 : 1) to afford compound THK-3 (4.1 g, 7.5 mmol) in 84.7% yield. ¹H NMR (300 MHz, CDCl₃) δ 8.31–8.59 (m, 4H), 8.14 (d, 1H, J = 8.4 Hz), 8.07 (d, 2H, J = 8.3 Hz), 7.88 (d, 1H, J = 8.5 Hz), 7.71–7.74 (m, 4H), 7.35–7.45 (m, 7H), 7.09 (d, 1H, J = 2.7 Hz), 4.25 (t, 2H, J = 5.1 Hz), 4.09 (t, 2H, J = 5.0 Hz), 1.08 (s, 9H).

4. Preparation of 4-(6-(2-((tert-butyldiphenylsilyl)oxy)ethoxy)quinolin-2-yl)aniline: THK-4 THK-3 (3 g, 5.5 mmol) was dissolved in 300 mL ethanol and cooled to 0 °C with an ice bath. Anhydrous Cu(OAc)₂ (1.31 g, 7.2 mmol) was added to the solution. NaBH₄ (10.41 g, 275.2 mmol) was added portionwise within 25 min. The mixture was cooled for 10 min in an ice bath. Then the solution was warmed to room temperature and stirred for 3.5 h. All the above reactions were carried out under nitrogen. The mixture was poured into 50 mL water and ethanol was evaporated. The aqueous layer was extracted with ethyl acetate. The organic layer was washed, dried, filtered, and concentrated to afford THK-4 (2.4 g, 4.6 mmol) in 83% yield. ¹H NMR (300 MHz, CDCl₃) δ 7.97–8.01 (m, 4H), 7.71–7.76 (m, 5H), 7.30–7.43 (m, 7H), 7.02 (d, 1H, J = 2.8 Hz), 6.80 (d, 2H, J = 8.6 Hz), 4.21 (t, 2H, J = 5.1 Hz), 4.07 (t, 2H, J = 5.1 Hz), 3.80 (brs, 2H), 1.07 (s, 9H); LC-MS: calculated for C₃₃H₃₄N₂O₂Si, 518.24; found [M+H] 519.2.

5. Preparation of tert-butyl(4-(6-(2-((tert-butyldiphenylsilyl)oxy)ethoxy)quinolin-2-yl)phenyl)carbamate: THK-5 THK-4 (2.4 g, 4.6 mmol), Boc₂O (3 g, 13.7 mmol), and triethylamine (1.8 g, 17.8 mmol) were dissolved in tetrahydrofuran (THF) (30 mL) and heated to 90 °C reflux for 16 h under nitrogen protection. The solvent was removed under reduced pressure. The residue was purified by column chromatography (Dichloromethane : Ethyl acetate = 8 : 1 → 4 : 1) to afford THK-5 (2.4 g, 3.9 mmol) in 84% yield. ¹H NMR (300 MHz, CDCl₃) δ 8.09 (d, 2H, J = 8.8 Hz), 8.03 (d, 1H, J = 8.6 Hz), 8.02 (d, 1H, J = 9.1 Hz), 7.79 (d, 1H, J = 8.6 Hz), 7.70–7.75 (m, 4H), 7.51 (d, 2H, J = 8.7 Hz), 7.32–7.45 (m, 7H), 7.04 (d, 1H, J = 2.8 Hz), 6.59 (s, 1H), 4.22 (t, 2H, J = 5.1 Hz), 4.07 (t, 2H, J = 5.1 Hz), 1.55 (s, 9H), 1.07 (s, 9H).

6. Preparation of tert-butyl (4-(6-(2-hydroxyethoxy)quinolin-2-yl)phenyl)carbamate: THK-6 THK-5 (1.01 g, 1.6 mmol) was dissolved in 60 mL THF. Bu₄NF · 3H₂O (2 g, 6.3 mmol) in 10 mL THF was added to the

solution. The reaction was allowed to stir at room temperature for 15–20 min under nitrogen. The solvent was removed under reduced pressure. The residue was purified by column chromatography (Dichloromethane : Ethyl acetate = 4 : 1) to afford THK-6 (0.5 g, 1.3 mmol) in 65% yield. ^1H NMR (300 MHz, CDCl_3) δ 9.55 (s, 1H), 8.27 (d, 1H, $J = 8.5$ Hz), 8.13 (d, 2H, $J = 8.8$ Hz), 8.02 (d, 1H, $J = 8.7$ Hz), 7.93 (d, 1H, $J = 8.9$ Hz), 7.61 (d, 2H, $J = 8.8$ Hz), 7.35–7.42 (m, 2H), 4.93 (t, 1H, $J = 5.5$ Hz), 4.13 (t, 2H, $J = 4.8$ Hz), 3.80 (dd, 2H, $J = 4.8, 5.4$ Hz), 1.50 (s, 9H).

7. Preparation of 2-((2-(4-((tert-butoxycarbonyl)amino)phenyl)quinolin-6-yl)oxy)ethyl 4-methylbenzenesulfonate: THK-7 THK-6 (0.48 g, 1.2 mmol) was dissolved in 27 mL dichloromethane. After the mixture was cooled to -5 °C in an ice/sodium chloride bath, triethylamine (0.842 g, 8.3 mmol) and DMAP (0.154 g, 1.3 mmol) were added. TsCl (1.07 g, 5.6 mmol) was added at -10 °C. Then the ice bath was removed and the mixture was allowed to stir overnight at room temperature. The above reaction was carried out under N_2 atmosphere. The solvent was removed under reduced pressure. The residue was purified by column chromatography (Dichloromethane : Ethyl acetate = 30 : 1 \rightarrow 8 : 1) to afford 600 mg of a yellow solid. Finally, pure product THK-7 (0.59 g, 1.1 mmol) was crystallized from ethyl acetate (20 mL) and N-hexane (35 mL) in 87.5% yield. ^1H NMR (300 MHz, CDCl_3) δ 9.56 (s, 1H), 8.23 (d, 1H, $J = 8.7$ Hz), 8.14 (d, 2H, $J = 8.8$ Hz), 8.03 (d, 1H, $J = 8.8$ Hz), 7.91 (d, 1H, $J = 9.0$ Hz), 7.82 (d, 2H, $J = 8.3$ Hz), 7.62 (d, 2H, $J = 8.8$ Hz), 7.46 (d, 2H, $J = 8.0$ Hz), 7.24–7.30 (m, 2H), 4.42–4.45 (m, 2H), 4.30–4.33 (m, 2H), 2.39 (s, 3H), 1.50 (s, 9H); LC-MS: calculated for $\text{C}_{29}\text{H}_{30}\text{N}_2\text{O}_6\text{S}$, 534.18; found $[\text{M}+\text{H}]$ 535.0.

C. Standard Synthesis for [18F]-THK523: tert-butyl (4-(6-(2-fluoroethoxy)quinolin-2-yl)phenyl)carbamate (THKF-2)

1. Preparation of 6-fluoro-2-(4-nitrophenyl)quinoline: THKF-1 THK-2 (0.5 g, 1.9 mmol), K_2CO_3 (2.6 g, 18.8 mmol), and $\text{BrCH}_2\text{CH}_2\text{F}$ (0.5 g, 3.9 mmol) were added to 24 mL acetonitrile. The mixture was refluxed for 16 h at room temperature under nitrogen. The solution was filtered and evaporated. The residue was purified by column chromatography (N-hexane : Ethyl acetate = 4 : 1 \rightarrow 3 : 1) to afford THKF-1 (0.55 g, 1.8 mmol) in 93.4% yield. ^1H NMR (300 MHz, CDCl_3) δ 8.31–8.39 (m, 4H), 8.18 (d, 1H, $J = 8.5$ Hz), 8.12 (d, 1H, $J = 9.3$ Hz), 7.90 (d, 1H, $J = 8.6$ Hz), 7.48 (dd, 2H, $J = 9.2, 2.8$ Hz), 7.14 (d, 2H, $J = 2.8$ Hz), 4.86 (dm, 2H, $J = 47.4$ Hz), 4.36 (dm, 2H, $J = 27.6$ Hz).

2. Preparation of tert-butyl (4-(6-(2-fluoroethoxy)quinolin-2-yl)phenyl)carbamate: THKF-2 THKF-1 (0.55 g, 1.8 mmol) was dissolved in ethanol (120 mL) under nitrogen. The solution was cooled to 0 °C and then anhydrous $\text{Cu}(\text{OAc})_2$ (0.42 g, 2.3 mol) was added. NaBH_4 (2 g, 52.9 mol) was added portionwise at -5 °C. The solution was allowed to stir at -5 °C for 30 min. Then the mixture was allowed to warm to room temperature while stirring for 1 h. The solvent was evaporated and the mixture was poured into

50 mL water. The aqueous layer was extracted with ethyl acetate. The organic layer was washed, dried, filtered, and concentrated. The residue was purified by flash column chromatography (N-hexane : Ethyl acetate = 6 : 1 → 3 : 1) to afford THKF-2 (0.21 g, 0.7 mmol) in 42% yield. ^1H NMR (300 MHz, CDCl_3) δ 7.97–8.04 (m, 4H), 7.76 (d, 1H, $J = 8.7$ Hz), 7.39 (dd, 1H, $J = 9.3, 2.7$ Hz), 7.07 (d, 1H, $J = 2.7$ Hz), 6.80 (d, 1H, $J = 8.6$ Hz), 4.88 (dm, 2H, $J = 47.4$ Hz), 4.34 (dm, 2H, $J = 27.6$ Hz); LC-MS: calculated for $\text{C}_{17}\text{H}_{15}\text{FN}_2\text{O}$, 283.13; found $[\text{M}+\text{H}]$ 283.0.

I. Radiosynthesis of [18F]-THK523

[18F]-THK523 was synthesized using an automated module (PET Science & Technology Co. Ltd., Beijing, China) [Figure 2: see original paper]. The non-carrier-added aqueous [18F]fluoride was produced by the $^{18}\text{O}(\text{p},\text{n})^{18}\text{F}$ nuclear reaction on an Eclipse HP Cyclotron (Siemens).

The synthesis of [18F]-THK523 includes five steps: 1) azeotropic evaporation, 2) nucleophilic substitution, 3) hydrolysis for deprotection, 4) purification, and 5) sterile filtration [Figure 3: see original paper]. Briefly, [18F]fluoride was trapped on a QMA cartridge and eluted into the reaction vial with 1.5 mL of a mixture of K_2CO_3 in water and KryptofixTM2.2.2 in acetonitrile (19/31 mmol/L) that was pre-added in bottle 1#. The mixture in the vial was evaporated at 116 °C for 206 s under a N_2 flow and was co-evaporated to dryness with anhydrous acetonitrile (2 mL) in bottle 2# at 116 °C for another 203 s. The tosylated precursor (2 mg in 1 mL acetonitrile) in bottle 3# was added to the dried K[18F] and the nucleophilic substitution reaction was carried out at 120 °C for 15 min to afford [18F]-THK. After excess acetonitrile was removed at 120 °C under a flow of nitrogen, HCl (1 mol/L, 250 μL) in bottle 4# was added to hydrolyze the Boc protecting group. The mixture was allowed to react at 105 °C for 5 min, and NaOH (2 mol/L, 125 μL) stored in bottle 5# was added followed by saturated NaHCO_3 (1 mol/L, 125 μL) in bottle 6# to neutralize the solution.

The neutralized solution was loaded on a C-18 Sep-Pak, which was further washed with water to remove free $^{18}\text{F}^-$, KryptofixTM2.2.2, and other polar byproducts. The cartridge was then eluted with ethanol (2 mL). The obtained crude product was collected and injected onto a semi-preparative column (Waters XBridgeTMprep Shield RP18 10 μm , 250 mm \times 10 mm, part No. 186003990, serial No. 101/123041GG01) at a flow rate of 4 mL/min (70% EtOH : 30% H_2O). The fraction containing [18F]-THK523 was collected from 6.0–6.5 min on semi-preparative HPLC and evaporated to dryness. To the residual, 10% ethanol in saline (5 mL) was added and the resulting solution was stabilized with ascorbic acid (2 mg, 0.011 mmol) before sterile filtration through a 0.22 μm membrane filter into a sterile vial.

J. Quality Control of [18F]-THK523


The labeling yield was determined by thin layer chromatography (TLC). Apply 2–4 μL of the crude reaction solution to activated silica gel G60 with fluorescence (F254) plates. Develop the chromatogram in a solvent system consisting of a mixture of ethyl acetate : n-hexane : triethylamine = 4 : 1 : 0.005 (v/v) until the solvent has moved about 3/4 of the length of the chromatogram. Remove the chromatogram and allow it to dry at room temperature. Determine the radioactivity distribution by cutting the chromatogram into 10 pieces of strips with equal length from sampling spot to where the solvent developed and counting each strip in a Wizard 1470 automatic gamma counter (U.S. Perkin Elmer Company) equipped with a multi-channel analyzer. The Rf value of [18F]-THK523 was 0.7–0.8, and the Rf value of free F-18 was 0.0. Two TLCs were run for each tested reaction condition and radiolabeling yields were obtained by averaging the yields of the two runs.

Radiochemical purity (RCP) was determined by analytical radio-HPLC (high-performance liquid chromatography). The final product (20 μL) was injected into the HPLC column (Purospher STAR LPRP-18e endcapped (5 μm), 250 mm \times 4.6 mm, sorbent Lot No. TA1752311, column No. 210072, acetonitrile/0.05% triethylamine in water = 8/2 (v/v), flow rate at 0.6 mL/min) at room temperature. The absorbance measured at 350 nm. The retention time (tR) of the standard and [18F]-THK523 were 6.12 and 5.93 min, respectively. The chemical identity was verified by co-injection with cold standard THKF-2.

K. Studies of Reaction Kinetics

The reaction conditions for the nucleophilic substitution were optimized by studying the reaction kinetics. Six vials, each containing a 30–50 mCi activity of $^{18}\text{F}^-$, were used. After azeotropic evaporation, THK-7 (1 mL, 3.7 mmol/L) in DMSO/CAN (1/5) was added to the vials and reacted at 25 $^\circ\text{C}$, 120 $^\circ\text{C}$, and 160 $^\circ\text{C}$ according to the procedure described above. Three reaction times—2 min, 15 min, and 30 min—were investigated at each temperature. After the same deprotecting reaction workup, solutions were sampled with a capillary and labeling yields were determined by TLC. Films were dried and cut into 10 sections, then counted with a γ -counter. Order of reaction (n), rate constant (k), and activation energy (Ea) of the labeling reaction were calculated with the CHEMKIN code, developed by Cao et al. [22], to quantitatively study the labeling reaction for optimal conditions of nucleophilic substitution.

The rate constant, k, for the formation of [18F]-THK523 was calculated as follows. The reaction can be expressed as (let $a = [\text{K}^{18}\text{F}]_0$, $b = [\text{THK}]_0$, $x = [\text{18F-THK523}]$):



$$t = 0: a, b, 0$$

$$t = t: a-x, b-x, x$$

$$\begin{aligned} d[18\text{F-THK523}] / dt &= k[\text{K18F}] [\text{THK-7}] \\ dx/dt &= k(a - x)(b - x) \\ kdt &= dx/[(a - x)(b - x)] \end{aligned}$$

Integrating each side of the equation between $t = 0 \rightarrow t$ and $x = 0 \rightarrow x$ gives:

$$kt = [\ln b(a - x) - \ln a(b - x)]/(a - b)$$

Since $a - x = 10^{-9} - 10^{-8}$ mol/L and $b = 10^{-5} - 10^{-4}$ mol/L, $b \gg a - x$, the equation can be simplified as:

$$\begin{aligned} kt &= [\ln a/(a - x)]/b \\ bkt &= \ln[1/(1 - x/a)] \end{aligned}$$

Since the Rf value of [18F]-THK523 and ^{18}F are 0.7 and 0.0, respectively, x/a represents the count percentage of [18F]-THK523 relative to ^{18}F . In fact, x/a is the labeling yield of [18F]-THK523, and it can be determined by TLC. Let $P = x/a$, the equation can be written as:

$$\ln[1/(1 - P)] = bkt$$

The plot of $\ln[1/(1 - P)]$ vs. t shows a linear relationship and the rate constant k for the [18F]-THK523 formation can be calculated from the slope.

III. RESULTS AND DISCUSSION

There are several types of fluorinated precursor including the following substitute groups: (a) -OTs (tosylate), (b) -OTf (triflate), (c) -OMs (mesylate), and (d) -ONs (nosylate). In general, the reactivity is $b > a > c > d$, while the stability for the groups is $d > a > c > b$. The tosylate was chosen for design of the [18F]-THK523 precursor because it compromises stability and reactivity. The synthesis of protected precursor THK-7 and cold standard THKF-2 are described in [Figure 4: see original paper] and [Figure 5: see original paper].

The synthesis of THK-7 involves the tosylation of a primary alcohol (THK-6). The Boc protection of the primary amino group in the intermediate THK-6 is important for several reasons. First, primary amino groups are generally more reactive towards tosyl chloride than primary alcohols. Therefore, under the reaction conditions used to form precursor THK-7, the amino group would be tosylated preferentially to the alcohol. Furthermore, it is difficult to remove the aminotosyl protecting group by acid hydrolysis in a short reaction time, which would be detrimental for the radiolabeling reaction.

The choice of Boc as the protective group has the following advantages: Boc can be easily hydrolyzed by acid in a short reaction time and the by-products are CO_2 and iso-butanol (or isobutene), which can be easily evaporated. In addition, the presence of a free amino group in precursor BF241, used in Ref. [16], could lead to low radiosynthesis yield, as THK-6 may interact with K18F to form hydrogen bonds reducing the nucleophilicity of $^{18}\text{F}^-$. Fluorine is the most electronegative element and can form hydrogen bonds with hydrogens on

the amino group, while the Boc-protected amino is a carbamate and is less prone to hydrogen bonding formation.

Compared to the work of Fodero-Tavoletti et al. [16], the most prominent advantage of using THK-7 instead of BF241 for radiolabeling of [18F]-THK523 was the improvement of radiochemical yields: $40 \pm 5\%$ and 24% (non-decay corrected) at end of synthesis for THK-7 and BF241, respectively. There were no significant differences in radiochemical purity and specific activity.

Therefore, Boc protection of the primary amino group in the precursor THK-7 is important to prevent side reactions, simplify product purification, and improve precursor stability.

Radiolabeling yield of [18F]-THK523 was assayed by TLC. As shown in , the radiolabeling yield is relatively high, ranging from 11.91% to 75.11% (decay corrected to end of bombardment, EOB) and the synthesis time is 45–55 min from EOB.

Radio-HPLC analysis showed the radiochemical purity (RCP) of [18F]-THK523 was $\geq 90\%$. As shown in [Figure 6: see original paper], the retention time (tR) of [18F]-THK523 and precursor THK-7 were 6.12 min and 12.94 min, respectively. The chemical identity of [18F]-THK523 was verified by co-injection with the non-labelled standard THKF-2 (tR = 5.93 min).

The specific activity of [18F]-THK523 was determined as 2.5 ± 0.5 Ci/ μ mol (end of synthesis, EOS).

Labeling of radiopharmaceuticals is influenced by reaction time and temperature. Their relationship can be evaluated by chemical kinetics. Although TsO, as a leaving group during nucleophilic substitution, has been applied in many ¹⁸F-labeled radiopharmaceuticals, different molecular structures lead to different labeling yields. Evaluation of chemical kinetics is authentically necessary because of the importance of reaction temperature for labeling reaction. Labeling yields of [18F]-THK523 at different temperatures and times are listed in . The mean labeling yields and mean specific radioactivity increase with temperature and time.

At 120 °C, the rate constants were 4.95, 4.97, 5.00, and 5.04 for THK-7 concentrations of 5.24, 26.2, 52.4, and 131 mmol/L, respectively ().

For reaction chemical kinetics parameters, rate constant (k) increases with temperature, indicating that high temperature is conducive to increased reaction rate. In order to improve radiosynthesis yield, it is necessary to increase reaction temperature. However, [18F]-THK523 may be unstable or the binding of C-O or C-N may break down under such high temperature.

As shown in , labeling yields were high at 120 °C for reactions of 15 min and 30 min (decay-corrected to EOB). Due to favorable thermal stability of [18F]-THK523 and for convenience in experimental operation, the optimal reaction

temperature was chosen as 120 °C when labeling yields were larger than 70% at 15 min. Calculations show the order of reaction is 1.

The chemical kinetics study demonstrated the labeling reaction of 2-(2-(4-(tert-butoxycarbonyl)phenyl)quinolin-6-yloxy)ethyl 4-methylbenzenesulfonate (THK-7) was quite facile. This method can be applied to study labeling reactions of other radiopharmaceuticals.

The plot of $\ln[1/(1 - P)]$ vs. t at four different concentrations of THK-7 didn't show significant linear relationship. The changes of rate constants under different concentrations of precursor are shown in . The rate constant k didn't change greatly with the concentration of THK-7.

The plot of $\ln[1/(1 - P)]$ vs. t at different temperatures showed a better linear relationship. The rate constant k increased with temperature, being 0.04, 4.85, and 5.18 at 25 °C, 120 °C, and 160 °C, respectively. This demonstrated that k of the reaction was temperature dependent. From the variation of the order of magnitude, k had a significant increase from temperature 25 °C (0.04) to temperature 120 °C (4.95). This provides further evidence validating the fact that temperature must be higher than 120 °C.

IV. CONCLUSION

[18F]-THK523 can be synthesized at CPCU with high synthesis yield ($70 \pm 5\%$, $n = 6$, decay-corrected to EOB). The total synthesis time was 45–55 min and the average specific activity was 2.5 ± 0.5 Ci/ μmol for [18F]-THK523 (EOS) under optimum conditions. The chemical kinetics for [18F]-THK523 showed that the reaction order was 1 because the concentration of precursor was much larger than that of ^{18}F . Low E_a was indicative of a low activation barrier. The fact that k increased with reaction temperature indicated that a rise in temperature favors an accelerated reaction rate. $t_{1/2}$ (the time for half of the reactant to react) shortened with increased reaction temperature, which means that increasing temperature accelerates the labeling reaction. The protected precursor used in this method should be applicable for high-yield automated production in a commercial synthesis module for clinical application in the future. The use of THK-7 might be suitable for routine production of high-yield [18F]-THK523 and the radiosynthesis can be easily performed on an automated synthesis module.

REFERENCES

- [1] Wimo A and Winblad B. Handbook of Clinical Neurology. Elsevier, 2008, 89: 137–146.
- [2] McKhann G, Drachman D, Folstein M, et al. Neurology, 1984, 34: 939–939.
- [3] Zhang W, Arteaga J, Cashion D K, et al. J Alzheimers Dis, 2012, 31: 601–612.
- [4] Bulic B, Pickhardt M, Mandelkow E M, et al. Neuropharmacology, 2010, 59: 276–289.

- [5] Pritchard S M, Dolan P J, Vitkus A, et al. J Cell Mol Med, 2011, 15: 1621–1635.
- [6] Aizenstein H J, Nebes R D, Saxton J A, et al. Arch Neurol-Chicago, 2008, 65: 1509–1617.
- [7] Lai M K, Chen C P, Hope T, et al. Neuroreport, 2010, 21: 666–676.
- [8] Arriagada P V, Growdon J H, Hedley-Whyte E T, et al. Neurology, 1992, 42: 631–639.
- [9] Villemagne V, Pike K, Darby D, et al. Neuropsychologia, 2008, 46: 1688–1797.
- [10] Braskie M N, Klunder A D, Hayashi K M, et al. Neurobiol Aging, 2010, 31: 1669–1778.
- [11] Karran E, Mercken M, De Strooper B. Nat Rev Drug Discov, 2011, 10: 698–712.
- [12] Långström B, Andrén P E, Lindhe Ö, et al. Mol Imaging Biol, 2007, 9: 161–175.
- [13] Matsumura K, Ono M, Yoshimura M, et al. Bioorg Med Chem, 2013, 21: 3356–3362.
- [14] Matsumura K, Ono M, Hayashi S, et al. Med Chem Comm, 2011, 2: 596–600.
- [15] Okamura N, Suemoto T, Furumoto S, et al. J Neurosci, 2005, 25: 10857–10962.
- [16] Fodero-Tavoletti M T, Okamura N, Furumoto S, et al. Brain, 2011, 134: 1089–1100.
- [17] Ono M, Hayashi S, Matsumura K, et al. ACS Chem Neurosci, 2011, 2: 269–275.
- [18] Xia C F, Arteaga J, Chen G, et al. Alzheimers Dement, 2013, 9: 666–676.
- [19] Matsumura K, Ono M, Kimura H, et al. ACS Med Chem Lett, 2011, 3: 58–62.
- [20] Liu J, Kepe V, Žabjek A, et al. Mol Imaging Biol, 2007, 9: 46–52.
- [21] Harada R, Okamura N, Furumoto S, et al. Eur J Nucl Med Mol I, 2013, 40: 125–132.
- [22] Cao G X, Zhou X Q, Liu Y T, et al. Nucl Sci Tech, 2012, 23: 040302.

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv — Machine translation. Verify with original.