

New Continuum Observations of the Andromeda galaxy M31 with FAST Postprint

Authors: Wenjun Zhang, Xiaohui Sun and Jie Wang

Date: 2023-12-15T00:00:00+00:00

Abstract

We present a new total intensity image of M31 at 1.248 GHz, observed with the Five-hundred-meter Aperture Spherical radio telescope (FAST) with an angular resolution of and a sensitivity of about 16 mK. The new FAST image clearly reveals weak emission outside the ring due to its high sensitivity on large-scale structures. We derive a scale length of 2.7 kpc for the cosmic ray electrons and find that the cosmic ray electrons propagate mainly through diffusion by comparing the scale length at 4.8 GHz. The spectral index of the total intensity varies along the ring, which can be attributed to the variation of the spectra of synchrotron emission. This variation is likely caused by the change of star formation rates along the ring. We find that the azimuthal profile of the non-thermal emission can be interpreted by an axisymmetric large-scale magnetic field with varying pitch angle along the ring, indicating a complicated magnetic field configuration in M31.

Full Text

Research in Astronomy and Astrophysics, 23:115022 (9pp), 2023 November
© 2023. National Astronomical Observatories, CAS and IOP Publishing
Ltd. Printed in China and the U.K.
<https://doi.org/10.1088/1674-4527/acfae1>

New Continuum Observations of the Andromeda Galaxy M31 with FAST

Wenjun Zhang¹, Xiaohui Sun¹, and Jie Wang²

¹ School of Physics and Astronomy, Yunnan University, Kunming 650500, China; zhangwenjun@mail.ynu.edu.cn, xhsun@ynu.edu.cn

² National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100101, China

Received 2022 December 3; revised 2023 July 23; accepted 2023 September 7;
published 2023 October 17

Abstract

We present a new total intensity image of M31 at 1.248 GHz, observed with the Five-hundred-meter Aperture Spherical radio Telescope (FAST) with an angular resolution of 4' and a sensitivity of about 16 mK. The new FAST image clearly reveals weak emission outside the ring due to its high sensitivity to large-scale structures. We derive a scale length of 2.7 kpc for the cosmic ray electrons and find that they propagate mainly through diffusion by comparing the scale length at 4.8 GHz. The spectral index of the total intensity varies along the ring, which can be attributed to the variation of synchrotron emission spectra. This variation is likely caused by changes in star formation rates along the ring. We find that the azimuthal profile of the non-thermal emission can be interpreted by an axisymmetric large-scale magnetic field with varying pitch angle along the ring, indicating a complicated magnetic field configuration in M31.

Key words: galaxies: ISM – galaxies: magnetic fields – radiation mechanisms: general – (ISM:) cosmic rays

1. Introduction

The Andromeda galaxy (M31) is a prominent spiral galaxy and the nearest major galaxy to the Milky Way (Beck et al. 1998). Due to its proximity and outstanding “ring” of star formation, M31 has been widely studied across various wavelengths, including X-ray (Pietsch et al. 2005), UV (Gil de Paz et al. 2007), near-IR (Jarrett et al. 2003), far-IR (Rice et al. 1988), and optical (Sarkisyan et al. 2020). These observations have improved our understanding of the formation and evolution of M31.

The radio total intensity of M31 is concentrated in a ring about 7–13 kpc from the center (Beck et al. 1998). In this ring, there are also concentrations of cold molecular gas (Nieten et al. 2006), warm neutral gas (Chemin et al. 2009), warm ionized gas (Devereux et al. 1994), and dust (Smith et al. 2012). All these indicate that the M31 ring is an important star-forming region and contributes most of the galaxy’s brightness.

There have been many studies based on radio observations of M31. The first radio observation of M31 was conducted with the Cambridge one-mile radio telescope at 408 and 1407 MHz with resolutions of 80' and 23', respectively (Pooley 1969). With continuing improvements in angular resolution and sensitivity, follow-up observations have revealed more detailed emission structures, such as those by the Effelsberg 100 m telescope at 4850 MHz with a resolution of 2.6' (Berkhuijsen et al. 1983).

In addition, the combination of interferometric observations from the Very Large Array and single-dish observations from the Effelsberg radio telescope at 1.5

GHz yielded an image with a high resolution of $45''$ covering all spatial scales (Beck et al. 1998).

The spectrum of M31 has been studied based on radio observations. The radio emission consists of synchrotron emission (non-thermal) and free-free emission (thermal). At frequencies below 10 GHz, the non-thermal emission is dominant (Battistelli et al. 2019), and the fraction of thermal emission is only about 24% at 1 GHz (Berkhuijsen et al. 2003).

Based on observations at 1.46 and 4.85 GHz, Berkhuijsen et al. (2003) obtained an overall non-thermal spectral index of $\alpha = -1 \pm 0.1$ ($S \propto \nu^{-\alpha}$, with S being the flux density at frequency ν). Beck et al. (2020) derived a total intensity spectral index of -0.71 ± 0.02 and a synchrotron emission index of -0.81 ± 0.03 using observations at 2.645 GHz, 4.85 GHz, and 8.35 GHz. The spectral index toward the ring increases to -0.5 because the thermal fraction is large and the spectrum thus tends to be flat (Fatigoni et al. 2021).

The Five-hundred-meter Aperture Spherical radio Telescope (FAST) is well suited to image extended sources such as M31 with its high sensitivity (Jiang et al. 2020). The M31 Halo is one of the key projects currently being conducted by FAST with guaranteed time (PI: Jie Wang), which focuses on HI mapping in a region of about 800 square degrees around M31 and pulsar searching in the M31 galaxy. The continuum and polarization data have also been recorded simultaneously, allowing us to analyze the total intensity distribution and spatial variation of spectral index to infer the properties of M31.

The paper is organized as follows: the observations and results are described in Section 2, discussed in Section 3, and summarized in Section 4.

2. Observations and Results

M31 was observed in drifting-scan mode (Jiang et al. 2020). In this mode, the 19-beam receiver is rotated by $23^\circ.4$ so that a declination strip of about $22''$ can be completed in Nyquist sampling for each individual scan. In total, 11 drifting scans were conducted during 2020 October 1–18. To calibrate system variations, a linearly polarized noise source of about 12.5 K (Jiang et al. 2019) is injected into the system for 2 s every 1000 s. The frequency range of the receiving system is 1.0–1.5 GHz, which is divided into 65,536 channels. The width of each channel is about 7.63 kHz. The backend outputs four channels: two for total intensity I1 and I2, and the other two for Stokes U and V.

We follow the data processing procedure developed by Sun et al. (2021). To obtain images at a single frequency channel, the following steps were taken: (1) mitigation of radio frequency interference (RFI); (2) correction of instrumental polarization leakage and angle, and temperature scale calibration with injected noise; (3) conversion to brightness temperature scale (Tb). We repeat this procedure and obtain maps of about 29,000 frequency channels. These maps are then smoothed to a common resolution of $4''$. (4) combination of all 11 scans.

2.1. Total Intensity Maps

We average all the frequency channel maps by taking median values and obtain the total intensity image at 1.248 GHz, as shown in Figure 1. Some scanning effects indicated by stripes along the declination direction still remain in the image, which were caused by drifting of the system. These effects would require scans in alternative directions to remove, as shown by Sun et al. (2022). The rms noise is about 16 mK.

A smaller area containing M31 is cut out from Figure 1, rotated by -53° following Beck et al. (1998), and shown in Figure 2 (top panel). For comparison, the map by Beck et al. (1998) combining interferometer and single-dish observations at 1.46 GHz is also shown in Figure 2 (middle panel).

As can be seen from Figures 1 and 2, there is strong radio emission toward the center and the ring area. The FAST image clearly reveals weak emission extending outside the ring toward the southwest. In contrast, this extended emission is almost absent in the image by Beck et al. (1998).

To study the emission from M31, we remove the bright compact sources marked in Figure 2 (top panel) by subtracting Gaussian fits to these sources. The resulting image is shown in Figure 2 (bottom panel), which is used to investigate variations of total intensity below. Note that background point sources that cannot be resolved with FAST still remain. We retrieve the flux intensities of these sources from the NVSS survey with a higher resolution of 45" (Condon et al. 1998) and find that their contribution to the intensity of the ring is less than about 5%. Their influence on the analyses below can thus be ignored.

2.2. Radial Profile of the Total Intensity

The radial variation of total intensity can be used to constrain the propagation of cosmic ray electrons. We derive the average intensity of 6"-width annuli from the center until a radius of about 110". The intensity versus radius is shown in Figure 3. The standard deviation from each annulus is calculated as the error bar.

The sharp peak near radius 0 is contributed by the nucleus of M31. The broad peak from a radius of about 27" to about 57" indicates the ring area with bright emission. With a distance to M31 of about 780 kpc (Stanek & Garnavich 1998), this angular range corresponds to about 6–13 kpc. The total intensity gradually decreases with increasing radius and flattens out beyond a radius of about 93".

We fit the total intensity radial profile for radii larger than 27" to a Gaussian, namely $T(r) = T_0 \exp[-(r-r_0)^2/(2L^2)]$, where r is the radius, r_0 is the peak radius, and L is the scale length. After accounting for the influence of beamwidth, we obtain a scale length of about 16.2 ± 0.1 ", or 3.7 ± 0.1 kpc.

2.3. Azimuthal Profile of the Total Intensity

The variation of total intensity with direction reflects the structure of the magnetic field. From the Gaussian fitting of the radial profile, we obtain the peak position of the ring at $r_0 = 42$. We use the annulus of $r_0 \pm L$ to derive the azimuthal profile of total intensity in the ring. The azimuthal angle starts from the major axis toward the northeast and increases eastwards (Figure 1) or starts from the major axis toward the left and increases counter-clockwise (Figure 2). An average intensity is derived for every 10° sector, and the profile is shown in Figure 4.

The emission is brightest for azimuthal angles between 30° and 90° , and becomes weaker for azimuthal angles between 150° and 270° , as can also be seen from Figure 2. There is no clear pattern for the azimuthal profile.

2.4. Spectral Index

The spectral index is related to emission properties such as non-thermal and thermal fractions and cosmic ray electron aging. We do not derive the spectral index for all pixels by fitting brightness temperature against frequency because of the influence of background emission. Instead, we divide the whole M31 area into eight regions (Figure 5) and obtain the spectral index using the Temperature–Temperature plot (TT-plot) method (Turtle et al. 1962), which is immune to background emission.

We take the median values of the first 10,000 and the last 10,000 channels to calculate the spectral index with TT-plot. The corresponding frequencies are 1.055 and 1.420 GHz, which have the largest frequency separation and hence allow for more accurate determination of the spectral index. The TT-plot results for the seven regions are shown in Figures 5 and 6. Note that the brightness temperature spectral index β , as we obtain, can be connected to the flux density spectral index α as $\alpha = \beta + 2$.

The spectral index for the total intensity is not uniform across M31. Toward the areas marked as 6, 1, and 2 in Figure 5 where the total intensity emission is weak, the spectra are steep with spectral index α between -0.8 and -0.7 . In contrast, in the areas marked as 3, 4, and 5 in Figure 5 where the emission is stronger, the spectra are shallower with spectral index α up to -0.44 ± 0.09 . There seems to be a tendency of intensity correlating with flatter spectrum. The nucleus region marked as 7 has the smallest spectral index of -0.90 ± 0.03 . The area outside M31 marked as 8, detected because of the high sensitivity of FAST, has a spectral index of $\alpha = -0.64 \pm 0.08$ and has not been well studied before.

3. Discussion

3.1. Properties of Cosmic Ray Electrons

The total intensity is composed of both thermal emission from free-free radiation and non-thermal emission from synchrotron radiation. We take the images of thermal and non-thermal emission derived by Tabatabaei et al. (2013) and obtain the thermal fraction versus azimuthal angle for the ring of M31, as shown in Figure 7. The thermal fraction is up to 20% for azimuthal angles between 30° and 90° , and about 10% for the rest. The thermal emission has a flat spectrum with $\alpha_{\text{th}} = -0.1$, which is much larger than that of non-thermal emission. Therefore, the mixture of thermal and non-thermal emission will bring the spectral index of the total emission to a larger value. Typically, for thermal fractions of 10% and 20%, the spectral intensity calculated from the frequency pair of 1.055 and 1.420 GHz is increased by 0.1 and 0.2, respectively, compared with the spectral index of non-thermal emission α_n .

For the ring of M31, the spectra for total intensity toward the north part in Figure 1 (or regions 3, 4, 5 on the left part in Figure 5) are shallower than those toward the south part. Taking into account the influence of thermal emission, the spectral index of non-thermal emission could be estimated by subtracting 0.1 or 0.2 from the total intensity spectral index. For example, $\alpha_n = -0.64$ for region 5 with a thermal fraction of about 20%. This implies that the non-thermal spectral index toward the northern part is larger than that toward the southern part of M31.

The asymmetry of the non-thermal spectral index between the northern and southern parts of M31 was also found by Fatigoni et al. (2021) based on observations at frequencies between 1.5 and 6.6 GHz. The star formation rate is also higher toward the north (Fatigoni et al. 2021). A higher star formation rate results in more stars and consequently more supernovae and supernova remnants that can produce more cosmic ray electrons. Therefore, the spectrum is flatter toward the north. Moreover, the presence of higher cosmic ray electron density results in stronger synchrotron radiation, which explains the correlation between large intensity and large spectral index.

For area 8 in Figure 5, it cannot be an extension from the ring. Otherwise, its spectrum should be steeper because of the aging of cosmic ray electrons as they propagate from the ring. From the Herschel image at 250 μm (Viaene et al. 2014), there is strong emission corresponding to this area, which seems to be part of a large ring structure surrounding M31. There could be star-forming activities that produce high-energy electrons. High-resolution observations are needed to detect radio emission related to the infrared emission observed by Herschel.

The scale length of the total emission is 3.7 kpc. The thermal emission is highly concentrated in the ring (Tabatabaei et al. 2013) and therefore has little influence on the scale length of non-thermal emission. In practice, we derive a scale

length of 3.8 kpc for the non-thermal emission after removing the thermal contribution. This corresponds to an exponential scale length of 5.4 kpc. Assuming energy equipartition between magnetic field and cosmic rays, the scale length of the non-thermal emission is equal to that of cosmic ray electrons multiplied by the factor $(3 - \alpha_{\text{n}})/2$ (Berkhuijsen et al. 2013). The scale length of cosmic ray electrons is thus about 2.7 kpc.

The diffusion length of cosmic ray electrons varies with frequency as $\sim \nu^{-0.4-0.5}$ (Mulcahy et al. 2016). It is therefore expected that the scale length of cosmic ray electrons is about 1.9 kpc at 4.8 GHz, which is consistent with the scale length of about 3.7 kpc for non-thermal emission and hence the scale length of about 1.9 kpc for cosmic ray electrons derived by Beck et al. (2020) at 4.8 GHz. This implies that cosmic ray transport occurs primarily through diffusion and that energy loss is not significant at these frequencies.

3.2. The Magnetic Field in the Ring

We obtain the azimuthal profile of the non-thermal emission within the ring by subtracting the thermal emission contribution (Figure 7) from the total emission (Figure 4). The result is shown in Figure 8. The non-thermal emission originates from synchrotron radiation, and its profile thus reflects the magnetic field configuration in M31.

The large-scale regular magnetic field in M31 mainly follows an axisymmetric pattern (Han et al. 1998; Fletcher et al. 2004). In this configuration, the magnetic field component perpendicular to the line of sight can be derived as $B_{\perp} = B \cos(\phi + \theta) \sin i$, where ϕ is the azimuth angle, θ is the pitch angle, and i is the inclination angle of M31. The non-thermal intensity (I) can be assessed as $I_{\text{nt}} = \int B_{\perp}^{1-\alpha_{\text{n}}} dl$ (Beck & Krause 2005). Here, energy equipartition between magnetic field and cosmic rays is assumed.

The random magnetic field has recently been classified into ordered and isotropic components to interpret the polarization observations of M31 (Beck et al. 2020). The ordered field can be generated from shear of differential rotation or compression of spiral arm shocks (Jaffe 2019). However, there are no indications of such large-scale structures inside the ring from radio observations. We therefore only consider the isotropic random magnetic fields. Since the path length along different lines of sight toward the ring is similar, the synchrotron emission from random fields is independent of azimuthal angle. The azimuthal variation of non-thermal emission is thus caused by the large-scale regular field.

The intensity of synchrotron radiation can be derived using the above equations. The inclination angle $i = 75^{\circ}$ (Chemin et al. 2009) is used. The pitch angle is uncertain. Berkhuijsen et al. (2003) obtained a pitch angle of $\theta = -13^{\circ} \pm 7^{\circ}$ by analyzing polarization observations at 6 cm. Fletcher et al. (2004) found a radial change of the pitch angle from -17° to -8° based on multi-frequency polarization observations. However, the recent analysis by Beck et al. (2020) suggested large variation of the pitch angle versus azimuthal angle for the total

magnetic field including both regular and ordered components.

Assuming a constant pitch angle for M31, we would expect the separation of the two intensity peaks to be 180° , which is clearly inconsistent with the observed profile shown in Figure 8. This implies variations of the pitch angle. We thus propose a simple model of the large-scale magnetic field with a pitch angle of -15° for the azimuthal angle range of 0° – 190° , and a pitch angle of 30° for the azimuthal angle range of 190° – 360° . Additionally, we impose a factor of 0.95 for the ratio of the two peaks, which we attribute to the variation of cosmic ray electron density. The resulting profile is presented in Figure 8, which reasonably reproduces the observed profile. A more sophisticated magnetic field model requires both total intensity and polarization observations at multiple frequencies, which is beyond the scope of this paper.

4. Summary

We obtain total intensity images from the FAST legacy survey of M31, covering the frequency range of 1.0–1.5 GHz. By averaging over all frequency channels, we derive the intensity image at 1.248 GHz, which reveals weak extended emission outside the ring because of the high sensitivity. The image allows us to analyze the propagation of cosmic ray electrons and the magnetic field configuration.

The main conclusions are as follows:

1. We derive a scale length of about 2.7 kpc for the cosmic ray electrons by fitting the radial profile of non-thermal emission. In comparison with the scale length at higher frequencies, we find that cosmic ray electrons propagate mainly through diffusion.
2. The spectral index of the total intensity in the ring, calculated from TT-plots, clearly exhibits azimuthal variation, which can be attributed to spectral variation of synchrotron radiation. The variation is likely caused by changes in star formation rates along the ring.
3. The weak emission detected outside the ring might not be extended emission from the ring. Instead, it might be produced by star-forming activities.
4. The azimuthal profile of non-thermal emission can be reproduced with an axisymmetric large-scale magnetic field pattern with pitch angle varying with azimuthal angle, indicating that the magnetic field configuration in the ring is complicated.

Acknowledgments

W.Z. and X.S. are supported by the National SKA Program of China (grant No. 2022SKA0120101). We thank the anonymous referee for comments that

helped improve the paper. We thank Dr. Fatemeh Tabatabaei for providing the thermal fraction map of M31.

ORCID iDs

Wenjun Zhang <https://orcid.org/0000-0003-3136-7756>

Xiaohui Sun <https://orcid.org/0000-0002-3464-5128>

References

- Battistelli, E. S., Fatigoni, S., Murgia, M., et al. 2019, ApJL, 877, L31
- Beck, R., Berkhuijsen, E. M., Gießbübel, R., & Mulcahy, D. D. 2020, A&A, 633, A5
- Beck, R., Berkhuijsen, E. M., & Hoernes, P. 1998, A&As, 129, 329
- Beck, R., & Krause, M. 2005, AN, 326, 414
- Berkhuijsen, E. M., Beck, R., & Hoernes, P. 2003, A&A, 398, 937
- Berkhuijsen, E. M., Beck, R., & Tabatabaei, F. S. 2013, MNRAS, 435, 1598
- Berkhuijsen, E. M., Wielebinski, R., & Beck, R. 1983, A&A, 117, 141
- Chemin, L., Carignan, C., & Foster, T. 2009, ApJ, 705, 1395
- Condon, J. J., Cotton, W. D., Greisen, E. W., et al. 1998, AJ, 115, 1693
- Devereux, N. A., Price, R., Wells, L. A., & Duric, N. 1994, AJ, 108, 1667
- Fatigoni, S., Radiconi, F., Battistelli, E. S., et al. 2021, A&A, 651, A98
- Fletcher, A., Berkhuijsen, E. M., Beck, R., & Shukurov, A. 2004, A&A, 414, 53
- Gil de Paz, A., Boissier, S., Madore, B. F., et al. 2007, ApJS, 173, 185
- Han, J. L., Beck, R., & Berkhuijsen, E. M. 1998, A&A, 335, 1117
- Jaffe, T. R. 2019, Galaxies, 7, 52
- Jarrett, T. H., Chester, T., Cutri, R., Schneider, S. E., & Huchra, J. P. 2003, AJ, 125, 525
- Jiang, P., Tang, N.-Y., Hou, L.-G., et al. 2020, RAA, 20, 064
- Jiang, P., Yue, Y., Gan, H., et al. 2019, SCPMA, 62, 959502
- Mulcahy, D. D., Fletcher, A., Beck, R., Mitra, D., & Scaife, A. M. M. 2016, A&A, 592, A123
- Nieten, C., Neininger, N., Guélin, M., et al. 2006, A&A, 453, 459
- Pietsch, W., Freyberg, M., & Haberl, F. 2005, A&A, 434, 483
- Pooley, G. G. 1969, MNRAS, 144, 101
- Rice, W., Lonsdale, C. J., Soifer, B. T., et al. 1988, ApJS, 68, 91
- Sarkisyan, A., Sholukhova, O., Fabrika, S., et al. 2020, MNRAS, 497, 687
- Smith, M. W. L., Eales, S. A., Gomez, H. L., et al. 2012, ApJ, 756, 40
- Stanek, K. Z., & Garnavich, P. M. 1998, ApJL, 503, L131
- Sun, X.-H., Gao, X.-Y., Reich, W., et al. 2022, RAA, 22, 125011
- Sun, X.-H., Meng, M.-N., Gao, X.-Y., et al. 2021, RAA, 21, 282
- Tabatabaei, F. S., Berkhuijsen, E. M., Frick, P., Beck, R., & Schinnerer, E. 2013, A&A, 557, A129
- Turtle, A. J., Pugh, J. F., Kenderdine, S., & Pauliny-Toth, I. I. K. 1962, MNRAS, 124, 297
- Viaene, S., Fritz, J., Baes, M., et al. 2014, A&A, 567, A71

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv — Machine translation. Verify with original.