

The transition to compulsion in addiction: insights from personality traits, social factors, and neurobiology

Authors: Haodong Su, Tongtong Ye, Chunyan Hu, Chunyan Hu

Date: 2024-05-20T00:00:00+00:00

Abstract

Compulsion stands as a central symptom of drug addiction; however, only a small fraction of drug users exhibit compulsive characteristics. Differences observed in Sign-trackers (ST) and Goal-trackers (GT) during Pavlovian conditioning may shed light on individual variances in drug addiction. Here, we focus on the behavioral attributes, formation processes, and neural mechanisms underlying ST and how they drive addiction towards compulsivity in humans. We will explore addiction from three interconnected levels: individual personality traits, social factors, and neurobiology. Furthermore, we distinguish between the processes of sensitization and habituation within ST. These nuanced distinctions across various aspects of addiction will contribute to our understanding of the addiction development process and the formulation of targeted preventive strategies.

Full Text

Preamble

The Transition to Compulsion in Addiction: Insights from Personality Traits, Social Factors, and Neurobiology

Authors: Haodong Su^{1,2,#}, Tongtong Ye^{1,2,#}, Chunyan Hu^{1,*}

¹ College of Humanities, Anhui Science and Technology University, Chuzhou, China

² Psychological Education Research Department, Anhui Science and Technology University, Chuzhou, China

Author Contribution Statement

Tongtong Ye: Conceptualization and research design; manuscript writing and revision

Haodong Su: Conceptualization and research design; manuscript writing and revision

These authors contributed equally to this work and should be considered co-first authors

- Corresponding author: Chunyan Hu (E-mail: 38375920@qq.com)

Abstract

Compulsion stands as a central symptom of drug addiction; however, only a small fraction of drug users exhibit compulsive characteristics. Differences observed in Sign-trackers (ST) and Goal-trackers (GT) during Pavlovian conditioning may shed light on individual variances in drug addiction. Here, we focus on the behavioral attributes, formation processes, and neural mechanisms underlying ST and how they drive addiction towards compulsivity in humans. We will explore addiction from three interconnected levels: individual personality traits, social factors, and neurobiology. Furthermore, we distinguish between the processes of sensitization and habituation within ST. These nuanced distinctions across various aspects of addiction will contribute to our understanding of the addiction development process and the formulation of targeted preventive strategies.

Keywords: Sign-tracker; Goal-tracker; Drug addiction; Individual differences

1. Introduction

Drug addiction represents a pressing global challenge in contemporary society, with approximately 243 million people worldwide grappling with substance abuse. This issue is accompanied by escalating societal costs, including increased healthcare expenditures, diminished productivity, and a surge in crime rates (Veber & Weidemann, 2018). The core symptom of drug addiction lies in compulsive drug use behavior, where individuals persistently seek and consume drugs despite severe negative consequences (Zou et al., 2017).

However, a key issue in addiction is that not everyone transitions from recreational, controlled drug use to uncontrolled, compulsive drug use. Only a small minority—approximately 15%-20% of individuals—cannot flexibly adjust their behavior. This implies that there are significant individual differences in the process of transitioning towards addiction (Volkow & Morales, 2015). Susceptibility factors propel individuals from initial drug use to maintenance and the development of addiction, and recognizing these susceptibility factors is crucial for addiction prevention (Moggi, 2018).

In the classical Pavlovian conditioning paradigm, the identification of Sign-tracker (ST) and Goal-tracker (GT) as two distinct phenotypes provides crucial

insights into understanding the driving forces behind individual susceptibility to addiction. Notably, these phenotypes may reflect inter-individual variations outlined in the incentive-sensitization theory. Specifically, these differences manifest as follows: (1) behaviorally, ST exhibits weakened inhibitory control, characterized by heightened novelty seeking and impulsivity traits, which may be associated with the two successive stages leading to compulsive transitions; (2) early negative experiences in ST, compared to GT, may be linked to the development of externalizing disorders; (3) the formation of ST possesses a distinct neural basis compared to GT. Given these points, it is imperative to incorporate the limited literature into a comprehensive framework encompassing psychological, social, and neurobiological factors. In this manuscript, we specifically review the unique aspects of ST across psychological, social, and neural domains, elucidating the driving roles of these three factors in the addiction process. Notably, we provide a more precise differentiation between the neural underpinnings of sensitization and habituation in addiction within the current manuscript.

2. The Incentive-Sensitization Theory: Sign-Tracker Versus Goal-Tracker

While consensus on the formation process of drug addiction remains elusive, the discourse has given rise to two classical theories, namely, the opponent process theory and the incentive-sensitization theory. The opponent-process theory posits that individuals initially experience the pleasurable effects of drug use (positive reinforcement). However, as tolerance, anxiety, and negative effects emerge, they experience relief from drug withdrawal symptoms (negative reinforcement) (Solomon & Corbit, 1974). In summary, the opponent-process theory suggests that addiction involves drug choices driven by negative states. However, this theory does not account for individual differences in addiction and cannot explain why individuals may exhibit strong drug motivations even when they are not in a withdrawal state (Kalivas & McFarland, 2003).

The incentive-sensitization theory provides a reasonable explanation for individual differences in addiction. According to this theory, addictive substances induce adaptive changes in the nervous system. These changes do not alter the pleasurable experience of the drug (“liking” the drug) but instead grant drug-related cues a strong motivational significance (“wanting” the drug). The incentive-sensitization theory successfully explains the separation of actions and intentions reported by many addicts and the fact that even after extended periods of abstinence, subtle environmental cues can trigger intense drug cravings (Hogarth & Hardy, 2018; Szumlinski et al., 2006). Individual differences in addiction stem from varying attributions, with the core characteristic of individual addiction being the attribution of reward to drug-related cues (Robinson & Berridge, 2000).

The phenotypes of Sign-trackers (ST) and Goal-trackers (GT) observed in the classic Pavlovian conditioned approach (PCA) paradigm, as well as the strong Pavlovian-instrumental transfer (PIT) effects displayed by ST in the

PIT paradigm, align perfectly with the incentive-sensitization theory. In PCA, neutral stimuli (e.g., a bell) repeatedly paired with unconditioned stimuli (US, e.g., food) become conditioned stimuli (CS), leading to conditioned responses (CR) in rodents (Pavlov, 2010). ST, as cue responders in the PCA, exhibit a stronger incentive quality for cues predicting rewards (e.g., lever press). They tend to approach these cues initially, and this cue's incentive can even persist in the absence of the US (Flagel & Robinson, 2017). GT, as goal responders, are less sensitive to cues predicting rewards. For instance, they may initially approach the food trough when it suggests a reward is available and do not invest as much time and effort in cue-related behaviors (Robinson & Flagel, 2009b) (Fig. 1).

In the PIT tests, ST individuals exhibit greater transfer effects, further highlighting the significant incentive value of CS for ST (Corbit & Balleine, 2011). As mentioned in the incentive-sensitization theory, this pathological incentive for cues may indeed be a driver of addiction. Correspondingly, research has reported that, compared to GT, ST individuals are more likely to predict compulsive drug use behavior in addiction. According to the 3-CRIT for judging compulsive drug use in animal models: (1) resistance to punishment (such as shock) during continued drug responses, (2) continued responses (drug craving) when the drug is unavailable, and (3) motivation to seek the drug under progressive ratio schedules (Deroche-Gamonet, Belin, & Piazza, 2004; Spanagel, 2017). Research findings indeed suggest that when subjected to relevant tests, ST individuals exhibit features of compulsion as outlined in the 3-CRIT criteria (Fitzpatrick, Geary, Creeden, & Morrow, 2019; Morrison, Bamkole, & Nicola, 2015; Saunders & Robinson, 2011). Despite reports of contradictory results, when the CS is devalued (CS associated with an aversive stimulus, such as lithium chloride), ST individuals do exhibit reduced responses to CS. However, this may be context-dependent, as ST can lose sensitivity to CS devaluation when the contextual environment is inconsistent, subsequently reverting to compulsive seeking (Amaya, Stott, & Smith, 2020; Derman, Schneider, Juarez, & Delamater, 2018). This aligns with real-world scenarios, such as the effectiveness of addiction treatment in controlled settings but susceptibility to cue-induced relapse in daily life. Furthermore, ST and drug abuse share common neural foundations (A. Tomie, Grimes, & Pohorecky, 2008). Thus, the potential differences between ST and GT phenotypes may indeed be related to individual variations in addiction.

3. Mapping Sign-Tracking and Goal-Tracking onto Human Behaviors

But another pivotal question revolves around whether these two phenotypes observed in animal models exhibit consistent or analogous patterns in humans. This consideration affects the potential for the susceptibility demonstrated within the ST phenotype to hold relevance for human translation. Previous research suggests that such a translation is not only possible but also reason-

able. In human studies, akin to findings in animal models, both ST and GT phenotypes have been identified. While intermediate types also exist (in line with animal models), a predominant bimodal distribution trend has emerged in human populations (Fig. 2). Moreover, investigations employing eye-tracking and functional magnetic resonance imaging (fMRI) techniques have provided evidence of distinct neural mechanisms for ST and GT in humans (Schad et al., 2020).

Some researchers have commented on the applicability of analogizing ST traits to humans. Similarly, studies have indicated that in humans, individuals who exhibit a focus on cues akin to ST may also be associated with more severe addiction and compulsivity (Colaizzi et al., 2020; Albertella et al., 2019).

Figure 2 [Figure 2: see original paper]. Distribution of Sign-Trackers and Goal-Trackers in Human and Animal Studies. a. In rodents, Sign-Trackers and Goal-Trackers exhibit a nearly symmetrical distribution (Saunders & Robinson, 2011). b. In human studies, Sign-Trackers and Goal-Trackers display a proportion similar to that observed in animal research (Schad et al., 2020).

In summary, it is appropriate to translate ST and GT from animal models to human research. Moreover, the behavioral characteristics exhibited in ST can help in understanding susceptibility factors in human addiction. There is increasing evidence that substance addiction is associated with habitual behavioral patterns. Therefore, in many studies, the behavioral characteristics of ST are referred to as “model-free” habitual behavioral patterns, while GT represents “model-based” goal-directed behaviors (Kasal et al., 2021; Schad et al., 2020). Second-order schedules can be used to investigate drug-related cue-reinforced seeking responses, such as studying the transition from goal-directed to habitual seeking controlled by CS in substances like heroin, cocaine, and alcohol (Di Ciano & Everitt, 2005; Everitt & Robbins, 2000; Giuliano et al., 2018).

4. Sensitization and Habituation of Cues: A Psychological and Neurobiological Separation

However, current research often conflates cue incentive sensitization and habituation within ST as the same process. This review argues that these two phenomena are not equivalent and are distinct in both psychological processes and neural foundations. In ST, cue sensitization remains goal-directed behavior, determined by the high incentive value of cues. This excessive attention to cues may be related to attentional bias and reflect poor attentional control in individuals. Its neural basis may be associated with dopamine neurons projecting from the ventral tegmental area (VTA) and substantia nigra compacta (SNc) to forebrain targets like the nucleus accumbens (NAc), encoding motivational behaviors (Berridge & Robinson, 2016; Budygin et al., 2020; Koshy Cherian et al., 2017; Kutlu et al., 2022).

The habituation within ST is related to a weakening of behavioral inhibition in individuals, reflecting a lack of planning in behavior. For example, compared

to GT, ST individuals exhibit earlier and more frequent lever-pressing behavior (Flagel et al., 2010). This pattern of weakened inhibition primarily manifests as difficulties in top-down control, potentially associated with deficits in cholinergic modulation in the cortex, the gradual waning of cortical control over subcortical structures, and the shift from the ventral to dorsal striatum (from ventral striatum (VTS) and dorsal medial striatum (DMS) to dorsal lateral striatum (DLS)). It is worth noting the synaptic plasticity between the cortex and striatum, which is linked to the Dopaminergic neuronal projection from the midbrain to nucleus accumbens medium spiny neurons (MSN) (Barker et al., 2015; Giuliano, Belin, & Everitt, 2019; Pascoli et al., 2014; Pascoli, Turiault, & Luscher, 2011; Sarter & Phillips, 2018). This further emphasizes that learning associated with high-incentive cue associations may form the foundation for the transition to habitual behaviors. The failure of goal-directed control and the dominance of habitual behavior patterns may serve as the basis for the shift from controlled drug use to compulsive drug use (Luscher, Robbins, & Everitt, 2020).

Hence, the behavioral characteristics, underlying factors, and biological basis exhibited by ST seem to shed light on individual differences in human addiction. First, concerning behavioral characteristics, as mentioned earlier, ST individuals are more prone to cue sensitization, which may be associated with attentional biases. Conversely, habituation may be linked to poorer behavioral inhibition, a trait often found in individuals with impulsivity (Schettino et al., 2022).

Individuals with high impulsivity traits often exhibit reduced attentional capacity and behavioral inhibition, which has been associated with impulsive behaviors and various substance use disorders in human studies (Dent & Isles, 2014; Hildebrandt, Dieterich, & Endrass, 2021; Potvin et al., 2018). It is worth noting that, in terms of impulsive choice, there is no difference between ST and GT, suggesting that ST may only demonstrate poor behavioral inhibition without necessarily being sensitive to issues of time or probability discounting (Flagel et al., 2010). Furthermore, the use of selectively bred High-Responder (BHR) and Low-Responder (LHR) rats has shown differences in incentive attribution, with BHR primarily showing ST characteristics. This implies that novelty-seeking traits may be susceptibility factors for addiction (Flagel, Waselus, Clinton, Watson, & Akil, 2014). There is already considerable evidence linking novelty-seeking personality traits to substance addiction, including nicotine and cocaine (Belin & Deroche-Gamonet, 2012; Perkins et al., 2008). Secondly, the driving role of social factors in ST is noteworthy. In humans, substance abuse or addiction is often associated with a range of other behavioral syndromes collectively referred to as “externalizing disorders,” which includes impulsivity (Krueger, Markon, Patrick, Benning, & Kramer, 2007). These “externalizing disorders” are linked to early-life environmental stress, developmental experiences, and attachment relationships (typically with caregivers), with positive parenting being a protective factor against these disorders (Robinson & Flagel, 2009a). In summary, this comprehensive review delves into the intricacies of the transition from recreational drug use to the compulsive and uncontrollable phase, elucidating the driving forces at play across the interconnected dimensions of personality traits,

social factors, and neurobiology.

5.1 Novelty Seeking

Novelty seeking refers to the tendency to initiate behavior in response to new stimuli and potential rewards. This trait was initially introduced as part of the biopsychosocial model proposed by Cloninger and colleagues in 1993. This model is based on complex interactions among genetics, psychology, social influences, culture, and spiritual dimensions, categorizing an individual's personality traits into two major aspects: temperament and character, consisting of a total of seven sub-dimensions (Cloninger, Svrakic, & Przybeck, 1993). Cloninger posited that novelty seeking is a component of an individual's temperament module, representing a non-learned instinctual behavior characterized by a high motivation for new stimuli in the environment (Beckmann, Marusich, Gipson, & Bardo, 2011). As mentioned earlier, selectively bred rats that exhibit different responses to novel stimuli also show differences in attribution to incentives. Rats with high novelty-seeking tendencies often display ST in PCA, suggesting that the novelty-seeking trait might be one of the susceptibility factors in the transition to compulsive behaviors.

However, research on novelty seeking in addiction has yielded mixed results. In human studies, the novelty-seeking trait can predict susceptibility in the initial stages of self-administration and compulsive drug use. Novelty-seeking levels measured in early adulthood can serve as predictive factors for the abuse of substances such as alcohol, nicotine, cannabis, and various other substances (Belin & Deroche-Gamonet, 2012; Foulds, Boden, Newton-Howes, Mulder, & Horwood, 2017; Stautz & Cooper, 2013). But another animal study that aligns with the 3-CRIT suggested that the high novelty-seeking phenotype cannot predict binge-like drinking behavior in mice (Radwanska & Kaczmarek, 2012). It's worth noting that Belin et al.'s research may reveal a complex structure of the novelty-seeking trait, with its different dimensions being related to different aspects of addiction. As defined by Zuckerman and others, novelty seeking is "a personality trait characterized by a tendency to actively seek out new and exciting sensations, and a willingness to take physical, social, legal, and other risks for the sake of experiencing these novel sensations (Zuckerman, Kuhlman, Joireman, Teta, & Kraft, 1993)." Questionnaire measurements of novelty-seeking traits in human studies also encompass dimensions such as impulsivity, exploratory excitement, and disorderliness (Garcia, Lester, Cloninger, & Robert Cloninger, 2017).

Therefore, novelty-seeking measured in animal models may not capture all the features of human novelty-seeking, which could be a reason for the inconsistencies between human and animal research findings. This notion is supported by studies that investigate subtypes of novelty-seeking. Researchers have further subdivided high novelty-preferring rats (BHR) into high novelty preference (HNP) and low novelty preference (LNP) subtypes based on their choice preference in free-choice procedures. The results suggest that HNP may be a susceptibility factor for compulsive cocaine use, promoting the transition from cocaine

use to compulsive behavior (Belin, Berson, Balado, Piazza, & Deroche-Gamonet, 2011). These findings indicate that making fine distinctions in animal models may help us better understand the mechanisms underlying compulsive drug use in addiction.

Novelty-seeking may predict the transition from drug use to compulsive behavior due to two potential factors. From a neural mechanism perspective, novelty-seeking and sensitization in ST may share a common biological basis. For example, many addictive drugs lead to an increase in dopamine levels in the mesolimbic system. Alcohol has complex effects on gamma-aminobutyric acid (GABA) and glutamate receptors, resulting in rapid changes in dopamine levels in the NAc (Abrahao, Salinas, & Lovinger, 2017). Cocaine, as a potent stimulant, increases dopamine levels by blocking the reuptake of dopamine at neuronal terminals, while nicotine can directly depolarize dopamine neurons (Luscher & Ungless, 2006). Among the neurotransmitters associated with substance addiction, some have also been reported in studies related to novelty-seeking behavior. For example, Rohan et al. revealed that exposure to a new environment could activate neural pathways shared with addiction (Rohan, Lowen, Rock, & Andersen, 2021). Behaviorally, novelty-seeking may encompass characteristics such as poor attention and impulsivity, all of which fall under the concept of behavioral inhibition. These traits may include compulsive cue-seeking behavior seen in ST.

5.2 The Trait of Impulsivity

The trait of impulsivity is a complex, multidimensional construct that can conceptually be divided into two main components: impulsive behavior and impulsive choice. High impulsivity has been associated with a range of psychiatric disorders, including bipolar disorder, attention-deficit/hyperactivity disorder, and borderline personality disorder, among others (Bayes, Parker, & Paris, 2019; Berg, Latzman, Bliwise, & Lilienfeld, 2015). Within the realm of personality traits, impulsivity is generally defined as “the tendency to make rapid, unplanned, or reward-driven responses to internal or external stimuli without adequately considering the potential consequences for oneself and others (Moeller, Barratt, Dougherty, Schmitz, & Swann, 2001).” The description of the impulsivity trait in humans aligns with the characteristics of ST. As previously discussed, we distinguished between sensitization and habituation in the formation of ST behavior, which is consistent with the attention deficits and lack of planning features observed in human impulsivity traits.

Firstly, there's sensitization. ST attributes the incentive value to cues rather than drugs, and even when the reward is lost (e.g., food), it cannot stop the attention to cues. For example, raccoons may become fixated on biting coins (US) and miss out on food (CS) rewards (Breland & Breland, 1961; Bucker & Theeuwes, 2017). Attention control deficits may be related to this behavioral pattern. For example, ST typically perform poorly in sustained attention tasks (SAT) (Pitchers, Wood, Skrzynski, Robinson, & Sarter, 2017). The attention

capture related to cue rewards may form the basis for incentive attribution, and this cue sensitization may predict compulsive behavior in addiction. In human studies, cue-reward-related attention capture has been found to predict an individual's addiction and compulsive behavior, and is associated with the severity of compulsive behavior. This attention bias may be the basis for cognitive inflexibility patterns (Albertella et al., 2019; Albertella et al., 2020).

Compulsive behavior can be understood as a focus on the immediate action despite adverse consequences, losing the association between behavior and consequences. Attentional narrowing may be a precursor to transitioning to these compulsive traits, but at this stage, it is still goal-oriented (Brown, Duka, & Forster, 2018). Therefore, the core features of compulsive use in addiction are the excessive habitual behavior (lack of planning) following cue sensitization and the inability to break free from habit-based control during drug use (Fig. 4 [Figure 4: see original paper]). Habitual behavior is based on stimulus-response associations and typically occurs after extensive training. Once habits are established, they require fewer cognitive resources, making the response often independent of outcome value, triggered by specific cues or stimuli (automatic attention to cues) (Dickinson, 1985). The reduced capacity for top-down behavioral inhibition observed in ST may make it more prone to habit formation. In other words, in the competition between goal-directed and habitual behavioral patterns, ST individuals may be more inclined to have their behavior dominated by habit-based patterns (Campus et al., 2019).

Therefore, the impulsivity trait, characterized by impulsiveness or a lack of goal-directedness, may link habit formation with compulsive drug-taking behavior. It is this deficient top-down control that makes it difficult to quit drug use and shift back to goal-directed behavior. Normally, when there is reward devaluation, individuals quickly revert to goal-directed behavior. However, individuals with high impulsivity traits, due to poor attention control and weakened top-down behavioral inhibition, are prone to remain sensitized to cues, maintaining habitual attention to cues and subsequent behavioral responses. In drug use, this habit-dominated behavioral pattern leads to the transition to compulsive drug-taking and eventually evolves into uncontrolled drug-seeking behavior.

6. Social Factors in Addiction

High impulsivity and addiction both fall under the category of externalizing disorders, and the development of such externalizing disorders is influenced by an individual's early life experiences. Therefore, to further understand the origins of impulsivity, attention deficits, and novelty-seeking behavior, we will delve into the psychodynamic perspective to comprehend the psychological processes involved in the transition towards compulsive drug use during an individual's drug use development.

Adverse experiences, the establishment of attachment relationships, and exposure to stress during an individual's early developmental stages may influence

susceptibility to compulsive behavior in addiction (Alvarez-Monjaras, Mayes, Potenza, & Rutherford, 2019). It's important to note that these adverse experiences, attachment, and stress are not isolated factors; they often interconnect and mutually affect each other. For instance, early separation from caregivers during childhood is both an adverse experience and an example of an insecure attachment relationship. While these factors overlap, they also have distinctions. For instance, early adverse experiences may be related to the dopamine neural system, attachment relationships emphasize social interactions and may be related to the oxytocin system, and stress may be associated with an individual's hypothalamic-pituitary-adrenal (HPA) axis function (Kim et al., 2017).

6.1 Early Adverse Experiences

Early adverse experiences in rodents are typically characterized by disrupted caregiving behaviors, such as premature separation from the mother. Research has shown that animals experiencing such adverse experiences tend to exhibit high novelty-seeking behaviors in adulthood, consistent with what was mentioned earlier. In addition to showing high seeking traits, they also demonstrate greater sensitivity to addictive substances (Brake, Zhang, Diorio, Meaney, & Gratton, 2004; Kim et al., 2017; Zimmerberg & Shartrand, 1992). These early adverse experiences may physiologically impact the activity of dopamine neurons in the brain. For instance, adolescent rats exposed to early maternal separation exhibit alterations in baseline dopamine levels in the striatum and prefrontal cortex (Llorente et al., 2010). Importantly, there are also changes in the levels of dopamine release induced by stimuli. For instance, an enhanced dopamine release in response to rewarding stimuli has been observed in the VST and hypothalamus (Arborelius & Eklund, 2007). The altered patterns of reward and motivation-related brain dopamine neuron activity resulting from these early adverse experiences have also been consistently observed in human research (Pruessner, Champagne, Meaney, & Dagher, 2004). These changes in dopamine neuron activity patterns may be related to the attribution of incentive to subsequent drug-related cues (Colaizzi et al., 2023; Hynes et al., 2018). Additionally, early adverse experiences influence the development of compulsive behavioral traits during adolescence (Brydges, Holmes, Harris, Cardinal, & Hall, 2015). All of these findings suggest that early adverse experiences in individuals may drive drug use behaviors in addiction towards compulsivity.

6.2 Attachment Relationships

Secure attachment relationships have a protective effect against substance abuse in adulthood (Arthur Tomie, 2018). The positive social interactions between individuals and their caregivers contribute to the development of executive functions and self-regulation. In contrast, a lack of soothing and positive attachment experiences can hinder the establishment of these functions, eventually manifesting as impulsive traits, particularly attention deficits and weakened behavioral inhibition within an individual's personality. For instance, rodent

models have demonstrated that rats with insufficient early social interaction experiences (poor attachment experiences) exhibit significant arousal towards reward-related cues, leading to a loss of behavioral inhibition (Lomanowska et al., 2011). On the other hand, the HPA axis is believed to directly influence the behavior of ST and GT within the PCA, with ST showing greater cortisol release during single PCA sessions (Beckmann & Bardo, 2012; Flagel, Akil, & Robinson, 2009). However, positive social interaction experiences can reduce HPA axis activity, inhibiting ST' s seeking responses to cues within the PCA (Beckmann & Bardo, 2012). Recent research suggests that attachment and addiction may share a common neural basis. Therefore, individuals with maladaptive attachment relationships may have their neural systems perpetuating the development of addictive behaviors (Burkett & Young, 2012; Rutherford, Williams, Moy, Mayes, & Johns, 2011).

6.3 Stress

Early life stress events can lead to maladaptive tendencies, such as children facing extremely harsh parenting styles often exhibiting higher levels of impulse control disorders and externalizing disorders (Robinson & Flagel, 2009a). In rodent studies, rats subjected to stress due to social isolation exhibited more cue-induced sensitization characteristics and showed heightened locomotor reactivity to novel stimuli during adolescence (Baarendse, Limpens, & Vanderschuren, 2014; Lomanowska et al., 2011). Furthermore, the HPA axis is involved in a series of cascading neurotransmitter and hormone regulatory processes related to stress. In chronic stress environments, sustained activation of the HPA axis can enhance extracellular DA release in the striatum, indirectly impacting neural pathways encoding motivation and reward within the brain (Nikulina, Lacagnina, Fanous, Wang, & Hammer, 2012; Wang et al., 2005). Overall, these findings suggest that early life stress events may shape susceptibility traits for compulsive behavior in addiction.

In summary, whether it' s attachment relationships or stressful events, their influence on individuals during early life is enduring and subtle, and these developmental social factors always impact inhibitory functions behaviorally. Physiologically, it is invariably associated with the neural foundations that encode motivation and goal-directed or habitual behaviors. From a psychological process perspective, it indeed drives the formation of impulsive behaviors, laying the foundation for the transition of drug use in addiction towards compulsivity.

7.1 Neural Basis of Cue Sensitization

In the preceding sections, we have discussed how personality traits and social factors drive the transition from drug use to compulsive drug use. However, another crucial question is how the brain' s relevant neural systems participate in the transition to addiction. First and foremost is the sensitization to cues, where the encoding of cues in a highly motivated state may rely on the phasic release of dopamine in the NAc. The use of fast-scan cyclic voltammetry allows

for the measurement of dopamine changes at a sub-second timescale. The results indicate that differences in dopamine responses between reward and prediction occur only in the ST system, with no significant changes observed in the GT system (Flagel et al., 2011). This suggests that the phasic changes in dopamine in the NAc may not encode the traditional “reward prediction error hypothesis,” which relates to the encoding of prediction and actual reward discrepancies. Instead, it may encode the incentive value of cues and is mediated by dopamine in the Nucleus Accumbens core (NAcc) (Dalley et al., 2005; Di Ciano, Cardinal, Cowell, Little, & Everitt, 2001; Montague, Dayan, & Sejnowski, 1996). This proposition was further confirmed by systemic administration of flupenthixol, a non-selective dopamine antagonist, which showed that blocking dopamine had an impact on learning in the ST system within the PCA, while the GT system remained largely unaffected (Flagel et al., 2011).

Furthermore, the sustained incentive for cues in the ST system is also associated with the dopamine content in the NAcc. During cue-induced reinstatement tests, the ST system exhibits higher responsiveness compared to the GT system. Blocking dopamine in the NAcc can attenuate this response in the ST system, leading to behavior similar to that of the GT system (Saunders, Yager, & Robinson, 2013). The upregulation of the dopamine transporter (DAT) in the ST system may be related to the increased dopamine concentration in the Nucleus Accumbens core (NAcc) induced by drug use. When dopamine is released from neurons into the extracellular space, DAT on the synaptic surface plays a primary role in clearing and recycling excess dopamine. The longer dopamine stays in the extracellular space, the more it interacts with neighboring neurons, leading to dysregulation in the system (Singer et al., 2016). Therefore, compared to the GT system, dopamine reuptake in the extracellular space of the NAcc occurs more rapidly in the ST system. However, some addictive substances can increase synaptic dopamine levels by blocking and inhibiting DAT. For example, directly injecting amphetamine into the NAcc can slow down the dopamine reuptake process in the ST system (Singer et al., 2016). The synaptic dopamine that is not cleared primarily binds to D1 and D2 receptors and plays a significant role in encoding the incentive salience (Flagel et al., 2016). Furthermore, in Long-Evans rats, the high expression of DAT phenotype can predict cocaine-like addictive behaviors (Yamamoto et al., 2013). Therefore, these results suggest that the transmission of dopamine ability in the NAcc may contribute to the encoding of incentive salience in the ST system and this may also be a susceptibility feature for the transition to compulsive drug use behavior.

The formation of the ST is therefore dependent on dopamine, and the pattern of dopamine release in the NAcc is crucial for the reward learning process. The dopamine neurons in the NAcc primarily originate from dense projections of the midbrain VTA and SNc, and this mesolimbic dopamine neural circuit plays a significant role in encoding cue-induced motivation (Volkow, Wise, & Baler, 2017). It is noteworthy that dopamine neurons output from the VTA and SNc exchange information in the striatum through MSN and glutamatergic neurons

from the cortex, forming a circuit with a spiral-like structure reminiscent of a serial loop (Haber, Fudge, & McFarland, 2000). This also underscores that changes in the neural system sensitized to cue-induced stimulation will further impact the cortical structure and ventral striatum regulation of target behaviors.

Another crucial brain region related to cue-induced motivation is the hippocampus. The hippocampus is involved in various types of memory, especially context-related cues (Burgess, Maguire, & O'Keefe, 2002; Martin & Clark, 2007). Anatomically, the hippocampus can be divided into the ventral hippocampus (VHipp) and the dorsal hippocampus (DHipp). VHipp projects glutamatergic neurons to the NAc and is thus potentially involved in cue-driven motivation through this pathway. For instance, it has been shown that damage to VHipp can impact the concentration of dopamine in the NAc and inhibit cue-seeking behaviors in the ST within the PCA (Fitzpatrick, Creeden, Perrine, & Morrow, 2016). In summary, the hippocampus, as a regulator of contextual or spatial stimuli, may play a crucial role in the motivational effects of cues on drug seeking (Selden, Everitt, Jarrard, & Robbins, 1991).

The basolateral amygdala (BLA) also plays a crucial role in cue-induced motivation for drug seeking, and it is the BLA-NAc connectivity loop that is particularly essential. Selective damage to either the BLA or NAc, effectively disconnecting the two, has no impact on self-administration behavior for cocaine but impairs cocaine seeking in secondary reinforcement procedures (Di Ciano & Everitt, 2004). This suggests that there are distinctions at the neural circuit level between drug seeking and drug taking. In summary, maintaining cue-controlled drug seeking requires the involvement of the BLA-NAc connection, and this connection forms a serial loop circuit with the involvement of the prefrontal cortex and striatum.

7.2 The Top-Down Loss of Control and the Transition to Dorsal Striatum Control Pattern

The projection of dopamine neurons from the midbrain to the striatum mediates the sensitization of cue-induced motivation, and these neural system's plastic changes also lay the foundation for subsequent habitual behavioral patterns. However, the core of transitioning to compulsive seeking in addiction is the inability to break free from habit-dominated behavioral patterns. Therefore, the neural basis of habituation likely represents a core susceptibility mechanism for compulsive seeking. In cue-induced motivation, an individual's behavior is still in a state of balance between goal-directed and habitual actions. As mentioned earlier, within the ST, not only is there sensitization to cues but also changes driven by habitual behaviors. Unlike goal-directed behavior (model-based), this form of habit-driven behavior (model-free) is considered a crucial foundation for the transition to compulsive seeking (Luscher et al., 2020). This gradual transition to a habitual pattern of behavior, which remains resistant to punishment, may have its neural basis in the difficulty of top-down control from cortical regions to subcortical structures and the dominance of the DLS

in behavior control (Furlong, Jayaweera, Balleine, & Corbit, 2014; Hyman & Malenka, 2001; Smith & Laiks, 2018).

It is widely recognized that the PFC plays a crucial role in maintaining goal-directed behavior. However, many addictive substances can impair PFC function. For example, in individuals with alcohol addiction, brain regions associated with goal-directed behavior (vmPFC and ventral striatum) have been found to be less active compared to control groups, while regions associated with habit (e.g., the nucleus accumbens shell, equivalent to the dorsolateral striatum in rodents) show increased activity (Ersche, Williams, Robbins, & Bullmore, 2013). Although this frontal lobe damage is related to substance intake, individual susceptibility factors may also play a role. Studies involving drug-addicted individuals and their non-addicted siblings have found that both groups exhibit impairments in frontal lobe function. High impulsivity traits are often associated with difficulties in frontal lobe-mediated behavioral inhibition, suggesting that impulsivity traits in personality may serve as susceptibility factors for addiction (Ersche et al., 2011; Ersche et al., 2012). This impairment of PFC function can also affect an individual's executive functions and lead to decision-making deficits, which may, in turn, drive the development of addiction (Bechara et al., 2001). Impairments in executive functions can result in poor inhibition of habitual behaviors, thereby prioritizing the output of habitual behaviors in response to cues (Hardwick, Forrence, Krakauer, & Haith, 2019).

As the glutamatergic neurons from the prefrontal cortex interact with the dopamine neurons projecting from the VTA and SNc to the striatum in a serial loop circuit, the balance between goal-directed and habitual behavior patterns is likely disrupted, leading to the establishment of habit-dominated behavior patterns (Haber et al., 2000) (Fig. 3). This shift in behavior patterns is thought to be the result of a transition from the ventral striatum to the dorsal striatum.

The important roles of the DMS and DLS in compulsive drug use have been confirmed in both human and animal research. For example, the anterior part of the DLS (aDLS) plays a prominent role in the transition to compulsive seeking; functional magnetic resonance imaging (fMRI) studies in humans have shown increased activation in the DMS when individuals who engage in recreational drug use see drug-related cues, while addicted individuals show enhanced DLS activity (Everitt & Robbins, 2005, 2016; Zhou et al., 2019). The “seeking-taking” chained procedure distinguishes compulsive seeking and taking of drugs. In this paradigm, animals must perform an action (such as pressing a lever) to “seek” another task that allows them to “take” the drug. After several weeks of training, a shift in control from the DMS to the DLS was observed in animals that were insensitive to reinforcement devaluation (Corbit, Nie, & Janak, 2012). This indicates a shift from goal-directed to habit-dominated behavior patterns in this paradigm. Furthermore, in the same training procedure, when rats exhibited habit-dominated behavior, inhibiting DLS activity forced the habit system offline, and rats again exhibited sensitivity to reinforcement devaluation

(Zapata, Minney, & Shippenberg, 2010). In summary, these results suggest that in the transition to compulsive seeking, seeking behavior dependent on the DMS gradually becomes dominated by habitual seeking responses dependent on DLS activity.

Therefore, as it stands, the significant individual differences observed in addiction are not only related to sensitization to cues during drug use but, more importantly, to an overreliance on the DLS during the drug use process and an inability to break free from habit-dominated behavior driven by the DLS. When faced with punishment, individuals who cannot disengage from the DLS will exhibit characteristics of compulsive drug use (Fig. 4). The dopaminergic neurons projecting from the VTA to the NAc are associated with GABAergic MSN in the NAc region. GABAergic MSN neurons project back to subcortical areas, inhibiting dopamine neurons located outside the VTA. These dopamine neurons subsequently project to the NAc shell. This circuit forms a feedback mechanism, and after several iterations, it reaches the SNc, which sends dopamine neuron outputs to the DLS. PFC regulates this serial loop circuit by sending powerful glutamatergic neurons to both the NAc and DLS (Haber et al., 2000; Ikemoto, 2007). Indeed, during the transition to compulsive drug use, the PFC gradually loses control over this serial loop circuit. In the competition between goal-directed and habit-directed behaviors, control is gradually shifted to the habit system dominated by the DLS. This shift in control dynamics is a crucial aspect of the development of compulsive drug-seeking behavior.

8. Preliminarily Exploring the Synergistic Dynamics of Personality Traits, Social Factors, and Neurobiology

In the preceding sections, we have elaborated on the potential driving roles of Personality Traits, Social Factors, and Neurobiology in the individual's progression towards addictive behavior. However, as mentioned, these three factors do not operate in isolation; rather, they interact in intricate ways to propel individuals towards compulsive drug-seeking behaviors. Therefore, a preliminary discussion on the complex interactions among these three factors would aid in a better understanding of the driving forces behind individual progression towards addiction.

Although individuals with high novelty-seeking and impulsivity tendencies may exhibit similar behavioral characteristics, not all of them will manifest susceptibility to addiction. This suggests the existence of other significant driving factors. It is noteworthy that environmental pressures (herein referred to as broader social factors), especially negative experiences in early life, may promote individuals to display externalizing disorders, characterized by elevated novelty-seeking and impulsivity. These negative experiences include early maternal deprivation, establishment of insecure social relationships, and familial adversities.

Overall, social environmental factors can influence individual behavioral traits,

and this influence may be enduring and stable. However, current research on the effects of early environmental pressures on these behaviors is predominantly conducted using rodent models, despite considerable similarities to human-related research and its implications for addiction susceptibility. Considering the ecological validity and translational potential of research findings, we recommend that future studies involving human subjects may appropriately consider the following methods: (1) Large-scale cross-lagged panel studies: By measuring the impact of stress and negative experiences, such as social factors, on individual novelty-seeking and impulsivity tendencies at different time points, explore the causal relationship between these factors and analyze their cross-lagged effects to determine the direction of the influence of these social factors on individual novelty-seeking and impulsivity traits; (2) Machine learning: Machine learning offers significant advantages in handling large-scale data. Using machine algorithms such as random forests, neural networks, etc., these algorithms can process extensive data and automatically detect nonlinear relationships between variables, potentially revealing underlying patterns of how social factors influence individual novelty-seeking and impulsivity traits.

The inclination towards novelty-seeking, which delineates an individual's propensity for heightened behavioral responses to novel stimuli and potential rewards, may signify an inherent neurobiological predisposition towards sensitization to cues. For instance, neural substrates involving DA neurons encoding circuitry originating from the VTA and SNc, projecting to forebrain targets such as the NAc, are implicated. Previous studies have also indicated that novelty-seeking may engage shared neural pathways with addiction (Berridge & Robinson, 2016; Budygin et al., 2020; Koshy Cherian et al., 2017; Kutlu et al., 2022; Rohan et al., 2021).

Furthermore, the lack of planning or goal-directedness within the trait of impulsivity may link habits with compulsive drug-seeking behaviors. It is precisely this top-down mediated lack of control that renders abandonment of drug use and restoration of goal-directedness challenging. Impulsivity thus emerges as a potent driving force for individuals transitioning into a habitual dominance of compulsive drug-seeking patterns (Luscher et al., 2020). These behavioral characteristics are associated with corresponding neurobiological substrates and may further shape and strengthen the neural connections associated with them.

As mentioned earlier, social environmental factors may influence individual novelty-seeking and impulsivity traits, and this influence similarly extends to neurobiological factors. For instance, the hypothalamic-pituitary-adrenal (HPA) axis, widely implicated in stress and stress-related responses, is also believed to directly influence behaviors related to sensation seeking and goal-directed behavior (Arthur Tomie, Tirado, Yu, & Pohorecky, 2004). Therefore, in future research, investigators may consider employing methods such as biomarkers and neuroimaging techniques. These tools, utilized to measure biological and neural system responses, can offer insights into the impact of social environmental factors such as early maternal deprivation, establishment of insecure social rela-

tionships, and familial adversities on the neurobiology of addiction.

9. Concluding Summary

In fact, in the fifth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) by the American Psychiatric Association, there has been a gradual shift away from emphasizing drug dependence in the diagnostic criteria for substance use disorders. Instead, a specific emphasis has been placed on the compulsive features exhibited by individuals with addiction. This includes behaviors such as dedicating substantial time and financial resources to drug-seeking, neglecting fundamental activities (such as essential employment and basic social interactions), and persisting in drug intake despite experiencing negative physical and psychological consequences (American Psychiatric, 2022).

This underscores the increasing support for compulsive drug-seeking behavior as a core symptom in drug addiction, highlighting the significance of understanding the transition of individual behaviors into compulsive drug use. In particular: (1) before the manifestation of compulsive drug-seeking behavior, individuals retain flexibility in adjusting their expectations of goal values. However, following the onset of compulsive drug-seeking behavior, individuals exhibit diminished sensitivity to the devaluation of drugs. As discussed earlier, this observation may suggest the existence of distinct stages of sensitization and habituation during addiction; (2) precisely delineating these stages in the transition to addiction may aid in comprehending the complex and rich neuroscientific findings. Numerous neural circuits and neurotransmitters have been implicated in addiction, and distinguishing these results based on the temporal dimension of addiction progression may provide greater clarity.

In summary, our preceding discourse systematically reviewed the distinct phenotypes of Sign-Tracker (ST) and Goal-Tracker (GT) within the Pavlovian Conditioned Approach (PCA) paradigm. The emphasis was placed on highlighting that the observed differences between these phenotypes may unveil inherent susceptibility traits in addiction. Furthermore, our exploration revealed that several features associated with the ST phenotype may be intricately linked to compulsive drug use in addiction. Notably, these features include suboptimal top-down behavioral control, attentional deficits, and impulsive behavioral traits (Arthur Tomie, 2018). These traits are not limited to animal models; similar or analogous traits have been found in human studies, suggesting the potential for translating individual differences identified in animal models to human research (Colaizzi et al., 2020; Garofalo & di Pellegrino, 2015).

Therefore, we summarized the role of personality traits, specifically novelty seeking and impulsivity, in addiction. Overall, individuals with these two traits exhibit core features of the ST phenotype: poor attention and difficulties in behavioral inhibition. Furthermore, we ascertain that the behavior of ST may be associated with early environmental factors and events, such as early stress and attachment relationships. This observation potentially unveils the societal

driving factors contributing to the development of compulsive drug use in humans.

Finally, we summarized the current understanding of the neurobiological mechanisms underlying compulsive drug use. These studies suggest that the transition from controlled drug use to uncontrolled, compulsive drug-seeking behavior is rooted in habitual behavior (Barker et al., 2015; Feil et al., 2010; Furlong et al., 2014). These susceptibility traits for addiction are likely influenced by genetics and experiences. Therefore, future research in this field may employ genetic studies, longitudinal tracking studies, and cross-addiction spectrum studies to measure the predictive utility of these susceptibility traits in addiction and to separate the pharmacological effects of addictive substances.

Reference

- Abraham, K. P., Salinas, A. G., & Lovinger, D. M. (2017). Alcohol and the Brain: Neuronal Molecular Targets, Synapses, and Circuits. *Neuron*, 96(6), 1223-1238. doi:10.1016/j.neuron.2017.10.032
- Albertella, L., Le Pelley, M. E., Chamberlain, S. R., Westbrook, F., Fontenelle, L. F., Segrave, R., . . . Yucel, M. (2019). Reward-related attentional capture is associated with severity of addictive and obsessive-compulsive behaviors. *Psychol Addict Behav*, 33(5), 495-502. doi:10.1037/adb0000484
- Albertella, L., Le Pelley, M. E., Chamberlain, S. R., Westbrook, F., Lee, R. S. C., Fontenelle, L. F., . . . Yucel, M. (2020). Reward-related attentional capture and cognitive inflexibility interact to determine greater severity of compulsivity-related problems. *J Behav Ther Exp Psychiatry*, 69, 101580. doi:10.1016/j.jbtep.2020.101580
- Alvarez-Monjaras, M., Mayes, L. C., Potenza, M. N., & Rutherford, H. J. (2019). A developmental model of addictions: integrating neurobiological and psychodynamic theories through the lens of attachment. *Attach Hum Dev*, 21(6), 616-637. doi:10.1080/14616734.2018.1498113
- Amaya, K. A., Stott, J. J., & Smith, K. S. (2020). Sign-tracking behavior is sensitive to outcome devaluation in a devaluation context-dependent manner: implications for analyzing habitual behavior. *Learn Mem*, 27(4), 136-149. doi:10.1101/lm.051144.119
- American Psychiatric Association. (2022). *Diagnostic and Statistical Manual of Mental Disorders* (5th ed.).
- Arborelius, L., & Eklund, M. B. (2007). Both long and brief maternal separation produces persistent changes in tissue levels of brain monoamines in middle-aged female rats. *Neuroscience*, 145(2), 738-750. doi:10.1016/j.neuroscience.2006.12.007
- Baarendse, P. J., Limpens, J. H., & Vanderschuren, L. J. (2014). Disrupted

social development enhances the motivation for cocaine in rats. *Psychopharmacology (Berl)*, 231(8), 1695-1704. doi:10.1007/s00213-013-3362-8

Barker, J. M., Corbit, L. H., Robinson, D. L., Gremel, C. M., Gonzales, R. A., & Chandler, L. J. (2015). Corticostriatal circuitry and habitual ethanol seeking. *Alcohol*, 49(8), 817-824. doi:10.1016/j.alcohol.2015.03.003

Bayes, A., Parker, G., & Paris, J. (2019). Differential Diagnosis of Bipolar II Disorder and Borderline Personality Disorder. *Curr Psychiatry Rep*, 21(12), 125. doi:10.1007/s11920-019-1109-4

Bechara, A., Dolan, S., Denburg, N., Hinds, A., Anderson, S. W., & Nathan, P. E. (2001). Decision-making deficits, linked to a dysfunctional ventromedial prefrontal cortex, revealed in alcohol and stimulant abusers. *Neuropsychologia*, 39(4), 376-389. doi:10.1016/s0028-3932(00)00136-6

Beckmann, J. S., & Bardo, M. T. (2012). Environmental enrichment reduces attribution of incentive salience to a food-associated stimulus. *Behav Brain Res*, 226(1), 331-334. doi:10.1016/j.bbr.2011.09.021

Beckmann, J. S., Marusich, J. A., Gipson, C. D., & Bardo, M. T. (2011). Novelty seeking, incentive salience and acquisition of cocaine self-administration in the rat. *Behav Brain Res*, 216(1), 159-165. doi:10.1016/j.bbr.2010.07.022

Belin, D., Berson, N., Balado, E., Piazza, P. V., & Deroche-Gamonet, V. (2011). High-novelty-preference rats are predisposed to compulsive cocaine self-administration. *Neuropsychopharmacology*, 36(3), 569-579. doi:10.1038/npp.2010.188

Belin, D., & Deroche-Gamonet, V. (2012). Responses to novelty and vulnerability to cocaine addiction: contribution of a multi-symptomatic animal model. *Cold Spring Harb Perspect Med*, 2(11). doi:10.1101/cshperspect.a011940

Berg, J. M., Latzman, R. D., Bliwise, N. G., & Lilienfeld, S. O. (2015). Parsing the heterogeneity of impulsivity: A meta-analytic review of the behavioral implications of the UPPS for psychopathology. *Psychol Assess*, 27(4), 1129-1146. doi:10.1037/pas0000111

Berridge, K. C., & Robinson, T. E. (2016). Liking, wanting, and the incentive-sensitization theory of addiction. *Am Psychol*, 71(8), 670-679. doi:10.1037/amp0000059

Brake, W. G., Zhang, T. Y., Diorio, J., Meaney, M. J., & Gratton, A. (2004). Influence of early postnatal rearing conditions on mesocorticolimbic dopamine and behavioural responses to psychostimulants and stressors in adult rats. *Eur J Neurosci*, 19(7), 1863-1874. doi:10.1111/j.1460-9568.2004.03286.x

Breland, K., & Breland, M. (1961). The misbehavior of organisms. *American Psychologist*, 16(11), 681-684. doi:10.1037/h0040090

Brown, C. R. H., Duka, T., & Forster, S. (2018). Attentional capture by alcohol-related stimuli may be activated involuntarily by top-down search goals. *Psy-*

- chopharmacology (Berl), 235(7), 2087-2099. doi:10.1007/s00213-018-4906-8
- Brydges, N. M., Holmes, M. C., Harris, A. P., Cardinal, R. N., & Hall, J. (2015). Early life stress produces compulsive-like, but not impulsive, behavior in females. *Behav Neurosci*, 129(3), 300-308. doi:10.1037/bne0000059
- Bucker, B., & Theeuwes, J. (2017). Pavlovian reward learning underlies value driven attentional capture. *Atten Percept Psychophys*, 79(2), 415-428. doi:10.3758/s13414-016-1244-1
- Budygin, E. A., Bass, C. E., Grinevich, V. P., Deal, A. L., Bonin, K. D., & Weiner, J. L. (2020). Opposite Consequences of Tonic and Phasic Increases in Accumbal Dopamine on Alcohol-Seeking Behavior. *iScience*, 23(3), 100877. doi:10.1016/j.isci.2020.100877
- Burgess, N., Maguire, E. A., & O'Keefe, J. (2002). The human hippocampus and spatial and episodic memory. *Neuron*, 35(4), 625-641. doi:10.1016/s0896-6273(02)00830-9
- Burkett, J. P., & Young, L. J. (2012). The behavioral, anatomical and pharmacological parallels between social attachment, love and addiction. *Psychopharmacology (Berl)*, 224(1), 1-26. doi:10.1007/s00213-012-2794-x
- Campus, P., Covelo, I. R., Kim, Y., Parsegian, A., Kuhn, B. N., Lopez, S. A., . . . Flagel, S. B. (2019). The paraventricular thalamus is a critical mediator of top-down control of cue-motivated behavior in rats. *Elife*, 8. doi:10.7554/eLife.49041
- Cloninger, C. R., Svrakic, D. M., & Przybeck, T. R. (1993). A psychobiological model of temperament and character. *Arch Gen Psychiatry*, 50(12), 975-990. doi:10.1001/archpsyc.1993.01820240059008
- Colaizzi, J. M., Flagel, S. B., Gearhardt, A. N., Borowitz, M. A., Kuplicki, R., Zotev, V., . . . Paulus, M. P. (2023). The propensity to sign-track is associated with externalizing behavior and distinct patterns of reward-related brain activation in youth. *Sci Rep*, 13(1), 4402. doi:10.1038/s41598-023-30906-3
- Colaizzi, J. M., Flagel, S. B., Joyner, M. A., Gearhardt, A. N., Stewart, J. L., & Paulus, M. P. (2020). Mapping sign-tracking and goal-tracking onto human behaviors. *Neurosci Biobehav Rev*, 111, 84-94. doi:10.1016/j.neubiorev.2020.01.018
- Corbit, L. H., & Balleine, B. W. (2011). The General and Outcome-Specific Forms of Pavlovian-Instrumental Transfer Are Differentially Mediated by the Nucleus Accumbens Shell. *Journal of Neuroscience*, 31(33), 11786-11794. doi:10.1523/jneurosci.2711-11.2011
- Corbit, L. H., Nie, H., & Janak, P. H. (2012). Habitual alcohol seeking: time course and the contribution of subregions of the dorsal striatum. *Biol Psychiatry*, 72(5), 389-395. doi:10.1016/j.biopsych.2012.02.024

- Dalley, J. W., Laane, K., Theobald, D. E., Armstrong, H. C., Corlett, P. R., Chudasama, Y., & Robbins, T. W. (2005). Time-limited modulation of appetitive Pavlovian memory by D1 and NMDA receptors in the nucleus accumbens. *Proc Natl Acad Sci U S A*, 102(17), 6189-6194. doi:10.1073/pnas.0502080102
- Dent, C. L., & Isles, A. R. (2014). An overview of measuring impulsive behavior in mice. *Curr Protoc Mouse Biol*, 4(2), 35-45. doi:10.1002/9780470942390.mo140015
- Derman, R. C., Schneider, K., Juarez, S., & Delamater, A. R. (2018). Sign-tracking is an expectancy-mediated behavior that relies on prediction error mechanisms. *Learn Mem*, 25(10), 550-563. doi:10.1101/lm.047365.118
- Deroche-Gamonet, V., Belin, D., & Piazza, P. V. (2004). Evidence for addiction-like behavior in the rat. *Science*, 305(5686), 1014-1017. doi:10.1126/science.1099020
- Di Ciano, P., Cardinal, R. N., Cowell, R. A., Little, S. J., & Everitt, B. J. (2001). Differential involvement of NMDA, AMPA/kainate, and dopamine receptors in the nucleus accumbens core in the acquisition and performance of pavlovian approach behavior. *J Neurosci*, 21(23), 9471-9477. doi:10.1523/JNEUROSCI.21-23-09471.2001
- Di Ciano, P., & Everitt, B. J. (2004). Direct interactions between the basolateral amygdala and nucleus accumbens core underlie cocaine-seeking behavior by rats. *J Neurosci*, 24(32), 7167-7173. doi:10.1523/JNEUROSCI.1581-04.2004
- Di Ciano, P., & Everitt, B. J. (2005). Neuropsychopharmacology of drug seeking: Insights from studies with second-order schedules of drug reinforcement. *Eur J Pharmacol*, 526(1-3), 186-198. doi:10.1016/j.ejphar.2005.09.024
- Dickinson, A. (1985). Actions and Habits: The Development of Behavioural Autonomy. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 308(1135), 67-78. doi:10.1098/rstb.1985.0010
- Ersche, K. D., Barnes, A., Jones, P. S., Morein-Zamir, S., Robbins, T. W., & Bullmore, E. T. (2011). Abnormal structure of frontostriatal brain systems is associated with aspects of impulsivity and compulsivity in cocaine dependence. *Brain*, 134(Pt 7), 2013-2024. doi:10.1093/brain/awr138
- Ersche, K. D., Jones, P. S., Williams, G. B., Turton, A. J., Robbins, T. W., & Bullmore, E. T. (2012). Abnormal brain structure implicated in stimulant drug addiction. *Science*, 335(6068), 601-604. doi:10.1126/science.1214463
- Ersche, K. D., Williams, G. B., Robbins, T. W., & Bullmore, E. T. (2013). Meta-analysis of structural brain abnormalities associated with stimulant drug dependence and neuroimaging of addiction vulnerability and resilience. *Curr Opin Neurobiol*, 23(4), 615-624. doi:10.1016/j.conb.2013.02.017
- Everitt, B. J., & Robbins, T. W. (2000). Second-order schedules of drug reinforcement in rats and monkeys: measurement of reinforcing efficacy

and drug-seeking behaviour. *Psychopharmacology (Berl)*, 153(1), 17-30. doi:10.1007/s002130000566

Everitt, B. J., & Robbins, T. W. (2005). Neural systems of reinforcement for drug addiction: from actions to habits to compulsion. *Nat Neurosci*, 8(11), 1481-1489. doi:10.1038/nn1579

Everitt, B. J., & Robbins, T. W. (2016). Drug Addiction: Updating Actions to Habits to Compulsions Ten Years On. *Annu Rev Psychol*, 67, 23-50. doi:10.1146/annurev-psych-122414-033457

Feil, J., Sheppard, D., Fitzgerald, P. B., Yucel, M., Lubman, D. I., & Bradshaw, J. L. (2010). Addiction, compulsive drug seeking, and the role of frontostriatal mechanisms in regulating inhibitory control. *Neurosci Biobehav Rev*, 35(2), 301-313. doi:10.1016/j.neubiorev.2010.03.001

Fitzpatrick, C. J., Creeden, J. F., Perrine, S. A., & Morrow, J. D. (2016). Lesions of the ventral hippocampus attenuate the acquisition but not expression of sign-tracking behavior in rats. *Hippocampus*, 26(11), 1424-1434. doi:10.1002/hipo.22619

Fitzpatrick, C. J., Geary, T., Creeden, J. F., & Morrow, J. D. (2019). Sign-tracking behavior is difficult to extinguish and resistant to multiple cognitive enhancers. *Neurobiol Learn Mem*, 163, 107045. doi:10.1016/j.nlm.2019.107045

Flagel, S. B., Akil, H., & Robinson, T. E. (2009). Individual differences in the attribution of incentive salience to reward-related cues: Implications for addiction. *Neuropharmacology*, 56 Suppl 1, 139-148. doi:10.1016/j.neuropharm.2008.06.027

Flagel, S. B., Chaudhury, S., Waselus, M., Kelly, R., Sewani, S., Clinton, S. M., . . . Akil, H. (2016). Genetic background and epigenetic modifications in the core of the nucleus accumbens predict addiction-like behavior in a rat model. *Proc Natl Acad Sci U S A*, 113(20), E2861-2870. doi:10.1073/pnas.1520491113

Flagel, S. B., Clark, J. J., Robinson, T. E., Mayo, L., Czuj, A., Willuhn, I., . . . Akil, H. (2011). A selective role for dopamine in stimulus-reward learning. *Nature*, 469(7328), 53-57. doi:10.1038/nature09588

Flagel, S. B., & Robinson, T. E. (2017). Neurobiological Basis of Individual Variation in Stimulus-Reward Learning. *Curr Opin Behav Sci*, 13, 178-185. doi:10.1016/j.cobeha.2016.12.004

Flagel, S. B., Robinson, T. E., Clark, J. J., Clinton, S. M., Watson, S. J., See-man, P., . . . Akil, H. (2010). An animal model of genetic vulnerability to behavioral disinhibition and responsiveness to reward-related cues: implications for addiction. *Neuropsychopharmacology*, 35(2), 388-400. doi:10.1038/npp.2009.142

Flagel, S. B., Waselus, M., Clinton, S. M., Watson, S. J., & Akil, H. (2014). Antecedents and consequences of drug abuse in rats selectively bred for

high and low response to novelty. *Neuropharmacology*, 76 Pt B, 425-436. doi:10.1016/j.neuropharm.2013.04.033

Foulds, J. A., Boden, J. M., Newton-Howes, G. M., Mulder, R. T., & Horwood, L. J. (2017). The role of novelty seeking as a predictor of substance use disorder outcomes in early adulthood. *Addiction*, 112(9), 1629-1637. doi:10.1111/add.13838

Furlong, T. M., Jayaweera, H. K., Balleine, B. W., & Corbit, L. H. (2014). Binge-like consumption of a palatable food accelerates habitual control of behavior and is dependent on activation of the dorsolateral striatum. *J Neurosci*, 34(14), 5012-5022. doi:10.1523/JNEUROSCI.3707-13.2014

Garcia, D., Lester, N., Cloninger, K. M., & Robert Cloninger, C. (2017). Temperament and Character Inventory (TCI). In *Encyclopedia of Personality and Individual Differences* (pp. 1-3). Springer.

Garofalo, S., & di Pellegrino, G. (2015). Individual differences in the influence of task-irrelevant Pavlovian cues on human behavior. *Front Behav Neurosci*, 9, 163. doi:10.3389/fnbeh.2015.00163

Giuliano, C., Belin, D., & Everitt, B. J. (2019). Compulsive Alcohol Seeking Results from a Failure to Disengage Dorsolateral Striatal Control over Behavior. *J Neurosci*, 39(9), 1744-1754. doi:10.1523/JNEUROSCI.2615-18.2018

Giuliano, C., Pena-Oliver, Y., Goodlett, C. R., Cardinal, R. N., Robbins, T. W., Bullmore, E. T., . . . Everitt, B. J. (2018). Evidence for a Long-Lasting Compulsive Alcohol Seeking Phenotype in Rats. *Neuropsychopharmacology*, 43(4), 765-777. doi:10.1038/npp.2017.105

Haber, S. N., Fudge, J. L., & McFarland, N. R. (2000). Striatonigrostriatal pathways in primates form an ascending spiral from the shell to the dorsolateral striatum. *J Neurosci*, 20(6), 2369-2382. doi:10.1523/JNEUROSCI.20-06-02369.2000

Hardwick, R. M., Forrence, A. D., Krakauer, J. W., & Haith, A. M. (2019). Time-dependent competition between goal-directed and habitual response preparation. *Nat Hum Behav*, 3(12), 1252-1262. doi:10.1038/s41562-019-0725-0

Hildebrandt, M. K., Dieterich, R., & Endrass, T. (2021). Disentangling substance use and related problems: urgency predicts substance-related problems beyond the degree of use. *BMC Psychiatry*, 21(1), 242. doi:10.1186/s12888-021-03240-z

Hogarth, L., & Hardy, L. (2018). Depressive statements prime goal-directed alcohol-seeking in individuals who report drinking to cope with negative affect. *Psychopharmacology (Berl)*, 235(1), 269-279. doi:10.1007/s00213-017-4765-8

Hyman, S. E., & Malenka, R. C. (2001). Addiction and the brain: the neurobiology of compulsion and its persistence. *Nat Rev Neurosci*, 2(10), 695-703.

doi:10.1038/35094560

Hynes, T. J., Thomas, C. S., Zumbusch, A. S., Samson, A., Petriman, I., Mrdja, U., . . . Lovic, V. (2018). Early life adversity potentiates expression of addiction-related traits. *Prog Neuropsychopharmacol Biol Psychiatry*, 87(Pt A), 56-67. doi:10.1016/j.pnpbp.2017.09.005

Ikemoto, S. (2007). Dopamine reward circuitry: two projection systems from the ventral midbrain to the nucleus accumbens-olfactory tubercle complex. *Brain Res Rev*, 56(1), 27-78. doi:10.1016/j.brainresrev.2007.05.004

Kalivas, P. W., & McFarland, K. (2003). Brain circuitry and the reinstatement of cocaine-seeking behavior. *Psychopharmacology (Berl)*, 168(1-2), 44-56. doi:10.1007/s00213-003-1393-2

Kasal, M. I., Besiroglu, L., Zorlu, N., Dikmeer, N., Bilge, A., Durmaz, E., . . . Sebold, M. (2021). Fronto-striatal structures related with model-based control as an endophenotype for obsessive-compulsive disorder. *Sci Rep*, 11(1), 11951. doi:10.1038/s41598-021-91447-5

Kim, S., Kwok, S., Mayes, L. C., Potenza, M. N., Rutherford, H. J. V., & Strathairn, L. (2017). Early adverse experience and substance addiction: dopamine, oxytocin, and glucocorticoid pathways. *Ann N Y Acad Sci*, 1394(1), 74-91. doi:10.1111/nyas.13140

Koshy Cherian, A., Kucinski, A., Pitchers, K., Yegla, B., Parikh, V., Kim, Y., . . . Sarter, M. (2017). Unresponsive Choline Transporter as a Trait Neuromarker and a Causal Mediator of Bottom-Up Attentional Biases. *J Neurosci*, 37(11), 2947-2959. doi:10.1523/JNEUROSCI.3499-16.2017

Krueger, R. F., Markon, K. E., Patrick, C. J., Benning, S. D., & Kramer, M. D. (2007). Linking antisocial behavior, substance use, and personality: an integrative quantitative model of the adult externalizing spectrum. *J Abnorm Psychol*, 116(4), 645-666. doi:10.1037/0021-843X.116.4.645

Kutlu, M. G., Zachry, J. E., Melugin, P. R., Tat, J., Cajigas, S., Isiktas, A. U., . . . Calipari, E. S. (2022). Dopamine signaling in the nucleus accumbens core mediates latent inhibition. *Nature Neuroscience*, 25(8), 1071-1081. doi:10.1038/s41593-022-01126-1

Llorente, R., O' Shea, E., Gutierrez-Lopez, M. D., Llorente-Berzal, A., Colado, M. I., & Viveros, M. P. (2010). Sex-dependent maternal deprivation effects on brain monoamine content in adolescent rats. *Neurosci Lett*, 479(2), 113-117. doi:10.1016/j.neulet.2010.05.039

Lomanowska, A. M., Lovic, V., Rankine, M. J., Mooney, S. J., Robinson, T. E., & Kraemer, G. W. (2011). Inadequate early social experience increases the incentive salience of reward-related cues in adulthood. *Behav Brain Res*, 220(1), 91-99. doi:10.1016/j.bbr.2011.01.033

- Luscher, C., Robbins, T. W., & Everitt, B. J. (2020). The transition to compulsion in addiction. *Nat Rev Neurosci*, 21(5), 247-263. doi:10.1038/s41583-020-0289-z
- Luscher, C., & Ungless, M. A. (2006). The mechanistic classification of addictive drugs. *PLoS Med*, 3(11), e437. doi:10.1371/journal.pmed.0030437
- Martin, S. J., & Clark, R. E. (2007). The rodent hippocampus and spatial memory: from synapses to systems. *Cell Mol Life Sci*, 64(4), 401-431. doi:10.1007/s00018-007-6336-3
- Moeller, F. G., Barratt, E. S., Dougherty, D. M., Schmitz, J. M., & Swann, A. C. (2001). Psychiatric aspects of impulsivity. *Am J Psychiatry*, 158(11), 1783-1793. doi:10.1176/appi.ajp.158.11.1783
- Moggi, F. (2018). Epidemiology, etiology and treatment of patients with psychosis and co-morbid substance use disorder. *Ther Umsch*, 75(1), 37-43. doi:10.1024/0040-5930/a000964
- Montague, P. R., Dayan, P., & Sejnowski, T. J. (1996). A framework for mesencephalic dopamine systems based on predictive Hebbian learning. *J Neurosci*, 16(5), 1936-1947. doi:10.1523/JNEUROSCI.16-05-01936.1996
- Morrison, S. E., Bamkole, M. A., & Nicola, S. M. (2015). Sign Tracking, but Not Goal Tracking, is Resistant to Outcome Devaluation. *Front Neurosci*, 9, 468. doi:10.3389/fnins.2015.00468
- Nikulina, E. M., Lacagnina, M. J., Fanous, S., Wang, J., & Hammer, R. P., Jr. (2012). Intermittent social defeat stress enhances mesocorticolimbic DeltaFosB/BDNF co-expression and persistently activates corticostriatal neurons: implications for vulnerability to psychostimulants. *Neuroscience*, 218, 15-26. doi:10.1016/j.neuroscience.2012.04.012
- Pascoli, V., Terrier, J., Espallergues, J., Valjent, E., O'Connor, E. C., & Luscher, C. (2014). Contrasting forms of cocaine-evoked plasticity control components of relapse. *Nature*, 509(7501), 459-464. doi:10.1038/nature13257
- Pascoli, V., Turiault, M., & Luscher, C. (2011). Reversal of cocaine-evoked synaptic potentiation resets drug-induced adaptive behaviour. *Nature*, 481(7379), 71-75. doi:10.1038/nature10709
- Pavlov, P. I. (2010). Conditioned reflexes: An investigation of the physiological activity of the cerebral cortex. *Ann Neurosci*, 17(3), 136-141. doi:10.5214/ans.0972-7531.1017309
- Perkins, K. A., Lerman, C., Coddington, S. B., Jetton, C., Karelitz, J. L., Scott, J. A., & Wilson, A. S. (2008). Initial nicotine sensitivity in humans as a function of impulsivity. *Psychopharmacology (Berl)*, 200(4), 529-544. doi:10.1007/s00213-008-1231-7
- Pitchers, K. K., Wood, T. R., Skrzynski, C. J., Robinson, T. E., & Sarter, M. (2017). The ability for cocaine and cocaine-associated cues to compete for

- attention. *Behav Brain Res*, 320, 302-315. doi:10.1016/j.bbr.2016.11.024
- Potvin, S., Pelletier, J., Grot, S., Hebert, C., Barr, A. M., & Lecomte, T. (2018). Cognitive deficits in individuals with methamphetamine use disorder: A meta-analysis. *Addict Behav*, 80, 154-160. doi:10.1016/j.addbeh.2018.01.021
- Pruessner, J. C., Champagne, F., Meaney, M. J., & Dagher, A. (2004). Dopamine release in response to a psychological stress in humans and its relationship to early life maternal care: a positron emission tomography study using [¹¹C]raclopride. *J Neurosci*, 24(11), 2825-2831. doi:10.1523/JNEUROSCI.3422-03.2004
- Radwanska, K., & Kaczmarek, L. (2012). Characterization of an alcohol addiction-prone phenotype in mice. *Addict Biol*, 17(3), 601-612. doi:10.1111/j.1369-1600.2011.00394.x
- Robinson, T. E., & Berridge, K. C. (2000). The psychology and neurobiology of addiction: an incentive-sensitization view. *Addiction*, 95 Suppl 2, S91-117. doi:10.1080/09652140050111681
- Robinson, T. E., & Flagel, S. B. (2009a). Dissociating the predictive and incentive motivational properties of reward-related cues through the study of individual differences. *Biol Psychiatry*, 65(10), 869-873. doi:10.1016/j.biopsych.2008.09.006
- Robinson, T. E., & Flagel, S. B. (2009b). Dissociating the predictive and incentive motivational properties of reward-related cues through the study of individual differences. *Neuropsychopharmacology*, 34(10), 2115-2127. doi:10.1038/npp.2009.42
- Rohan, M. L., Lowen, S. B., Rock, A., & Andersen, S. L. (2021). Novelty preferences and cocaine-associated cues influence regions associated with the salience network in juvenile female rats. *Pharmacol Biochem Behav*, 203, 173117. doi:10.1016/j.pbb.2021.173117
- Rutherford, H. J., Williams, S. K., Moy, S., Mayes, L. C., & Johns, J. M. (2011). Disruption of maternal parenting circuitry by addictive process: rewiring of reward and stress systems. *Front Psychiatry*, 2, 37. doi:10.3389/fpsy.2011.00037
- Sarter, M., & Phillips, K. B. (2018). The neuroscience of cognitive-motivational styles: Sign- and goal-trackers as animal models. *Behav Neurosci*, 132(1), 1-12. doi:10.1037/bne0000226
- Saunders, B. T., & Robinson, T. E. (2011). Individual variation in the motivational properties of cocaine. *Neuropsychopharmacology*, 36(8), 1668-1676. doi:10.1038/npp.2011.48
- Saunders, B. T., Yager, L. M., & Robinson, T. E. (2013). Cue-evoked cocaine "craving" : role of dopamine in the accumbens core. *J Neurosci*, 33(35), 13989-14000. doi:10.1523/JNEUROSCI.0450-13.2013

- Schad, D. J., Rapp, M. A., Garbusow, M., Nebe, S., Sebold, M., Obst, E., . . . Huys, Q. J. M. (2020). Dissociating neural learning signals in human sign- and goal-trackers. *Nat Hum Behav*, 4(2), 201-214. doi:10.1038/s41562-019-0765-5
- Schettino, M., Ceccarelli, I., Tarvainen, M., Martelli, M., Orsini, C., & Ottaviani, C. (2022). From skinner box to daily life: Sign-tracker phenotype cosegregates with impulsivity, compulsivity, and addiction tendencies in humans. *Cogn Affect Behav Neurosci*, 22(3), 567-581. doi:10.3758/s13415-022-01014-y
- Selden, N. R., Everitt, B. J., Jarrard, L. E., & Robbins, T. W. (1991). Complementary roles for the amygdala and hippocampus in aversive conditioning to explicit and contextual cues. *Neuroscience*, 42(2), 335-350. doi:10.1016/0306-4522(91)90379-3
- Singer, B. F., Guptaroy, B., Austin, C. J., Wohl, I., Lovic, V., Seiler, J. L., . . . Aragona, B. J. (2016). Individual variation in incentive salience attribution and accumbens dopamine transporter expression and function. *Eur J Neurosci*, 43(5), 662-670. doi:10.1111/ejn.13134
- Smith, R. J., & Laiks, L. S. (2018). Behavioral and neural mechanisms underlying habitual and compulsive drug seeking. *Prog Neuropsychopharmacol Biol Psychiatry*, 87(Pt A), 11-21. doi:10.1016/j.pnpbp.2017.09.003
- Solomon, R. L., & Corbit, J. D. (1974). An opponent-process theory of motivation. I. Temporal dynamics of affect. *Psychol Rev*, 81(2), 119-145. doi:10.1037/h0036128
- Spanagel, R. (2017). Animal models of addiction. *Dialogues Clin Neurosci*, 19(3), 247-258. Retrieved from <https://www.ncbi.nlm.nih.gov/pubmed/29302222>
- Stautz, K., & Cooper, A. (2013). Impulsivity-related personality traits and adolescent alcohol use: a meta-analytic review. *Clin Psychol Rev*, 33(4), 574-592. doi:10.1016/j.cpr.2013.03.003
- Szumliński, K. K., Abernathy, K. E., Oleson, E. B., Klugmann, M., Lominac, K. D., He, D. Y., . . . Kalivas, P. W. (2006). Homer isoforms differentially regulate cocaine-induced neuroplasticity. *Neuropsychopharmacology*, 31(4), 768-777. doi:10.1038/sj.npp.1300890
- Tomie, A. (2018). Sign-Tracking and Drug Addiction. In J. D. Jentsch (Ed.), *The Neuroscience of Drug Addiction* (pp. 123-145). Springer.
- Tomie, A., Grimes, K. L., & Pohorecky, L. A. (2008). Behavioral characteristics and neurobiological substrates shared by Pavlovian sign-tracking and drug abuse. *Brain Res Rev*, 58(1), 121-135. doi:10.1016/j.brainresrev.2007.12.003
- Tomie, A., Tirado, A. D., Yu, L., & Pohorecky, L. A. (2004). Pavlovian autoshaping procedures increase plasma corticosterone and levels of norepinephrine and serotonin in prefrontal cortex in rats. *Behavioural Brain Research*, 153(1), 99-106. doi:10.1016/j.bbr.2003.11.006

Veber, M., & Weidemann, A. J. S. Z. F. S. A. (2018). United Nations Office on Drugs and Crime: World Drug Report. Retrieved from <https://www.unodc.org/wdr2017/index.html>

Volkow, N. D., & Morales, M. (2015). The Brain on Drugs: From Reward to Addiction. *Cell*, 162(4), 712-725. doi:10.1016/j.cell.2015.07.046

Volkow, N. D., Wise, R. A., & Baler, R. (2017). The dopamine motive system: implications for drug and food addiction. *Nat Rev Neurosci*, 18(12), 741-752. doi:10.1038/nrn.2017.130

Wang, B., Shaham, Y., Zitzman, D., Azari, S., Wise, R. A., & You, Z. B. (2005). Cocaine experience establishes control of midbrain glutamate and dopamine by corticotropin-releasing factor: a role in stress-induced relapse to drug seeking. *J Neurosci*, 25(22), 5389-5396. doi:10.1523/JNEUROSCI.0955-05.2005

Yamamoto, D. J., Nelson, A. M., Mandt, B. H., Larson, G. A., Rorabaugh, J. M., Ng, C. M., . . . Zahniser, N. R. (2013). Rats classified as low or high cocaine locomotor responders: a unique model involving striatal dopamine transporters that predicts cocaine addiction-like behaviors. *Neurosci Biobehav Rev*, 37(8), 1689-1700. doi:10.1016/j.neubiorev.2013.07.002

Zapata, A., Minney, V. L., & Shippenberg, T. S. (2010). Shift from goal-directed to habitual cocaine seeking after prolonged experience in rats. *J Neurosci*, 30(46), 15457-15463. doi:10.1523/JNEUROSCI.4072-10.2010

Zhou, X., Zimmermann, K., Xin, F., Zhao, W., Derckx, R. T., Sassmannshausen, A., . . . Becker, B. (2019). Cue Reactivity in the Ventral Striatum Characterizes Heavy Cannabis Use, Whereas Reactivity in the Dorsal Striatum Mediates Dependent Use. *Biol Psychiatry Cogn Neurosci Neuroimaging*, 4(8), 751-762. doi:10.1016/j.bpsc.2019.04.006

Zimmerberg, B., & Shartrand, A. M. (1992). Temperature-dependent effects of maternal separation on growth, activity, and amphetamine sensitivity in the rat. *Dev Psychobiol*, 25(3), 213-226. doi:10.1002/dev.420250306

Zou, Z., Wang, H., d' Oleire Uquillas, F., Wang, X., Ding, J., & Chen, H. (2017). Definition of Substance and Non-substance Addiction. *Adv Exp Med Biol*, 1010, 21-41. doi:10.1007/978-981-10-5562-1_2

Zuckerman, M., Kuhlman, D. M., Joireman, J., Teta, P., & Kraft, M. (1993). A comparison of three structural models for personality: The Big Three, the Big Five, and the Alternative Five. *Journal of Personality and Social Psychology*, 65(4), 757-768. doi:10.1037/0022-3514.65.4.757

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv – Machine translation. Verify with original.