

## A New Approach of Data-driven Simulation and its Application to Solar Active Region 12673 Postprint

**Authors:** Zhi-Peng Liu, Chao-Wei Jiang, Xin-Kai Bian, Qing-Jun Liu, Peng Zou and Xue-Shang Feng

**Date:** 2025-01-07T00:00:00+00:00

### Abstract

The solar coronal magnetic field is a pivotal element in the study of eruptive phenomena, and understanding its dynamic evolution has long been a focal point in solar physics. Numerical models, driven directly by observation data, serve as indispensable tools in investigating the dynamics of the coronal magnetic field. This paper presents a new approach to electric field inversion, which involves modifying the electric field derived from the DAVE4VM velocity field using ideal Ohm's law. The time series of the modified electric field is used as a boundary condition to drive a magnetohydrodynamics model, which is applied to simulate the magnetic field evolution of active region 12673. The simulation results demonstrate that our method enhances the magnetic energy injection through the bottom boundary, as compared with energy injection calculated directly from the DAVE4VM code, and reproduces the evolution of the photospheric magnetic flux. The coronal magnetic field structure is also in morphological similarity to the coronal loops. This new approach will be applied to the high-accuracy simulation of eruption phenomena and provide more details on the dynamical evolution of the coronal magnetic field.

### Full Text

#### Preamble

Research in Astronomy and Astrophysics, 24:125005 (10pp), 2024 December © 2024 National Astronomical Observatories, CAS and IOP Publishing Ltd. All rights, including for text and data mining, AI training, and similar technologies, are reserved. Printed in China. <https://doi.org/10.1088/1674-4527/ad862b>

A New Approach of Data-driven Simulation and its Application to Solar Active Region 12673

Zhi-Peng Liu, Chao-Wei Jiang, Xin-Kai Bian, Qing-Jun Liu, Peng Zou, and Xue-Shang Feng

Institute of Space Science and Applied Technology, Harbin Institute of Technology, Shenzhen 518055, China; chaowei@hit.edu.cn

Received 2024 July 26; revised 2024 September 30; accepted 2024 October 8; published 2024 December 6

## Abstract

The solar coronal magnetic field is a pivotal element in the study of eruptive phenomena, and understanding its dynamic evolution has long been a focal point in solar physics. Numerical models driven directly by observation data serve as indispensable tools for investigating the dynamics of the coronal magnetic field. This paper presents a new approach to electric field inversion, which involves modifying the electric field derived from the DAVE4VM velocity field using ideal Ohm's law. The time series of the modified electric field is used as a boundary condition to drive a magnetohydrodynamics model, which is applied to simulate the magnetic field evolution of active region 12673. The simulation results demonstrate that our method enhances the magnetic energy injection through the bottom boundary, as compared with energy injection calculated directly from the DAVE4VM code, and reproduces the evolution of the photospheric magnetic flux. The coronal magnetic field structure also shows morphological similarity to the coronal loops. This new approach will be applied to high-accuracy simulation of eruption phenomena and provide more details on the dynamical evolution of the coronal magnetic field.

Key words: Sun: coronal mass ejections (CMEs) – Sun: flares – magnetohydrodynamics (MHD) – methods: numerical

## 1. Introduction

Solar eruptive activity significantly impacts space weather, with solar flares and coronal mass ejections being the primary events originating from solar active regions (ARs, Webb & Howard 2012). These phenomena release accumulated free magnetic energy in the coronal magnetic field, resulting in radiation emission and acceleration and heating of the surrounding plasma (Forbes 2000). The energetic plasmas carrying the Sun's magnetic field may interfere with the Earth's magnetic field and spacecraft, potentially causing disruptive effects on human activities. Therefore, understanding the dynamic processes of these eruptive events is crucial for mitigating their impact.

It is widely acknowledged that variations of the coronal magnetic field breed these eruptions. Nonetheless, direct observations of the coronal magnetic field are extremely challenging (Lin et al. 2004). While it is feasible to infer the trajectories of coronal magnetic field lines from some coronal loops using extreme-ultraviolet (EUV) and X-ray imaging (Brosius & White 2006; Tomczyk et

al. 2008; Liu 2009), obtaining detailed information remains elusive. The primary source of magnetic field data available to us predominantly originates from the photosphere. Consequently, numerical simulation emerges as an important tool for studying the coronal magnetic field. One prominent numerical approach, the Nonlinear Force-Free Field (NLFFF) model (Schrijver et al. 2006; Metcalf et al. 2008), derives the 3D coronal magnetic field through extrapolation from photospheric vector magnetograms. This modeling technique, refined over several decades, has significantly enhanced our understanding of the 3D coronal magnetic field configuration (Wiegelmann 2008; Regnier 2013; Guo et al. 2017; Wiegelmann & Sakurai 2021).

The magnetic field in the NLFFF model is in static equilibrium (in which the magnetic pressure-gradient force is balanced by the magnetic tension force), thus impeding detailed investigations of the triggering and driving processes of eruption events, during which the field evolves dynamically and the forces are significantly unbalanced. Consequently, a data-driven magnetohydrodynamics (MHD) model, utilizing vector magnetograms as boundary conditions, has been proposed to simulate the dynamic evolution of the coronal magnetic field. This time-dependent, data-driven model adeptly simulates the gradual, quasi-static evolution of the coronal magnetic field (Wu et al. 2006, 2009; Chitta et al. 2014), the rapid progression of solar eruptions (Jiang & Feng 2013; Kliem et al. 2013; Guo et al. 2021), and the transition from quasi-static to fast eruption phases (Jiang et al. 2016a, 2016b; Guo et al. 2019; Zhong et al. 2021).

According to the different levels of simplification made to the MHD equations, these models can be classified into three groups: the magneto-frictional model (Cheung et al. 2015; Price et al. 2019; Yardley et al. 2021), the zero- $\beta$  MHD model (Inoue et al. 2018; Guo et al. 2021; Kaneko et al. 2021), and the full MHD model (Fan et al. 2011; Feng et al. 2012; Guo et al. 2024). Further detailed information about these models can be found in Jiang et al. (2022).

On the other hand, according to the ways of implementing the boundary condition of the data-driven model, it can be subdivided into the magnetic field (B) driven (Wu et al. 2006; Feng et al. 2015, 2017; Jiang et al. 2016a, 2016b), the velocity (V) driven (Guo et al. 2019; Hayashi et al. 2019; Jiang et al. 2021; Zhong et al. 2021; Wang et al. 2022, 2023) and the electric field (E) driven (Cheung & DeRosa 2012; Hayashi et al. 2018; Pomoell et al. 2019; Price et al. 2019). The B-driven approach uses the observed magnetic field directly at the bottom boundary of the model, which is very straightforward, but this introduces magnetic divergence errors from the boundary (Tóth 2000). The V-driven approach uses the photospheric flow that can be recovered from a time sequence of observed magnetograms, and this velocity-recovering technique is relatively mature, such as the local correlation tracking technique (November & Simon 1988; Chae 2001; Fisher & Welsch 2008) and the differential affine velocity estimator (DAVE) method (Schuck 2006, 2008; Schuck & Antiochos 2019). The V-driven model gives a good representation of the surface motions of magnetic flux, such as shearing and rotation of sunspots, but cannot be sufficiently self-

consistent for the calculation of emerging flux, as the vertical component of the photospheric velocity is difficult to recover by currently available methods such as DAVE4VM. Jiang et al. (2021) found that there is a build-up of magnetic fields at locations of small velocities around the AR and that the simulated magnetograms differ more and more from the observations as time goes on. The E-driven approach, which uses the time series of the electric field as the bottom boundary, has the advantage that the magnetic field satisfies its solenoidal condition, and the corresponding disadvantage is that the inversion of the electric field is more difficult and complex (Fisher et al. 2010, 2012, 2015, 2020; Kazachenko et al. 2014; Lumme et al. 2017).

Although direct observation of the electric field in the solar photosphere is feasible using the Stark effect (Wien 1916), the limited sensitivity of measuring instruments poses significant challenges (Moran & Foukal 1991). A common approach involves indirectly deriving the electric field from the photospheric magnetic field. Typical practice involves dividing the electric field into an inductive part and a non-inductive part. While the inductive component can be derived by solving Faraday's law, the non-inductive component is specified using different assumptions (Fisher et al. 2010; Cheung & DeRosa 2012; Cheung et al. 2015; Lumme et al. 2017): neglecting the non-inductive part ( $\nabla \cdot \mathbf{E} = 0$ , where  $\phi$  is the non-inductive potential), the emergence of a twisted field, and uniform vortical motion ( $\nabla^2 \phi = -\Omega B_z$ ). The free parameters  $U$  and  $\Omega$  in the latter two assumptions require additional velocity fields for estimation, such as Dopplergrams or DAVE4VM velocity. Kazachenko et al. (2014) describe the PTD-Doppler-FLCT-Ideal (PDFI) method, which, as the name suggests, includes the PTD method (which stands for poloidal-toroidal decomposition first introduced by Chandrasekhar 1961), the Dopplergram velocity, the optical flow of the FLCT method and the ideal constraint ( $\mathbf{E} \cdot \mathbf{B} = 0$ ). The PTD method serves as the inductive component, while the remaining elements are considered non-inductive components. A detailed implementation of the PDFI method can be found in Fisher et al. (2020).

Another relatively straightforward approach involves using FLCT or DAVE4VM code to process vector magnetograms to obtain the velocity field, followed by deriving the electric field using ideal Ohm's law ( $\mathbf{E} = -\mathbf{V} \times \mathbf{B}$ ). However, due to the large noise levels in weak magnetic field regions, velocity inversion is highly susceptible to outliers, potentially leading to the manifestation of unphysical phenomena (Schuck 2008). Thus, correction procedures are necessary after the inversion of the electric field.

In this paper, we present a new approach to correct the DAVE4VM-based electric field and perform a new E-driven MHD simulation for the one-day evolution of AR 12673 using the corrected electric field. The details of the electric field inversion are given in Section 2. The model setup and parameters are described in Section 3. The results of the simulation are shown in Section 4, and we conclude in Section 5.

## 2. Electric Field Inversion

We use the Space-weather HMI Active Region Patches (Bobra et al. 2014) data product which provides the AR's vector magnetograms with a time cadence of 12 minutes and a pixel (grid) size of 0.5. Before using the DAVE4VM code, we first rebin the data to a grid size of  $\Delta x = \Delta y = 1$  to reduce the computing time, and any data gap is filled with a simple linear interpolation in time. The DAVE4VM needs inputs of spatial (in the horizontal direction) and temporal derivatives of magnetic field and window size. All the derivatives of the magnetic field (as mentioned in this section) are calculated by central difference, for example, where  $nC$  is the time index. The window size is chosen as 11.

From the DAVE4VM code and the vector magnetogram  $B_O$ , we obtain the vector velocity  $v_D$  and thus the electric field by assuming that  $E_D = -v_D \times B_O$ . The electric field can be decomposed into two components, i.e.,  $E_D = (E_h, E_z)$ , where  $h$  denotes the horizontal component and  $z$  the vertical component. However, using  $E_D$  in the magnetic induction equation ( $B_O/t = -\nabla \times E_D$ ) does not recover the evolution of  $B_O$ . Since the horizontal velocity ( $v_h$ ) is often more accurate than the vertical velocity ( $v_z$ ), the vertical electric field  $E_z$  should be more accurate than the horizontal electric field  $E_h$ , and therefore we have more freedom to modify  $E_h$ . Our purpose is to modify  $E_D$  to a new electric field  $E$  such that using the magnetic induction equation  $B_z$  can be fully recovered.

Before this, we smoothed the data  $B_O$  and  $E_D$  since in our data-driven model, the implementation of bottom boundary conditions is based on numerical differences. These data were spatially smoothed using a Gaussian smoothing with FWHM of 8, and temporally smoothed using a boxcar average of 120 minutes, and the smoothed version of the two fields is also denoted by  $B_O$  and  $E_D$ .

The horizontal electric field  $E_h$  can be decomposed into two parts:  $E_h = E_h^I + E_h^N$ , where  $E_h^I$  is the inductive part,  $E_h^N$  is the non-inductive part, and  $E_h^I = \nabla \times (f\hat{z})$ . The two scalars  $f(x, y)$  and  $(x, y)$  are functions of  $x$  and  $y$ .

Using the vertical component of the induction equation, we have  $B_z/t = -\nabla_h \times E_h = -\nabla_h \times (E_h^I + E_h^N) = -\nabla_h \times E_h^I = -\nabla_h \times (\nabla_h \times f\hat{z}) = \nabla_h^2 f$ , and solving this 2D Poisson equation in a rectangular region  $A = [0, L_x] \times [0, L_y]$  obtains  $f$  and thus  $E_h^I$ . In the paper, we used the procedure `IMSL{POISSON2D}` in IDL to solve the Poisson equation. Two choices may be used for the boundary conditions. If we use the Neumann boundary conditions such that  $f/n = 0$  on the boundary  $C$ , we have that at the  $x = 0$  and  $x = L_x$  boundaries and at the  $y = 0$  and  $y = L_y$  boundaries, which means that  $E_h^I$  is perpendicular to the boundary lines. This requires that  $\int_C E_h \cdot dl = 0$ , which is, however, not generally satisfied by the magnetogram.

If we use the Dirichlet boundary conditions  $f = 0$  on the boundary  $C$ , at the  $x = 0$  and  $x = L_x$  boundaries we have  $E_{\hat{I}} = -f/y = 0$  and at the  $y = 0$  and  $y = L_y$  boundaries  $E_{\hat{I}} = f/x = 0$ , which means that  $E_{\hat{I}}$  is parallel to the boundary lines. Then we can specify the non-inductive part by assuming  $E_{\hat{N}} = \hat{h}$ , which requires that  $\hat{h}^2 = \hat{h} \cdot E_{\hat{h}}$ , and solving this 2D Poisson equation obtains  $\hat{h}$  and thus  $E_{\hat{N}}$ . Also, two types of boundary conditions can be used. If we use the Neumann boundary conditions, then  $E_{\hat{N}}$  will be parallel to boundary lines, and if we use the Dirichlet boundary conditions,  $E_{\hat{N}}$  will be perpendicular to boundary lines. A good choice is that the total field  $E_{\hat{h}}$  is parallel to the boundary lines, and thus the Dirichlet boundary conditions should be used for  $E_{\hat{I}}$  and the Neumann boundary conditions for  $E_{\hat{N}}$ . By combining the two parts, we have the final electric field that can be used in the numerical code as  $E = (E_{\hat{I}} + E_{\hat{N}}, E_z)$ .

### 3. Model

We carried out a data-driven simulation using the DARE-MHD code (Jiang et al. 2016a). This model has been used in numerous data-driven simulations for solar coronal evolution and eruptions (e.g., Jiang et al. 2021, 2022; Wang et al. 2022, 2023). In this study, the simulation volume is a cubic box in the Cartesian coordinate system, with a size of 512 (corresponding to around 368 Mm) and the origin point  $(x, y, z) = (0, 0, 0)$  at the center of the bottom boundary. The computational volume is resolved by an adaptive mesh refinement grid in the simulation, in which the highest resolution is 1 and the lowest resolution is 8. To save computing time, we multiplied the observed magnetic field by a factor of 0.025 (therefore the maximal magnetic field  $B_z$  in the model is around 55 G initially), and increased the cadence of the input maps by 68.6 times. The other parameters utilized in the model are the same as those in Jiang et al. (2021) except that the kinetic viscosity coefficient is set to a small value of  $\nu = 0.1\Delta x^2/\Delta t$ . In addition, all the variables in the numerical model are normalized using typical coronal values:  $\rho_{\text{cor}} = 2.29 \times 10^{-15} \text{ g cm}^{-3}$  (density),  $T_{\text{cor}} = 1 \times 10^6 \text{ K}$  (temperature),  $H_{\text{cor}} = 11.52 \text{ Mm}$  (typical length in coronal scale),  $B_{\text{cor}} = 1.86 \text{ G}$  (magnetic field), and  $v_{\text{cor}} = 110 \text{ km s}^{-1}$  (velocity). In the rest of the paper, all variables and quantities are mentioned as normalized values, if they are not specified.

#### 3.1. Initial Conditions

AR 12673 is a very flare-productive region and has been studied by many authors (e.g., Yang et al. 2017; Chertok et al. 2018; Hou et al. 2018; Inoue et al. 2018; Jiang et al. 2018; Liu et al. 2018; Verma 2018; Morosan et al. 2019; Romano et al. 2019; Zou et al. 2019, 2020; Guo et al. 2024), delving into the intricate magnetic field structure of the AR and the mechanism of its eruptions. It appeared at the east solar limb on 2017 August 31, and disappeared on September 30, as observed by the Solar Dynamics Observatory (SDO). Figure 1(a) shows the location of AR 12673 on the solar disk on September 6 00:00

UT. From September 4 to 10, there were four X-class flares and twenty-seven M-class flares observed, and two X-class flares occurred on September 6. The first X-class flare (X2.2) started at 08:57 UT, reached its peak at 09:10 UT, and ended at 09:17 UT. Subsequently, the second X-class flare (X9.3) began at 11:53 UT, peaked at 12:02 UT, and ended at 12:10 UT. Remarkably, the second flare marked the largest flare in solar cycle 24.

We used SDO/HMI vector magnetograms for AR 12673 in a one-day period from 2017 September 6 00:00 UT to 7 00:00 UT. The magnetograms have a resolution of  $0.5''$  and a cadence of 720 s. The resolution used in our simulation is  $1''$ , which is rebinned from the original data in order to reduce the computational time. There is a data gap of magnetograms (from 06:00 UT to 08:36 UT) and we used a simple linear interpolation in time to fill the data gap. We chose the magnetogram of 00:00 UT on 2017 September 6 as the initial map and constructed the 3D magnetic field. First, we smooth the vector magnetogram using Gaussian smoothing with FWHM of  $8''$ , as shown in Figure 1(b), in order to filter out the small-scale structures and reduce the Lorentz force to make it easier to reach the equilibrium state. Then the initial 3D magnetic field is obtained by magnetic field extrapolation using the CESE-MHD-NLFFF code (Jiang & Feng 2013). Finally, we input the extrapolated field to the DARE-MHD model, along with an isothermal plasma atmosphere of temperature  $T = 1$  MK stratified by solar gravity with density  $\rho = 1$  at the bottom boundary (same as Jiang et al. 2021), and relaxed them to an MHD equilibrium, which is used as the initial condition of subsequent data-driven simulation.

### 3.2. Boundary Conditions

Our model does not use a ghost cell outside of the actual computational volume. The simulation volume extends from  $z = 0$ , and the  $z = 0$  plane is exactly the bottom surface where the data-driven boundary conditions are applied. In our code, all the MHD variables are assigned at the grid nodes (i.e., corner of the grid cells) rather than the cell center (as shown in Figure 2).

At the bottom surface, we update the magnetic field using the induction equation:  $\partial \mathbf{B} / \partial t = -\nabla \times \mathbf{E}$ . Equation (8) is discretized using a forward difference in time, a central difference in the horizontal direction on the surface, and a one-sided 2nd order difference in the  $z$ -direction. In practice we found such a scheme could be unstable when  $B_z$  is small, i.e., near the polarity inversion line (PIL). Therefore, a surface diffusion term is added to the induction equation for numerical stability. Specifically, taking the  $B_x$  component as an example, the scheme is:

$$B_{x,i,j,0}^{n+1} = B_{x,i,j,0}^n + \Delta t [(\nabla \times \mathbf{E})_x + \frac{1}{2} \frac{h^2}{\Delta z} B_x]_{i,j,0}^n$$

where the subscripts  $i, j, k$  denote the grid points in  $x, y, z$  directions, respectively, and  $k = 0$  for the points at the bottom boundary.  $E_{i,j,0}$  is specified by the inverted electric field, while for  $k > 0$ ,  $E_{i,j,k} = -\mathbf{v}_{i,j,k} \times \mathbf{B}_{i,j,k}$ .

With the magnetic field updated, we also need to update the plasma density, velocity, and pressure at the bottom boundary. Here the density and pressure are simply fixed to be their initial values. Note that the electric field contains both the ideal and resistive part ( $\mathbf{E} = -\mathbf{v} \times \mathbf{B} + \mathbf{j}$ ). If we assume that the current density is parallel to the magnetic field,  $\mathbf{j} = \alpha \mathbf{B}$  (which is the force-free assumption), the velocity can be updated by:

$$\mathbf{v} = (\mathbf{E} \times \mathbf{B})/B^2$$

In most cases, the force-free assumption is valid in the corona, thus it also applies to the boundary surface (base of the corona).

In addition, since the time step of the model  $\Delta t$  is determined by the CFL condition, which is around 2 s and is much smaller than the inputting cadence of the electric field (which is  $\Delta t_c/68.6 = 10.5$  s), interpolation of the electric field along time is required to provide the bottom-surface electric field needed by the model at each time step. We utilize a cubic spline interpolation scheme to maintain the continuity of the first-order time derivatives of the electric field (i.e., the changing rate of the electric field is continuous). In this way, the changing rate of the magnetic field at the bottom surface is also continuous.

#### 4. Result

First, we study the evolution of magnetic energy by comparing the volume-integrated value (i.e., magnetic energy stored in the corona) with the cumulative value from the bottom boundary (i.e., magnetic energy injected from the bottom boundary). This is an important way to check the consistency of the simulation. It is expected that, without eruption, the magnetic energy stored in the corona should be close to (but less than, since there is energy dissipation in the corona) the energy injected from the bottom boundary. We calculated the Poynting flux through the bottom boundary as follows,  $\mathbf{S} = (1/4\pi)\mathbf{E} \times \mathbf{B}$ , since  $-\mathbf{j} \cdot \mathbf{S} = -(\mathbf{j} \cdot \mathbf{E}) = (1/4\pi)\mathbf{B} \cdot \mathbf{B}/t$ , and then integrated it over time as the magnetic energy injected into the corona. Besides, for comparison, we also calculated the Poynting flux used directly in the observed field  $\mathbf{B}_O$  and the DAVE4VM-derived electric field  $\mathbf{E}_D$ , which are different from the values in our simulation. Figure 3(a) presents the results.

The blue and red curves represent the DAVE4VM-based and simulated magnetic energy injections, respectively, with the initial magnetic energy added up, while the black line depicts the variation of total magnetic energy in the model data-driven simulation. Overall, the three different values of magnetic energy increase in the simulated time interval, which is in agreement with the increase of the total unsigned magnetic flux (the green line). As expected, the magnetic energy injected from the bottom boundary in the simulation (i.e., the red line) exceeds slightly the volume-integrated magnetic energy (compare the red and black curves). Their deviation increases systematically with time, owing to numerical resistivity in the model.

Initially, the magnetic energy injection in the simulation briefly aligns with the DAVE4VM-based magnetic energy injection. However, with time, the simulated magnetic energy injection begins to surpass that of the DAVE4VM-based injection. This difference continues to increase, reaching around 500 (in a normalized unit of energy) at the end of the simulation, constituting 40% of the DAVE4VM-based magnetic energy injection. Interestingly, the volume-integrated magnetic energy is rather close to the DAVE4VM-based energy injection (e.g., compare the blue and black curves). We note that the DAVE4VM-based energy injection (blue curve) exhibits a brief decrease during the data gap of magnetograms (from  $t = 30$  to  $43$ ), which is not seen in the energy injection curve of the simulation.

Figure 3(b) depicts the evolution of free magnetic energy (the red curve) and kinetic energy (the black curve). In the whole simulation, the kinetic energy maintains very small values of below  $10^{-3}$  of the magnetic energy, indicating that the system evolves in a quasi-static way. Therefore, the simulation did not reproduce the eruptions. Before  $t = 50$ , the free magnetic energy and the total magnetic energy both exhibit a gradual increase. Following this period, the increase of the total magnetic energy experiences two brief decelerations, corresponding to the two decreases in free magnetic energy. Since no eruption is found in the simulation, the free energy decreases do not correspond to the two flares in observation.

We further look into how the simulated magnetic field at the bottom boundary differs from the observed photospheric magnetic field. Figures 4(a)–(c) compare the magnetograms ( $B_x$ ,  $B_y$  and  $B_z$ , separately) in the simulation and the observation at different times. Overall, the distributions of the simulated field resemble the observed one throughout the entire evolution, except for some small-scale discrepancies. Unlike the velocity-driven simulation, the electric field-driven simulation does not exhibit any unreasonable significant magnetic field pile-up at the edge of the main magnetic polarities (see Jiang et al. 2021).

Figure 4(d) presents the scatter plots of the magnetic field at the bottom boundary of the model with the magnetograms, where black, red and green dots represent  $B_x$ ,  $B_y$  and  $B_z$  values, respectively. The numbers indicate the Pearson correlation coefficients of the simulated values and magnetograms. It is evident that the simulated  $B_z$  values align very well with the magnetogram data, while  $B_x$  and  $B_y$  exhibit a slightly lower correspondence. The linear Pearson correlation coefficients are also shown in Figure 4(d), and their evolutions with time are shown in Figure 5. Note that our calculations of the correlation coefficients are based on the entire magnetogram and the full range of the magnetic field, which is different from some other authors who focus on particular regions and values (Price et al. 2019). The results reveal Pearson correlation coefficients above 0.6 for  $B_x$ , exceeding 0.7 for  $B_y$  and consistently approaching 1 for  $B_z$ .

Last, we study the coronal magnetic structure by comparing the EUV observed images with the simulated magnetic field in Figure 6. The first ((a), (e), (i)) and third ((c), (g), (k)) columns are the SDO/AIA observed images in EUV

wavelength of 171 Å and 131 Å, respectively. Panels ((b), (f), (j)) are the magnetic field lines at the corresponding moments. Panels ((d), (h), (l)) are the synthetic images generated from the current density in the corona, following the method as proposed by Cheung & DeRosa (2012) and Jiang et al. (2016b). As can be seen, in the early stage of the simulation, a good morphological similarity is achieved between the simulated emission and the observed EUV images. However, after the X9.3 flare time (near  $t = 60$ ), the magnetic field structure diverges from observations due to the absence of eruption and its significant changes in the simulation. Nevertheless, our simulation shows the formation of a magnetic flux rope along the main PIL before the flare time, and such a C-shaped flux rope is also revealed in previous studies by Jiang et al. (2018) and Guo et al. (2024) using different modeling techniques including NLFFF extrapolation and magneto-frictional model. After the flare time, this flux rope starts to rise gradually and reaches a higher position at the end of the simulation without showing impulsive eruption. As shown by the yellow lines in Figure 6(j), the flux rope still exists after the flare time at  $t = 75$ . There are several reasons why the flux rope did not erupt, for example, the magnetic free energy is not sufficient enough owing to the loss of energy by numerical diffusion in the corona; the boundary conditions have not fully reproduced the transverse field of magnetograms.

## 5. Conclusion and Discussion

In this paper, we first presented a new approach for photospheric electric field inversion by employing the photospheric velocity derived from vector magnetograms with the DAVE4VM code. With the photospheric velocity and magnetic field given, the electric field is simply derived based on Ohm's law, but the magnetic flux distribution cannot be recovered using the magnetic induction equation. This motivated us to modify the horizontal component of the derived electric field by decomposing it into an inductive part and a non-inductive part, where the inductive part is directly solved using the vertical component of the induction equation.

After obtaining the time series of the corrected electric field, we studied the magnetic dynamic evolution of AR 12673 from September 6 to 7, 2017 using a data-driven MHD model driven by the electric field. The result shows that the new approach enhances the magnetic energy injection through the bottom surface by 40% compared to that of the values by the original DAVE4VM-derived electric field. On the other hand, due to the energy dissipation in the corona, the volume-integrated magnetic energy in the corona is rather close to the DAVE4VM-based energy injection. The evolution of both total and free magnetic energy reflects the variation in the bottom magnetic field. Therefore we compared the magnetic field at the bottom boundary in the simulation with the observed magnetograms. As expected, the evolution of the photospheric vertical magnetic field is well recovered with a correlation coefficient of essentially unity. The main structures of the horizontal components are recovered,

and all correlation coefficients are mainly above 0.7. We also compared the simulated coronal magnetic field with EUV observations. At an early stage, the coronal magnetic field structure in our simulation exhibits a notable resemblance to EUV observations. However, at a later stage (after  $t = 60$ ) disparities emerge between the simulated magnetic field structure and the post-eruption observations. This discrepancy arises from the absence of eruption phenomenon in the simulation, which precludes large-scale magnetic field reconnection and reconfiguration.

In future studies, we will consider using more advanced electric field inversion methods, e.g., the PDFI\_{SS} code (Fisher et al. 2020), to provide an electric field that can hopefully recover all three components of the photospheric magnetic field. Moreover, a high-accuracy simulation is required to reduce the magnetic energy loss due to numerical resistivity, such that most of the energy injected from the bottom boundary can be accumulated in the coronal field until an eruption is initiated.

## Acknowledgments

This work was jointly supported by the National Natural Science Foundation of China (NSFC 42174200), Shenzhen Science and Technology Program (grant No. RCJC20210609104422048), Shenzhen Key Laboratory Launching Project (No. ZDSYS20210702140800001), Guangdong Basic and Applied Basic Research Foundation (2023B1515040021), and the Fundamental Research Funds for the Central Universities (grant No. HIT.OCEF.2023047).

## References

- Bobra, M. G., Sun, X., Hoeksema, J. T., et al. 2014, *SoPh*, 289, 3549  
Brosius, J. W., & White, S. M. 2006, *ApJL*, 641, L69  
Chae, J. 2001, *ApJL*, 560, L95  
Chandrasekhar, S. 1961, *Hydrodynamic and Hydromagnetic Stability* (Oxford: Clarendon)  
Chertok, I. M., Belov, A. V., & Abunin, A. A. 2018, *SpWea*, 16, 1549  
Cheung, M. C. M., & DeRosa, M. L. 2012, *ApJ*, 757, 147  
Cheung, M. C. M., Pontieu, B. D., Tarbell, T. D., et al. 2015, *ApJ*, 801, 83  
Chitta, L. P., Kariyappa, R., van Ballegoijen, A. A., DeLuca, E. E., & Solanki, S. K. 2014, *ApJ*, 793, 112  
Fan, Y. L., Wang, H. N., He, H., & Zhu, X. S. 2011, *ApJ*, 737, 39  
Feng, X., Jiang, C., Xiang, C., Zhao, X., & Wu, S. T. 2012, *ApJ*, 758, 62  
Feng, X., Li, C., Xiang, C., et al. 2017, *ApJS*, 233, 10  
Feng, X., Ma, X., & Xiang, C. 2015, *JGRA*, 120, 10159  
Fisher, G., & Welsch, B. 2008, *Subsurface and Atmospheric Influences on Solar Activity ASP Conference Series*, 383, ed. R. Howe et al. (San Francisco, CA: ASPC), 373  
Fisher, G. H., Abbett, W. P., Bercik, D. J., et al. 2015, *SpWea*, 13, 369  
Fisher, G. H., Kazachenko, M. D., Welsch, B. T., et al. 2020, *ApJS*, 248, 2  
Fisher, G. H., Welsch, B. T., & Abbett, W. P. 2012, *SoPh*, 277, 153  
Fisher, G. H., Welsch, B. T., Abbett, W. P., & Bercik, D. J. 2010, *ApJ*, 715, 242  
Forbes, T. G. 2000, *JGR*, 105, 23153  
Guo, J. H., Ni, Y. W., Guo, Y., et al. 2024, *ApJ*, 961, 140  
Guo, Y., Cheng, X., &

Ding, M. D. 2017, ScChD, 60, 1408 Guo, Y., Xia, C., Keppens, R., Ding, M. D., & Chen, P. F. 2019, ApJL, 870, L21 Guo, Y., Zhong, Z., Ding, M. D., et al. 2021, ApJ, 919, 39 Hayashi, K., Feng, X., Xiong, M., & Jiang, C. 2018, ApJ, 855, 11 Hayashi, K., Feng, X., Xiong, M., & Jiang, C. 2019, ApJL, 871, L28 Hou, Y. J., Zhang, J., Li, T., Yang, S. H., & Li, X. H. 2018, A&A, 619, 10 Inoue, S., Shiota, D., Bamba, Y., & Park, S.-H. 2018, ApJ, 867, 83 Jiang, C., Bian, X., Sun, T., & Feng, X. 2021, FrP, 9, 1 Jiang, C., & Feng, X. 2013, ApJ, 769, 144 Jiang, C., Feng, X., Guo, Y., & Hu, Q. 2022, Innov, 3, 100236 Jiang, C., Wu, S. T., Feng, X., & Hu, Q. 2016a, NatCo, 7, 11522 Jiang, C., Wu, S. T., Yurchyshyn, V., et al. 2016b, ApJ, 828, 62 Jiang, C., Zou, P., Feng, X., et al. 2018, ApJ, 869, 13 Kaneko, T., Park, S.-H., & Kusano, K. 2021, ApJ, 909, 155 Kazachenko, M. D., Fisher, G. H., & Welsch, B. T. 2014, ApJ, 795, 17 Kliem, B., Su, Y. N., van Ballegooijen, A. A., & DeLuca, E. E. 2013, ApJ, 779, 129 Lin, H., Kuhn, J. R., & Coulter, R. 2004, ApJ, 613, L177 Liu, L., Cheng, X., Wang, Y., et al. 2018, ApJL, 867, L5 Liu, Y. 2009, AnGeo, 27, 2771 Lumme, E., Pomoell, J., & Kilpua, E. K. 2017, SoPh, 292, 191 Metcalf, T. R., DeRosa, M. L., Schrijver, C. J., et al. 2008, SoPh, 247, 269 Moran, T., & Foukal, P. 1991, SoPh, 135, 179 Morosan, D. E., Carley, E. P., Hayes, L. A., et al. 2019, NatAS, 3, 452 November, L., & Simon, G. 1988, ApJ, 333, 427 Pomoell, J., Lumme, E., & Kilpua, E. 2019, SoPh, 294, 41 Price, D. J., Pomoell, J., Lumme, E., & Kilpua, E. K. 2019, A&A, 628, A114 Regnier, S. 2013, SoPh, 288, 481 Romano, P., Elmhamdi, A., & Kordi, A. S. 2019, SoPh, 294, 4 Schrijver, C. J., Derosa, M. L., Metcalf, T. R., et al. 2006, SoPh, 235, 161 Schuck, P. W. 2006, ApJ, 646, 1358 Schuck, P. W. 2008, ApJ, 683, 1134 Schuck, P. W., & Antiochos, S. K. 2019, ApJ, 882, 151 Tomczyk, S., Card, G. L., Darnell, T., et al. 2008, SoPh, 247, 411 Tóth, G. 2000, JCoPh, 161, 605 Verma, M. 2018, A&A, 612, A101 Wang, X., Jiang, C., & Feng, X. 2023, ApJL, 942, L41 Wang, X., Jiang, C., Feng, X., Duan, A., & Bian, X. 2022, ApJ, 938, 61 Webb, D. F., & Howard, T. A. 2012, LRSP, 9, 1 Wiegelmann, T. 2008, JGRA, 113, A03S02 Wiegelmann, T., & Sakurai, T. 2021, LRSP, 18, 1 Wien, W. 1916, AnP, 354, 842 Wu, S. T., Wang, A. H., Gary, G. A., et al. 2009, AdSpR, 44, 46 Wu, S. T., Wang, A. H., Liu, Y., & Hoeksema, J. T. 2006, ApJ, 652, 800 Yang, G., Xin, C., & MingDe, D. 2017, ScChD, 60, 1408 Yardley, S. L., Mackay, D. H., & Green, L. M. 2021, SoPh, 296, 10 Zhong, Y., Dai, Y., & Ding, M. D. 2021, ApJ, 916, 37 Zou, P., Jiang, C., Feng, X., et al. 2019, ApJ, 870, 97 Zou, P., Jiang, C., Wei, F., et al. 2020, ApJ, 890, 10

*Note: Figure translations are in progress. See original paper for figures.*

*Source: ChinaXiv — Machine translation. Verify with original.*