

Stellar Parameters, Extinction, and Distances for Stars in SMSS DR2 by the SPar Method (Post-print)

Authors: Mingxu Sun, Bingqiu Chen, Baokun Sun, Tao Wang, Zheng Yu, Baisong Zhang, Lin Zhang, Yuxi Bao, Guangya Zeng, Ming Yang and Wenyuan Cui

Date: 2025-06-13T16:51:30+00:00

Abstract

The availability of large data sets containing stellar parameters, distances, and extinctions for stars in the Milky Way, particularly within the Galactic disk, is essential for advancing our understanding of the Galaxy's stellar populations, structure, kinematics, and chemical evolution. In this study, we present a catalog of stellar parameters, including effective temperature (T_{eff}), metallicity ($[\text{Fe}/\text{H}]$), absolute magnitudes (M_G), distances (d), and reddening values ($E(\text{GBP} - \text{GRP})$), for a sample of 141 million stars from the SkyMapper Southern Survey. These parameters are derived using the SPar algorithm, which employs a fitting procedure to match multi-band photometric observations and estimate the stellar properties (T_{eff} , $[\text{Fe}/\text{H}]$, M_G , d , and $E(\text{GBP} - \text{GRP})$) on an individual star basis, following the methodology outlined in our previous work. This study successfully determines stellar parameters and extinction values simultaneously for stars located in high and low Galactic latitudes. The resulting stellar parameters and extinction values are in good agreement with results from other studies, demonstrating the robustness of our method. We derive a temperature dispersion of 195 K and a metallicity dispersion of 0.31 dex when comparing our results with spectroscopic data. The catalog produced in this work provides a valuable resource for future research on the Galactic metallicity distribution function, the structure of the Galaxy, three-dimensional extinction mapping, and other related astrophysical topics.

Full Text

Preamble

Research in Astronomy and Astrophysics, 25:057002 (15pp), 2025 May © 2025. National Astronomical Observatories, CAS and IOP Publishing Ltd. All rights, including for text and data mining, AI training, and similar technologies, are reserved. Printed in China. <https://doi.org/10.1088/1674-4527/adc5e1> CSTR: 32081.14.RAA.adc5e1 Stellar Parameters, Extinction, and Distances for Stars in SMSS DR2 by the SPar Method Mingxu Sun¹, Bingqiu Chen², Baokun Sun², Tao Wang², Zheng Yu², Baisong Zhang², Lin Zhang², Yuxi Bao², Guangya Zeng², Ming Yang³, and Wenyuan Cui¹ ¹ Department of Physics, Hebei Key Laboratory of Photophysics Research and Application, Hebei Normal University, Shijiazhuang 050024, China ² South-Western Institute for Astronomy Research, Yunnan University, Kunming 650500, China; bchen@ynu.edu.cn ³ Key Laboratory of Space Astronomy and Technology, National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100101, China Received 2024 October 9; revised 2025 March 4; accepted 2025 March 24; published 2025 May 28

Abstract

The availability of large data sets containing stellar parameters, distances, and extinctions for stars in the Milky Way, particularly within the Galactic disk, is essential for advancing our understanding of the Galaxy's stellar populations, structure, kinematics, and chemical evolution. In this study, we present a catalog of stellar parameters, including effective temperature (T_{eff}), metallicity ($[\text{Fe}/\text{H}]$), absolute magnitudes (M_G), distances (d), and reddening values ($E(\text{GBP} - \text{GRP})$), for a sample of 141 million stars from the SkyMapper Southern Survey. These parameters are derived using the SPar algorithm, which employs a fitting procedure to match multi-band photometric observations and estimate the stellar properties (T_{eff} , $[\text{Fe}/\text{H}]$, M_G , d , and $E(\text{GBP} - \text{GRP})$) on an individual star basis, following the methodology outlined in our previous work. This study successfully determines stellar parameters and extinction values simultaneously for stars located in high and low Galactic latitudes. The resulting stellar parameters and extinction values are in good agreement with results from other studies, demonstrating the robustness of our method. We derive a temperature dispersion of 195 K and a metallicity dispersion of 0.31 dex when comparing our results with spectroscopic data. The catalog produced in this work provides a valuable resource for future research on the Galactic metallicity distribution function, the structure of the Galaxy, three-dimensional extinction mapping, and other related astrophysical topics.

Key words: (ISM:) dust, extinction -Stars -stars: low-mass -(Galaxy:) solar neighborhood

1. Introduction

Understanding the formation and evolution of the Milky Way requires precise measurements of stellar atmospheric parameters, distances, and extinction for large samples of stars, particularly those in the Galactic disk. Such measurements allow us to characterize the properties of stellar populations and their spatial distributions across the Galaxy (Casagrande et al. 2011; Peng et al. 2013; Bland-Hawthorn & Gerhard 2016; Gaia Collaboration et al. 2018; Whitten et al. 2021).

Large-scale spectroscopic surveys have revolutionized the field by providing precise stellar parameters for tens of millions of stars. Notable examples include the Sloan Extension for Galactic Understanding and Exploration (SEGUE; Yanny et al. 2009), the Large Sky Area Multi-Object Fiber Spectroscopic Telescope (LAMOST; Luo et al. 2015), the Apache Point Observatory Galactic Evolution Experiment (APOGEE; Abdurro'uf et al. 2022), the Galactic Archaeology with HERMES (GALAH; Buder et al. 2021), and the Dark Energy Spectroscopic Instrument (DESI; DESI Collaboration et al. 2022). These surveys have provided high-quality data sets with broad coverage of parameter space, enabling the inference of stellar properties from photometric data. Complementing spectroscopic surveys that provide stellar properties, narrow- and medium-band photometric surveys, such as the SkyMapper Southern Survey (SMSS; Wolf et al. 2018; Onken et al. 2019), the Javalambre/Southern Photometric Local Universe Survey (J/S-PLUS; Cenarro et al. 2019), and the Sloan Digital Sky Survey (SDSS; York et al. 2000), have demonstrated the ability to estimate stellar parameters with high accuracy through their carefully designed filter sets.

Recent studies have demonstrated the efficacy of photometric methods for deriving stellar parameters. For example, Yuan et al. (2015a, 2015b) used empirical metallicity-dependent stellar loci to estimate photometric metallicity for 500,000 FGK dwarf stars in Stripe 82. Chen et al. (2019) derived intrinsic colors and extinction values for 23 million stars in the Galactic disk using optical and near-IR photometry, with spectroscopic data from LAMOST and SDSS serving as training samples. Extending the methods of Yuan et al. to red giants, Zhang et al. (2021) developed metallicity-dependent stellar loci to estimate metallicities from SDSS photometry, enabling the identification of metal-poor red giants. Leveraging the sensitivity of the SMSS *uvgriz* filters, Huang et al. (2019) constructed empirical relationships between photometric colors and stellar parameters, deriving accurate atmospheric parameters for approximately one million red giants. Building on this, Huang et al. (2022) estimated stellar parameters for 24 million stars using SMSS photometry, and Chiti et al. (2021) utilized a grid-based synthetic photometry approach to derive metallicities for over 250,000 stars from SMSS data. Other notable efforts include Xu et al. (2022), who estimated metallicities for 27 million stars using Gaia EDR3 photometry with LAMOST spectroscopic data as training samples, and Yang et al. (2022), who derived stellar parameters from J-PLUS photometry (Cenarro et al. 2019). For stars at high Galactic latitudes, extinction effects are minimal, allowing for pre-

cise parameter estimates using two-dimensional extinction maps (e.g., Schlegel et al. 1998).

However, in the low-latitude Galactic disk, where dust extinction is significant, simultaneous measurements of extinction and metallicity are essential to reduce systematic errors in parameter estimation (Andrae et al. 2022; Sun et al. 2023). This highlights the need for integrated approaches that can account for extinction effects while deriving stellar parameters.

In this study, we employ the SPar algorithm (Sun et al. 2023) to estimate stellar atmospheric parameters, extinction, and distances for a sample of 140 million stars from SMSS DR2. The SPar algorithm relies on empirical stellar libraries trained on LAMOST spectroscopic data (Luo et al. 2015). Our data set combines uvgriz photometry from SMSS with JHKS measurements from the Two Micron All Sky Survey (2MASS; Skrutskie et al. 2006), W1W2 photometry from the Wide-field Infrared Survey Explorer (WISE; Wright et al. 2010), and GGBPGRP data from Gaia DR3 (Gaia Collaboration et al. 2023). By matching observations to the stellar template library, we derive key stellar parameters, including effective temperature (T_{eff}), metallicity ($[\text{Fe}/\text{H}]$), absolute magnitude (M_G), distance (d), and extinction ($E(\text{GBP} - \text{GRP})$) for individual stars. Unlike previous algorithms such as StarHorse (Anders et al. 2019) and GSP-Phot (Andrae et al. 2022), which rely on theoretical stellar models that may introduce systematic uncertainties (Green et al. 2021), our approach is based on empirical relationships directly derived from observational data. By leveraging the sensitivity of the SMSS uv filters to metallicity and utilizing a robust training data set, we achieve accurate estimates of stellar parameters and extinction. Compared to prior studies using SMSS data, our work stands out for its larger sample size and the simultaneous estimation of stellar parameters and extinction, enabling precise characterization of stars in the Galactic disk and improving our understanding of the Galactic disk's structure and evolution.

2. Data

In this study, we utilize the uvgriz photometric data from SMSS DR2 (Onken et al. 2019; Wolf et al. 2018). SMSS is a hemispheric survey conducted at Siding Spring Observatory in Australia using the SkyMapper Telescope. The catalog contains a total of 505 million sources. The survey reaches depths of 19.7 to 21.7 mag across its six bands: u, v, g, r, i, and z. Compared to the single u band in SDSS, characterized by $(\lambda_{\text{cen}}/\text{FWHM}) = (358 \text{ nm}/55 \text{ nm})$, SMSS uses two separate filters in the blue: a violet v band (384 nm/28 nm) and a more ultraviolet u band (349 nm/42 nm). This configuration enhances the sensitivity to variations in metallicity. The internal photometric precision of SMSS DR2 is 1% for the u and v bands and 0.7% for the griz bands. Over 21,000 square degrees are covered in certain filters, with more than 7000 square degrees surveyed deeply across all six bands.

In Figure 1 [Figure 1: see original paper], we present the spatial distribution

of stars in SMSS within the Galactic Cartesian coordinate system. The X-axis points toward the Galactic center, the Y-axis toward the direction of Galactic rotation, and the Z-axis toward the north Galactic pole. The XYZ values for individual stars are calculated using their Galactic coordinates (l , b) and distances (d), where distances are simply derived as the inverse of Gaia DR3 parallaxes ($d = 1/\varpi$). Stars without Gaia parallaxes or with negative parallaxes are excluded, accounting for approximately 22% of the total sample. Notably, distance uncertainties increase significantly for stars at greater distances due to the large errors in Gaia parallax measurements. This distribution clearly traces the disk structure of the Milky Way.

To enhance the data set, we combine SMSS data with infrared photometry from the 2MASS and WISE catalogs. The 2MASS survey, conducted across the entire sky, uses three filters: J, H, and K_S (Skrutskie et al. 1997). Its source catalog contains 470 million objects. The WISE survey, using the W1, W2, W3, and W4 bands, provides additional infrared data. For this study, we use the AllWISE Source Catalog (Kirkpatrick et al. 2014), which contains 748 million sources. To ensure high data quality, we limit our analysis to the W1 and W2 bands due to their superior sensitivity and angular resolution compared to W3 and W4. Furthermore, we incorporate Gaia DR3 data (Gaia Collaboration et al. 2023), which provides essential astrometric parameters, including positions, parallaxes, and proper motions, as well as photometry in the G, GBP, and GRP bands. Gaia DR3 encompasses 1.8 billion sources, making it an invaluable resource for cross-matching and parameter estimation.

The SMSS data is cross-matched with 2MASS, AllWISE, and Gaia using a matching radius of 1.5'', resulting in the creation of the SMSS-2MASS-AllWISE-Gaia (SSAG) catalog. To ensure data quality, we apply the following selection criteria: (1) sources must have measurements in the SMSS gri bands and at least one additional band, and (2) photometric errors in the SMSS gri bands and at least one additional band must be less than 0.1 mag. After applying these criteria, the SSAG catalog includes a total of 188,464,828 sources.

Figure 2 [Figure 2: see original paper] presents a histogram illustrating the contributions of different catalogs to the merged data set. Our selection criteria, which require all SMSS sources to have complete observations in the gri bands and photometric errors below 0.1 mag in these bands, exclude a large portion of the original SMSS sources. Despite this, the majority of SMSS sources in the final sample are successfully matched with Gaia data, and over half of the sources are also cross-matched with 2MASS and WISE.

Figure 3 [Figure 3: see original paper] displays the color-magnitude diagrams ($g - r$ versus i) for the original SMSS catalog and the merged SSAG catalog. The overall distributions in both diagrams show no significant differences, indicating that the merging process preserves the consistency and integrity of the data. The $g - r$ values primarily range from -2.5 to 2.5 mag, while the i magnitudes span from 9 to 21 mag. This consistency suggests that, while the sample size is reduced due to the selection criteria, the proportion of different stellar

populations within the sample remains unchanged.

3. Method

The methodology employed in this study is based on the SPar algorithm described by Sun et al. (2023). This algorithm establishes relationships between stellar parameters—effective temperature (T_{eff}), metallicity ($[\text{Fe}/\text{H}]$), and absolute magnitude in the G band (M_{G})—and the absolute magnitudes across all filters using empirical stellar templates. For each star in the SSAG catalog, we perform a minimum χ^2 fitting of the multiband photometry and parallax data to derive initial parameter estimates. These estimates are then refined using a Markov Chain Monte Carlo (MCMC) analysis, which provides the final stellar parameters along with their uncertainties.

Below, we provide a concise summary of the methodology, while further details can be found in Sun et al. (2023):

1. We construct an empirical stellar library by selecting a sample of stars from LAMOST and Gaia DR3 that have well-determined atmospheric parameters, distances, and extinction values. This sample is supplemented with optical and near-infrared photometry from SMSS, 2MASS, and WISE. The library includes stellar parameters and the corresponding absolute magnitudes for each filter.
2. A Random Forest Regression algorithm is applied to the stellar library to map the relationships between the stellar parameters (T_{eff} , $[\text{Fe}/\text{H}]$, and M_{G}) and the absolute magnitudes in each passband. This process generates empirical stellar templates used for parameter estimation.
3. Distance estimation is performed using observed magnitudes in optical bands and the empirical stellar templates. For a given star, the stellar templates, combined with assumed values of T_{eff} , $[\text{Fe}/\text{H}]$, M_{G} , and $E(\text{GBP} - \text{GRP})$, are used to predict the “distance modulus-corrected” magnitudes M_x in the x filter, using the relation: $M_x(T_{\text{eff}}, [\text{Fe}/\text{H}], M_{\text{G}}) + A_x(E(\text{GBP} - \text{GRP}))$, where $A_x(E(\text{GBP} - \text{GRP}))$ represents the extinction in filter x . The distance modulus is then derived by subtracting from the observed magnitudes m_x : $m_x = M_x + A_x + \mu$. This calculation is restricted to optical filters, specifically Gaia G, GBP, GRP, and SMSS gri, because the Gaia magnitudes have high photometric accuracies and SMSS gri bands are available for all sources in the final merged catalog. The standard magnitude relation, $m_x = M_x + A_x + \mu$, is then used to simulate the apparent magnitudes in the individual passbands.
4. The observed magnitudes across all available filters and the observed parallaxes are modeled as a function of four free parameters: T_{eff} , $[\text{Fe}/\text{H}]$, M_{G} , and $E(\text{GBP} - \text{GRP})$. Initially, a coarse grid search is performed using a minimum χ^2 method to identify the best-fit values of T_{eff} , $[\text{Fe}/\text{H}]$,

M_G , and $E(GBP - GRP)$ for each star in the SSAG catalog.

5. Stars with acceptable fits, defined as those with $\chi^2 < 10$, are selected. The best-fit parameters from the χ^2 minimization serve as initial values for an MCMC analysis. This step refines the parameter estimates and provides their final values along with associated uncertainties.

4. Results and Discussion

In this section, we present the derived stellar parameters for our sample, including effective temperatures (T_{eff}), metallicities ($[Fe/H]$), absolute magnitudes (M_G), reddening values ($E(GBP - GRP)$), and distances (d). We also analyze and discuss the overall distribution of these parameters, as well as the spatial distribution of the stars within the sample.

Figure 4 [Figure 4: see original paper] illustrates the observational data used in this study, including multi-band photometry and Gaia parallax measurements (when available), along with the χ^2 distribution for all stars in the SSAG sample. As expected, the χ^2 distribution peaks near 2 and exhibits an extended tail. Of the 188,464,828 stars in the SSAG catalog, approximately 75% (140,599,779 stars) have χ^2 values below 10, which we use as a threshold for selecting high-quality fits. The stellar parameters for these stars were refined using the MCMC analysis. For the sources with $\chi^2 < 10$, the majority have more than eight independent measurements, combining magnitudes from different photometric filters and Gaia parallax. Nearly all stars in this subset have observations in the SMSS gri bands and Gaia G, GBP, and GRP bands. Additionally, approximately 25 million stars (18%) have observations in the SMSS u band, while 28 million stars (20%) have data in the SMSS v band. This broad coverage across multiple filters enables robust parameter estimation.

A detailed table containing the derived parameters for the 141 million stars with $\chi^2 < 10$ is available on the website at doi:10.12149/101372. This publicly accessible data set provides a valuable resource for further studies of stellar populations and Galactic structure.

Figure 5 [Figure 5: see original paper] illustrates the parameter distributions for the 141 million stars in our catalog, including effective temperature (T_{eff}), metallicity ($[Fe/H]$), absolute magnitude (M_G), extinction ($E(GBP - GRP)$), and distance. The majority of stars have T_{eff} values between 5000 and 6000 K, reflecting the dominance of mid-type stars in the sample. Metallicity values ($[Fe/H]$) are primarily concentrated in the range -0.4 to 0.2 dex, and M_G values are clustered between 3 and 6 mag. For the reddening values ($E(GBP - GRP)$), most stars exhibit values between 0 and 0.3 mag. Distances are predominantly within the range of 1000–3000 pc.

To emphasize the uniqueness of our stellar sample, Table 1 provides a comparative overview of the stellar parameter ranges from this study and those from Huang et al. (2022), Xu et al. (2022), and Zhang et al. (2023). Compared to

Huang et al. (2022) and Xu et al. (2022), our work has a significantly larger sample size and benefits from the simultaneous estimation of stellar parameters and extinction, which allows for more precise characterization of stellar properties, particularly in the Galactic disk. Unlike the sample in Zhang et al. (2023), which includes stars from brighter magnitude ranges, our catalog extends to much fainter stars, providing a complementary data set for probing stellar populations at further distances.

In this section, we will discuss the precision of the resultant effective temperature (T_{eff}), metallicity ($[\text{Fe}/\text{H}]$), extinction ($E(\text{GBP} - \text{GRP})$), and distance for the stars in our catalog. To evaluate the accuracy of our parameter estimates, we compare our results with those from previous studies, including large-scale spectral surveys. The parameter distributions are examined spatially across the Milky Way to identify trends and correlations, providing a foundation for further exploration of stellar populations and Galactic structure. We also compare our findings with related studies to highlight the scientific implications of our work and identify potential avenues for future research.

4.1. Discussion of T_{eff} and $[\text{Fe}/\text{H}]$

In this section, we compare the derived effective temperatures (T_{eff}) and metallicities ($[\text{Fe}/\text{H}]$) with results from multiple spectroscopic surveys, including LAMOST DR11 (L11; Luo et al. 2015), GALAH DR3 (G3; Buder et al. 2021), and APOGEE DR17 (A17; Abdurro' uf et al. 2022). The training sample used in this study is based on LAMOST DR8, which partially overlaps with L11 but remains independent from G3 and A17. These surveys are treated as benchmarks to assess the accuracy of our results. Additionally, we include three more catalogs for comparison: Gaia23 (Gaia Collaboration et al. 2023), Z23 (Zhang et al. 2023), and H22 (Huang et al. 2022). Cross-matching was performed between the selected catalogs and spectroscopic surveys, and we use the common sources shared between these catalogs and the spectroscopic surveys for comparison.

Figure 6 [Figure 6: see original paper] compares the T_{eff} values derived in this study with those from the spectroscopic surveys. Our results show good consistency with L11, G3, and A17, with median differences ($\bar{\Delta}$) and standard deviations (σ) of $\bar{\Delta} = -52$ K and $\sigma = 195$ K for L11, $\bar{\Delta} = 21$ K and $\sigma = 193$ K for G3, and $\bar{\Delta} = -18$ K and $\sigma = 187$ K for A17. The dispersion in differences increases with effective temperature. For G3, minor vertical stripes are visible in the comparison, which are also present when comparing Gaia23, Z23, and H22 with G3. This could indicate systematic effects related to the G3 observation template. Additionally, G3 and A17 exhibit two distinct parameter clustering regions, likely due to selection effects in these surveys. When compared with Gaia23, Z23, and H22, our T_{eff} results have accuracy comparable to Z23 and H22, and slightly better than Gaia23. Notably, Gaia23 shows a systematic overestimation of T_{eff} around 5000 K and at higher temperatures compared to the spectroscopic surveys. In particular, Gaia23 exhibits distinct horizontal stripes in comparison with L11, G3, and A17. Above 6600 K, deviations in

Z23 and H22 become apparent, primarily due to a few outlier points. H22, for instance, only includes stars with T_{eff} below 6600 K, which limits comparisons in this range.

For $[\text{Fe}/\text{H}]$, our results also exhibit good agreement with the spectroscopic surveys, as shown in Figure 7 [Figure 7: see original paper]. The median differences and standard deviations are $\mu = -0.02$ dex and $\sigma = 0.33$ dex for L11, $\mu = -0.01$ dex and $\sigma = 0.27$ dex for G3, and $\mu = -0.04$ dex and $\sigma = 0.33$ dex for A17. The dispersion increases at lower metallicities, particularly for $[\text{Fe}/\text{H}] < -1$, where the differences become more pronounced. For $[\text{Fe}/\text{H}] < -2$, the differences are even larger, though this region contains relatively few sources. Our $[\text{Fe}/\text{H}]$ results have accuracy comparable to those of Z23 and H22 and better than Gaia23. In the metallicity range of -0.5 – 0 dex, Gaia23 tends to underestimate values compared to spectroscopic surveys, whereas our results, Z23, and H22 show better agreement. Gaia23 also exhibits horizontal stripes in the $[\text{Fe}/\text{H}]$ comparisons, similar to those seen in the T_{eff} results. In the comparison plots of G23 and Z23, subsets of stars exhibit significant discrepancies in $[\text{Fe}/\text{H}]$ when compared to spectroscopic results. However, such discrepancies are absent in our results and those of H22. Upon further examination, we identified that the stars exhibiting significant deviations in metallicity in the G23 and Z23 studies are primarily disk dwarf stars. However, they account for only about 1% of the dwarf star sample in the common data set between these studies and the spectroscopic surveys. No significant trends in distance or extinction were identified among them. These deviations are likely statistical in nature, resulting from the Gaussian tail of measurement errors.

Figures 6 and 7 demonstrate that our results are in good consistent with the three spectroscopic surveys. When combining all three surveys, the typical median and dispersion values are $\mu = -19$ K and $\sigma = 197$ K for T_{eff} , and $\mu = -0.02$ dex and $\sigma = 0.31$ dex for $[\text{Fe}/\text{H}]$. In Figure 8 [Figure 8: see original paper], we present the median differences across various ranges when comparing our results with the spectroscopic surveys. For T_{eff} , there are no significant trends in the median differences (ΔT_{eff}) across different T_{eff} ranges. Our results closely align with those of Z23 and H22 and show slightly better consistency than Gaia23. For $[\text{Fe}/\text{H}]$, our results maintain a similar level of agreement with Z23 and H22 when $[\text{Fe}/\text{H}] > -1$ dex. However, for $[\text{Fe}/\text{H}] < -2$ dex, larger discrepancies are observed compared to the spectroscopic surveys.

When all measurements from the input catalogs are available ($n = 15$), our results show improved agreement with the spectroscopic surveys, with differences of $\mu = -13$ K and $\sigma = 176$ K for T_{eff} , and $\mu = -0.02$ dex and $\sigma = 0.25$ dex for $[\text{Fe}/\text{H}]$. However, not all stars have complete measurements across all input catalogs. To evaluate the effect of missing data, we randomly selected 10,000 sources with complete measurements ($n = 15$) and tested how the absence of certain input catalogs impacts the derived parameters. Figure 9 [Figure 9: see original paper] illustrates the median differences between our results and the spectroscopic surveys when specific input catalogs are excluded. For T_{eff} , the

deviations generally increase as T_{eff} decreases, although the absence of any single input catalog does not substantially affect the overall deviations across the full T_{eff} range. In contrast, for $[\text{Fe}/\text{H}]$, the deviations become more pronounced as $[\text{Fe}/\text{H}]$ decreases. Notably, the absence of SMSS uv data leads to significantly larger deviations in $[\text{Fe}/\text{H}]$, while the absence of other input catalogs has less impact on $[\text{Fe}/\text{H}]$ deviations.

To investigate parameter correlations, we selected an example star (SMSS id = 62096272) and performed an MCMC analysis using 32 walkers and 5000 steps. Figure 10 [Figure 10: see original paper] shows the corner plot with one- and two-dimensional projections of the posterior distributions of the parameters. The contours illustrate covariances, while the histograms display Gaussian-like distributions. We observe that variations in effective temperature (T_{eff}) significantly affect parameters such as extinction ($E(\text{GBP} - \text{GRP})$) and absolute magnitude (M_G), but have a smaller impact on metallicity ($[\text{Fe}/\text{H}]$). We also note a minor degeneracy between extinction and metallicity, as metallicity can influence stellar color.

4.2. Discussion of $E(\text{GBP} - \text{GRP})$ and d

Figure 11 [Figure 11: see original paper] presents a comparison of derived $E(\text{GBP} - \text{GRP})$ values from this study with those obtained using the star-pair algorithm based on spectroscopic survey parameters (Sun et al. 2023). For reference, we also include comparisons using the $E(\text{GBP} - \text{GRP})$ values from Gaia23 and Z23. All three data sets show good agreement with results from the spectroscopic data, though differences in the extinction laws adopted lead to variations in the slopes of the comparisons. Specifically, the slope for our results compared to the star-pair method is 1.0, while it is 1.2 for Gaia23 and 0.86 for Z23. In terms of residual dispersion, our results (0.05 mag) are comparable to those of Z23 (0.06 mag), whereas Gaia23 shows a larger residual dispersion of 0.11 mag.

We further compare our $E(\text{GBP} - \text{GRP})$ values with the $E(B - V)$ estimates from G19 (Green et al. 2019) and SFD98 (Schlegel et al. 1998), as shown in Figure 12 [Figure 12: see original paper]. Strong linear correlations are observed, with residual variances of 0.14 mag for G19 and 0.13 mag for SFD98. For the G19 comparison, $E(B - V)/E(\text{GBP} - \text{GRP}) = 0.68$, consistent with previous studies (Chen et al. 2019; Sun et al. 2021). When comparing with SFD98, we limit the sample to stars located more than 200 pc above the Galactic plane. In this case, the color excess ratio is approximately 0.85, slightly higher than the G19 value. This difference may be attributed to SFD98's known 14% overestimate of $E(B - V)$ (Schlafly et al. 2010; Yuan et al. 2013).

To evaluate the accuracy of our distance estimates (d), Figure 13 [Figure 13: see original paper] compares our results with parallax and distance measurements from Gaia DR3 (Bailer-Jones et al. 2021). Of the total sample, 99.9% (140,450,761 sources) have Gaia parallax measurements. For most stars, our dis-

tance estimates are in good agreement with Gaia's values, with approximately 62% of the sources showing relative discrepancies of less than 30%. However, at larger distances, our estimates tend to be slightly higher than those from Gaia, likely due to the increasing uncertainties in parallax measurements over greater distances.

4.3. Flags

Table 2 provides an overview of the contents of the final catalog, including detailed descriptions of all fields. To ensure that users can assess the reliability of our derived parameters, we have implemented a comprehensive flagging system. These flags indicate the quality and completeness of the input data and are essential for assessing the trustworthiness of our results. The key flags include `flag_Fe/H` (indicating the availability of the high-quality SMSS *v* band in the input catalogs), `n` (the number of input measurements, such as multi-band photometry and Gaia parallax), Chi-square (the minimal χ^2 value), `bands` (the names of the photometric bands used), and `flag_par` (indicating the inclusion of Gaia parallax).

For `flag_Fe/H`, a value of 1 indicates the presence of the SMSS *v* band with an error less than 0.1 mag in the input catalogs, contributing to more reliable metallicity ($[Fe/H]$) measurements. If the *v* band is absent or does not meet this quality criterion, `flag_Fe/H` is set to 0. The flag `n` represents the total number of input measurements, with a maximum value of 15, which includes multi-band photometry and Gaia parallax. While a higher `n` typically suggests greater accuracy, the reliability of the results also depends on the type and quality of the observations. The Chi-square value, as described in Section 3 and Sun et al. (2023), represents the quality of the parameter fitting, with our catalog restricting χ^2 to values below 10 to ensure reliability. The `bands` field lists the specific photometric bands used in the input catalogs. These include Gaia G, GBP, and GRP; 2MASS J, H, and KS; WISE W1 and W2; and SMSS uvgriz. Finally, `flag_par` indicates whether Gaia parallax measurements were included in the input, with a value of 1 for inclusion and 0 for exclusion. This provides users with an additional measure of the completeness of the input data used to derive stellar parameters.

4.4. Galactic Metallicity Distribution

The inclusion of the SMSS uv bands in the input star catalogs significantly improves the accuracy of $[Fe/H]$ measurements. For this study, we focused on stars from the input catalogs that include the SMSS *v* band with an error below 0.1 mag (indicated by `flag_Fe/H` = 1) to analyze the metallicity distribution of the Milky Way. To validate our results, we compared the derived $[Fe/H]$ values with those from Huang et al. (2022) (H22). As shown in Figure 14 [Figure 14: see original paper], the $[Fe/H]$ values from this study are in excellent agreement with those of H22, exhibiting a median difference of 0 dex and a dispersion of

0.35 dex. Importantly, our data set includes $[\text{Fe}/\text{H}]$ measurements for stars in low Galactic latitude regions, which were not covered in H22's work.

The spatial distribution of median metallicity ($[\text{Fe}/\text{H}]$) across the R-Z plane is presented in Figure 15 [Figure 15: see original paper]. For this analysis, we selected stars with high-quality $[\text{Fe}/\text{H}]$ and distance measurements, applying stringent criteria: relative distance errors below 20% ($|\Delta d|/d < 0.2$), metallicity uncertainties less than 0.15 dex ($|\Delta[\text{Fe}/\text{H}]| < 0.15$ dex), and the presence of the SMSS v band ($\text{flag}_{[\text{Fe}/\text{H}]} = 1$). The figure highlights the Galactic metallicity gradient, with a metal-rich region concentrated in the Galactic disk at lower vertical distances (Z). As Z increases, there is a clear transition to the metal-poor populations characteristic of the Galactic halo.

The metallicity distribution reveals radial variations within the Galactic disk. In the inner disk ($R < 8$ kpc), metallicity declines with increasing radius, while the outer disk displays significant flaring, characterized by metal-rich stars at higher vertical distances. However, near the midplane ($Z \approx 0$ kpc) at $R = 14$ – 20 kpc, the average metallicity is not as high. This discrepancy is likely due to significant extinction, which reduces the number of observable stars and increases metallicity measurement uncertainties. Since our metallicity template library has an upper limit, metal-rich stars near this boundary may be misclassified as having lower metallicities. The smaller number of stars in this region further exacerbates this bias. A more detailed investigation of this phenomenon has not been undertaken in this paper due to its scope limitations. Nonetheless, the results presented here offer a comprehensive view of the Milky Way's stellar metallicity distribution, providing valuable insights into the Galaxy's chemical evolution. Our catalog serves as a robust foundation for future studies of the Galactic metallicity distribution function and related astrophysical topics.

4.5. The Dust Extinction Distribution in the Southern Sky

We present new Galactic extinction maps in the top panels of Figure 16 [Figure 16: see original paper], constructed using the color excess $E(\text{GBP} - \text{GRP})$ derived in this study. To ensure high accuracy, these maps include stars with a color excess uncertainty below 0.05 mag and a fractional distance uncertainty of $|\Delta d|/d < 0.2$.

Figures 16 and 17 compare the extinction maps from this work with those from Guo et al. (2021). In Figure 16, the extinction distribution for nearby stars within $d < 0.5$ kpc highlights the intricate structures of local molecular clouds. When extended to $d < 5$ kpc, the maps reveal the broader features of the Galactic dust disk. Covering an area of approximately 25,529 square degrees, these maps provide a detailed view of the dust distribution and confirm many of the structures reported by Guo et al. (2021).

Figure 17 [Figure 17: see original paper] focuses on the region $340^\circ < l < 370^\circ$ and $-30^\circ < b < 30^\circ$, comparing the extinction maps over this specific area. The results from our study are consistent with those of Guo et al. (2021) in

terms of the observed dust structures. However, our maps feature significantly more complete sky coverage, enabling a more comprehensive characterization of the Galactic disk. Additionally, the larger data set used in this study allows for higher spatial resolution and greater depth, enhancing the reliability of the extinction measurements.

5. Summary

In this study, we analyzed a sample of stars from the SMSS DR2 and applied the SPar algorithm to derive stellar parameters, including effective temperature (T_{eff}), metallicity ($[\text{Fe}/\text{H}]$), absolute magnitude (M_G), extinction ($E(\text{GBP} - \text{GRP})$), and distance (d) for 141 million stars. This work leverages the extensive multi-band photometric data from SMSS uvgriz, 2MASS JHKs, WISE W1W2, and Gaia GGBPGRP, complemented by Gaia parallaxes when available. The SPar algorithm employs a fitting procedure to match these observations against a stellar template library trained on data from LAMOST and Gaia, enabling precise determination of stellar physical parameters, extinction, and distances for individual stars. The resulting catalog is publicly accessible at doi:10.12149/101372.

Our work represents a significant advancement in the characterization of stellar parameters for a vast region of the Galactic disk. Compared to previous studies within the SMSS framework, this study has significantly expanded both the sample size and the sky coverage. The derived stellar parameters, particularly T_{eff} and $[\text{Fe}/\text{H}]$, exhibit excellent agreement with values from spectroscopic surveys, with deviations characterized by a dispersion of 195 K for T_{eff} and 0.31 dex for $[\text{Fe}/\text{H}]$. These results are consistent with those reported in Z23 and H22, and show better precision than those from G23. Additionally, reddening estimates ($E(\text{GBP} - \text{GRP})$) and distance (d) align well with prior studies, further validating the robustness of our approach.

The new data set presented in this work offers a valuable resource for advancing our understanding of Galactic stellar populations, the interstellar dust distribution, and the chemical evolution of the Milky Way. Furthermore, the successful application of the SPar algorithm to such a large data set demonstrates its potential for future large-scale surveys, such as Mephisto and the Chinese Space Station Telescope (CSST) optical survey. These surveys promise to provide even larger data sets, enabling the determination of stellar physical parameters, extinction, and distances on an unprecedented scale.

Acknowledgments

We would like to thank the referee for providing us with detailed and constructive feedback that has significantly enhanced the quality of the manuscript. This work is partially supported by the National Natural Science Foundation of China (grants 12203016, 12173034, 12322304, and 12173013), Natural Science Foundation of Hebei Province (grants A2022205018, A2021205006, and 226Z7604G),

Yunnan University grant C619300A034, and Science Foundation of Hebei Normal University (grant L2022B33). W.Y.C. acknowledges the science research grants from the China Manned Space Project. M.X.S. acknowledges the support of Physics Postdoctoral Research Station at Hebei Normal University. We acknowledge the science research grants from the China Manned Space Project with Nos. CMS-CSST-2021-A09, CMS-CSST-2021-A08, and CMS-CSST-2021-B03.

This research made use of the cross-match service provided by CDS, Strasbourg. Guo Shou Jing Telescope (the Large Sky Area Multi-Object Fiber Spectroscopic Telescope LAMOST) is a National Major Scientific Project built by the Chinese Academy of Sciences. Funding for the project has been provided by the National Development and Reform Commission. LAMOST is operated and managed by the National Astronomical Observatories, Chinese Academy of Sciences.

This work presents results from the European Space Agency (ESA) space mission Gaia. Gaia data are being processed by the Gaia Data Processing and Analysis Consortium (DPAC). Funding for the DPAC is provided by national institutions, in particular the institutions participating in the Gaia MultiLateral Agreement (MLA). The Gaia mission website is <https://www.cosmos.esa.int/gaia>. The Gaia archive website is <https://archives.esac.esa.int/gaia>.

The national facility capability for SkyMapper has been funded through ARC LIEF grant LE130100104 from the Australian Research Council, awarded to the University of Sydney, the Australian National University, Swinburne University of Technology, the University of Queensland, the University of Western Australia, the University of Melbourne, Curtin University, Monash University, and the Australian Astronomical Observatory. SkyMapper is owned and operated by The Australian National University's Research School of Astronomy and Astrophysics. The survey data were processed and provided by the SkyMapper Team at ANU. The SkyMapper node of the All-Sky Virtual Observatory (ASVO) is hosted at the National Computational Infrastructure (NCI). Development and support of the SkyMapper node of the ASVO has been funded in part by Astronomy Australia Limited (AAL) and through the Australian Government's Education Investment Fund (EIF) and National Collaborative Research Infrastructure Strategy (NCRIS), particularly the National eResearch Collaboration Tools and Resources (NeCTAR) and the Australian National Data Service (ANDS) Projects.

This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration.

Mingxu Sun Bingqiu Chen Baisong Zhang Ming Yang Wenyuan Cui ORCID iDs
<https://orcid.org/0000-0002-2473-9948> <https://orcid.org/0000-0003-2472-4903>
<https://orcid.org/0009-0003-2888-6317> <https://orcid.org/0000-0001-8247-4936>
<https://orcid.org/0000-0003-1359-9908>

References

Abdurro' uf, Accetta, K., Aerts, C., et al. 2022, ApJS, 259, 35 Anders, F., Khalatyan, A., Chiappini, C., et al. 2019, A&A, 628, A94 Andrae, R., Fouesneau, M., Sordo, R., et al. 2022, A&A, 674, A27 Bailer-Jones, C. A. L., Rybizki, J., Fouesneau, M., Demleitner, M., & Andrae, R. 2021, AJ, 161, 147 Bland-Hawthorn, J., & Gerhard, O. 2016, ARA&A, 54, 529 Buder, S., Sharma, S., Kos, J., et al. 2021, MNRAS, 506, 150 Casagrande, L., Schönrich, R., Asplund, M., et al. 2011, A&A, 530, A138 Cenarro, A. J., Moles, M., Cristóbal-Hornillos, D., et al. 2019, A&A, 622, A176 Chen, B. Q., Huang, Y., Yuan, H. B., et al. 2019, MNRAS, 483, 4277 Chiti, A., Frebel, A., Mardini, M. K., et al. 2021, ApJS, 254, 31 DESI Collaboration, Abareshi, B., Aguilar, J., et al. 2022, AJ, 164, 207 Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, A&A, 616, A1 Gaia Collaboration, Vallenari, A., Brown, A. G. A., et al. 2023, A&A, 674, A1 Green, G. M., Rix, H.-W., Tschesche, L., et al. 2021, ApJ, 907, 57 Green, G. M., Schlafly, E., Zucker, C., Speagle, J. S., & Finkbeiner, D. 2019, ApJ, 887, 93 Guo, H. L., Chen, B. Q., Yuan, H. B., et al. 2021, ApJ, 906, 47 Huang, Y., Beers, T. C., Wolf, C., et al. 2022, ApJ, 925, 164 Huang, Y., Chen, B. Q., Yuan, H. B., et al. 2019, ApJS, 243, 7 Kirkpatrick, J. D., Schneider, A., Fajardo-Acosta, S., et al. 2014, ApJ, 783, 122 Luo, A. L., Zhao, Y.-H., Zhao, G., et al. 2015, RAA, 15, 1095 Onken, C. A., Wolf, C., Bessell, M. S., et al. 2019, PASA, 36, e033 Peng, X., Du, C., Wu, Z., Ma, J., & Zhou, X. 2013, MNRAS, 434, 3165 Schlafly, E. F., Finkbeiner, D. P., Schlegel, D. J., et al. 2010, ApJ, 725, 1175 Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525 Skrutskie, M. F., Schneider, S. E., Stiening, R., et al. 1997, in *Astrophysics and Space Science Library, The Impact of Large Scale Near-IR Sky Surveys*, ed. F. Garzon et al., Vol. 210 (Berlin: Springer), 25 Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, AJ, 131, 1163 Sun, M., Chen, B., Guo, H., et al. 2023, AJ, 166, 126 Sun, M., Jiang, B., Yuan, H., & Li, J. 2021, ApJS, 254, 38 Whitten, D. D., Placco, V. M., Beers, T. C., et al. 2021, ApJ, 912, 147 Wolf, C., Onken, C. A., Luvaul, L. C., et al. 2018, PASA, 35, e010 Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, AJ, 140, 1868 Xu, S., Yuan, H., Niu, Z., et al. 2022, ApJS, 258, 44 Yang, L., Yuan, H., Xiang, M., et al. 2022, A&A, 659, A181 Yanny, B., Rockosi, C., Newberg, H. J., et al. 2009, AJ, 137, 4377 York, D. G., Adelman, J., Anderson, J. E. J., et al. 2000, AJ, 120, 1579 Yuan, H. B., Liu, X. W., & Xiang, M. S. 2013, MNRAS, 430, 2188 Yuan, H., Liu, X., Xiang, M., Huang, Y., & Chen, B. 2015a, ApJ, 799, 134 Yuan, H., Liu, X., Xiang, M., Huang, Y., & Chen, B. 2015b, ApJ, 803, 13 Zhang, R.-Y., Yuan, H.-B., Liu, X.-W., et al. 2021, RAA, 21, 319 Zhang, X., Green, G. M., & Rix, H.-W. 2023, MNRAS, 524, 1855

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv –Machine translation. Verify with original.