

Two-proton radioactivity from excited states within the generalized liquid drop model

Authors: Long, Mr. Yi, Wang, Mr. Yue-Hua, Zhan, Mr. Dong, Li, Prof. Shuang, Deng, Prof. Jun-Gang, Mu, Dr. Lin, Cheng, Dr. Jun-Hao, Zhang, Dr. Hongfei, Deng, Prof. Jun-Gang

Date: 2026-04-23T09:39:43+00:00

Abstract

Unlike ground-state two-proton ($2p$) radioactivity, excited-state $2p$ emission is typically characterized by high decay energy and angular momentum, making it an important probe for studying the nuclear structure of extremely proton-rich nuclei. However, describing excited-state $2p$ radioactivity half-lives also poses a challenge to theoretical models. To examine whether the generalized liquid drop model (GLDM) with the improved proximity energy (Prox. 77-Set 13), optimized using ground-state $2p$ radioactivity experimental data, can be extended to describe excited-state $2p$ radioactivity, this work calculates the half-lives for $^{14}\text{O}^*$, $^{17}\text{Ne}^*$, $^{18}\text{Ne}^*$, $^{29}\text{S}^*$, and $^{94}\text{Ag}^*$. The present calculations are compared with experimental data and calculations obtained from the effective liquid drop model (ELDM), the unified fission model (UFM), and the Coulomb and proximity potential model (CPPM). The results calculated by this work are in good agreement with both the experimental data and the results of other theoretical models, validating the applicability of the GLDM with improved proximity energy for describing excited-state $2p$ radioactivity. This work also systematically investigates the dependence of excited-state $2p$ radioactivity half-life on decay energy and angular momentum, which will shed new light on the properties of exotic proton-rich nuclei.

Full Text

Preamble

Two-proton radioactivity from excited states within the generalized liquid drop model* Yi Long,^{1, 2} Yue-Hua Wang,^{1, 2} Dong Zhan,^{1, 2} Shuang Li,^{1, 2} Jun-Gang Deng,^{1, 2}, † Lin Mu,^{1, 2}, ‡ Jun-Hao Cheng,³ § and Hong-Fei Zhang⁴, ¶

College of Mathematics and Physics, China Three Gorges University, Yichang 443002, People's Republic of China Center for Astronomy and Space Sciences, China Three Gorges University, Yichang 443002, People's Republic of China Southwestern Institute of Physics, Chengdu, 610003, People's Republic of China School of Physics, Xi'an Jiaotong University, Xi'an 710049, People's Republic of China Unlike ground-state two-proton (2p) radioactivity, excited-state 2p emission is typically characterized by high decay energy and angular momentum, making it an important probe for studying the nuclear structure of extremely proton-rich nuclei. However, describing excited-state 2p radioactivity half-lives also poses a challenge to theoretical models. To examine whether the generalized liquid drop model (GLDM) with the improved proximity energy (Prox. 77-Set 13), optimized using ground-state 2p radioactivity experimental data, can be extended to describe excited-state 2p radioactivity, this work calculates the half-lives for $^{14}\text{O}^*$, $^{17}\text{Ne}^*$, $^{18}\text{Ne}^*$, $^{19}\text{S}^*$, and $^{94}\text{Ag}^*$. The present calculations are compared with experimental data and calculations obtained from the effective liquid drop model (ELDM), the unified fission model (UFM), and the Coulomb and proximity potential model (CPPM). The results calculated by this work are in good agreement with both the experimental data and the results of other theoretical models, validating the applicability of the GLDM with improved proximity energy for describing excited-state 2p radioactivity. This work also systematically investigates the dependence of excited-state 2p radioactivity half-life on decay energy and angular momentum, which will shed new light on the properties of exotic proton-rich nuclei.

Keywords

2p radioactivity, generalized liquid drop model, half-life, excited-state

INTRODUCTION

Two-proton (2p) radioactivity is a unique decay mode that plays a crucial role in studying the nuclear structure of extremely proton-rich nuclei near the proton drip line [1-5]. It was first proposed by Goldansky et al. [6, 7], who suggested that for the even- Z nuclei near or beyond the proton drip line, one-proton emission is energetically or dynamically forbidden due to the proton pairing effects, leading to the simultaneous emission of two protons instead [6, 7]. The early experiments aimed to investigate 2p radioactivity in light nuclei [1]. However, due to the limitations in experimental equipment and detection techniques, no clear evidence of 2p radioactivity was obtained [8, 9]. It was not until 2002 that 2p radioactivity was first observed in Fe at GSI [10] and GANIL [11], providing the first experimental confirmation of this decay mode. Later on, this phenomenon was successively observed in Zn [12], ^{48}Ni [13-16], and ^{67}Kr [17]. Overall, these observations not only confirmed the theoretical predictions but also established 2p radioactivity as a key probe for studying nuclear structure, including nuclear radius, spin and parity, deformation effect, tensor force effect, and so on [18-22].

The study of excited-state 2p radioactivity is a topic of considerable current interest in nuclear physics, which can provide rich nuclear structure information [2, 23-28]. Typically, there are two paths to populate the excited-state of the parent nucleus that can undergo 2p radioactivity. One is the β -delayed 2p radioactivity proposed by Janecke in 1965 [29], in which a parent nucleus first undergoes β decay to form an intermediate nucleus in an excited-state, which subsequently emits two protons [29]. The other is to populate the excited-state of the parent nucleus through nuclear reactions (such as p-capture, relativistic Coulomb excitation, and fragmentation reactions) including $^{14}\text{O}^*$ [30], $^{17}\text{Ne}^*$, $^{18}\text{Ne}^*$ [28, 31-34], S [35], and Ag [36].

In contrast to ground-state two-proton (2p) radioactivity, excited-state 2p emission is typically characterized by high decay energy and angular momentum [2, 23-27]. The high decay energy leads to an extremely short half-life, significantly shorter than that observed for ground-state 2p radioactivity [2, 23-27]. This poses significant challenges to the experimental identification of this decay mode and to precise theoretical predictions of the half-life for excited-state 2p radioactivity. Moreover, the emitted proton pair may carry away non-zero angular momentum from excited states, offering a unique probe for accessing the single-particle states occupied by the emitted protons [28]. Consequently, the investigation of excited-state 2p radioactivity is of considerable interest.

There are three main mechanisms proposed to describe 2p radioactivity. (i) Direct 2p emission. A strongly correlated proton pair is emitted simultaneously from the parent nucleus [6]. (ii) Sequential emission. The parent nucleus emits a proton, forming an intermediate state, which then emits another proton [37]. (iii) Three-body emission. Two protons are emitted simultaneously from the parent nucleus with a large opening angle [38]. For investigating the characteristics of 2p radioactivity, several theoretical models are applied within the picture of direct 2p emission, such as the density-dependent cluster model (DDCM) [39], the effective liquid drop model (ELDM) [24], the unified fission model (UFM) [25], the generalized liquid drop model (GLDM) [40-42], the Coulomb and proximity potential model (CPPM) [27, 43], the two-potential approach with Skyrme-Hartree-Fock (TPA-SHF) [44], the phenomenological model with a screened electrostatic barrier (SEB) [45], the Gamow-like model (GLM) [46], the Skyrme-Hartree-Fock-Bogoliubov theory (SHFB) [21, 22], and so on [47]. These theoretical models, constructed within the Gamow tunneling picture [48], differ primarily in the construction of the interaction potential between the daughter nucleus and the emitted 2p pair. Among these theoretical models, through the introduction of quasi-molecular shape mechanism and proximity energy, the GLDM can continuously describe the shape evolution from the parent nucleus to two separate fragments [49]. The proximity energy is used to describe the interaction in the neck region during the shape evolution and thus plays an essential role in constructing the potential between the daughter nucleus and the emitted 2p pair [49]. These features enable the GLDM to be successfully applied to describe various decay processes, including α decay [49-54], fission [55], fusion [56], proton radioactivity [57], and two-proton radioac-

tivity [41]. In our previous work, 15 different versions of the proximity energy were employed within the GLDM to construct the total interaction potential between the daughter nucleus and the emitted 2p pair [41]. A systematic study based on the latest experimental ground-state 2p radioactivity half-lives was performed to assess the predictive power of these proximity potentials. Among these formalisms, Prox. 77-Set 13 yields the smallest root-mean-square deviation from the experimental data, indicating that this parametrization provides the optimal description of ground-state 2p radioactivity half-lives [41].

Excited-state 2p radioactivity, which has been observed in nuclei including 14 O* [30], 17 Ne* [28, 34], 18 Ne* [31, 32], 29 S* [58], and 94 Ag* [36, 59], is typically accompanied by high decay energy and non-zero angular momentum transfer. Consequently, it serves as a probe for studying the dependence of the excited-state 2p radioactivity half-life on both angular momentum and decay energy. However, the high angular momentum significantly governs the centrifugal potential, thereby raising the total interaction potential. Meanwhile, the high decay energy may lead to difficulties in determining the classical turning points within the Wentzel-Kramers-Brillouin (WKB) approximation. Therefore, the theoretical description of excited-state 2p radioactivity half-lives remains challenging. It is thus of interest to examine whether the GLDM with improved proximity energy Prox. 77-Set 13 parametrization can be extended to effectively describe excited-state 2p radioactivity.

In this work, the GLDM with Prox. 77-Set 13 is adopted to study the 2p radioactivity half-lives for 14 O* [30], 17 Ne* [28, 34], 18 Ne* [31, 32], 29 S* [58], and 94 Ag* [36, 59]. The calculations obtained from this work are compared with the experimental data [28, 30–32, 34, 36, 59] and with the results calculated by the ELDM [24], the UFM [25], and the CPPM [27]. The 2p radioactivity half-lives calculated by this work are in good agreement with the available experimental data and show satisfactory consistency with calculations taken from three other theoretical models. Furthermore, this work systematically investigates the variation of excited-state 2p radioactivity half-lives with decay energy. The dependence of half-life on angular momentum is also examined, with 94 Ag* taken as a case. The organization of this paper is outlined as follows. Section II presents the theoretical framework for the GLDM and 2p radioactivity half-life. Section III provides detailed calculations and discussion. The paper concludes with a brief summary in Sec. IV.

II. THEORETICAL FRAMEWORK

A. Generalized liquid drop model

Under the framework of the generalized liquid drop model (GLDM), the total interaction potential includes the following components [49]:

$$E = E_V + E_S + E_C + E_{Prox} + E_l, \quad (1)$$

where E_V , E_S , E_C , E_{Prox} , and E_l represent the volume energy, the surface energy, the Coulomb energy, the proximity energy, and the centrifugal potential, respectively.

For one body shapes, E_V , E_S , E_C are calculated by

$$E_V = -15.494(1 - 1.8I^2)A, \quad (2)$$

$$E_S = 17.9439(1 - 2.6I^2)A^{2/3}(S/4\pi R_0^2), \quad (3)$$

$$E_C = 0.6e^2(Z^2/R_0) \times 0.5 \int (V(\theta)/V_0)(R(\theta)/R_0)^3 \sin \theta d\theta, \quad (4)$$

here $I = \frac{N-Z}{A}$ represents relative neutron excess of the parent nucleus. $V(\theta)$ is the electrostatic potential at the surface. R_0 is the radius of the parent nucleus, which can be expressed as follows

$$R_i = 1.28A_i^{1/3} - 0.76 + 0.8A_i^{-1/3} (i = 0, 1, 2). \quad (5)$$

For two body shapes, E_V , E_S , E_C are obtained by

$$E_V = -15.494[(1 - 1.8I_1^2)A_1 + (1 - 1.8I_2^2)A_2], \quad (6)$$

$$E_S = 17.9439[(1 - 2.6I_1^2)A_1^{2/3} + (1 - 2.6I_2^2)A_2^{2/3}], \quad (7)$$

$$E_C = 0.6e^2Z_1^2/R_1 + 0.6e^2Z_2^2/R_2 + e^2Z_1Z_2/r, \quad (8)$$

here A_i , Z_i , R_i , and I_i ($i = 1, 2$) represent the mass numbers, the proton numbers, the radii, and the relative neutron excesses of the emitted 2p pair and the daughter nucleus, respectively.

The proximity energy was initially proposed by Blocki to describe the interaction energy in the neck structure after contact in heavy-ion reactions [60]. Subsequently, it was introduced into the GLDM to describe the interaction energy within the neck or gap formed during the separation process between the daughter nucleus and emitted particle in decay processes [49]. In previous work, the capability of the GLDM with 15 different versions of the proximity energy to describe the ground-state 2p radioactivity half-lives was examined against the latest experimental data [41]. The calculations with Prox. 77-Set 13 were found to reproduce the experimental half-lives most satisfactorily [41]. The present work examines whether this parametrization can be extended to excited-state 2p radioactivity. The Prox. 77-Set 13 [61] is given as follows

$$E_{Prox}(r) = 4\pi\gamma b\bar{R}\phi(\xi), \quad (9)$$

where γ is the surface energy coefficient, defined by

$$\gamma = \gamma_0(1 - K_s A_s^2), \quad (10)$$

here $A_s \equiv \frac{N-Z}{A}$ represents the asymmetry parameter. In this work, $\gamma_0 = 0.911445$ (MeV/fm²) and $K_s = 2.2938$ [61]. The mean curvature radius \bar{R} can be calculated by

$$\bar{R} = \frac{C_1 C_2}{C_1 + C_2}, \quad (11)$$

here the matter radii C_1 and C_2 for the emitted 2p pair and the daughter nucleus are given by

$$C_i = R_i \left[1 - \left(\frac{b}{R_i} \right)^2 \right] (i = 1, 2). \quad (12)$$

The universal function $\phi(\xi)$ can be expressed as

$$\phi(\xi) = \begin{cases} -\frac{1}{2}(\xi - \xi_0)^2 - 0.0852(\xi - \xi_0)^3, & 0 < \xi \leq 1.2511, \\ -3.347 \exp\left(-\frac{\xi}{0.75}\right), & \xi \geq 1.2511, \end{cases} \quad (13)$$

here $\xi_0 = 2.54$ and $\xi = \frac{r - C_1 - C_2}{b}$ serve as the separation distance between the surfaces of the emitted 2p pair and the daughter nucleus. The width parameter b is taken as unity.

The centrifugal potential is expressed as

$$E_l(r) = \frac{\hbar^2 l(l+1)}{2\mu r^2}, \quad (14)$$

where l corresponds to the angular momentum taken away by the emitted 2p pair.

B. The 2p radioactivity half-life

Under the picture of direct diproton emission [6], 2p radioactivity is treated as a process in which a strongly correlated proton pair is emitted after tunneling through the Coulomb barrier of the parent nucleus [6]. This tunneling process is governed by the same quantum tunneling mechanism as α decay [62-72], proton radioactivity [57, 73], and cluster radioactivity [74-78].

The 2p radioactivity half-life can be calculated with the decay constant λ as

$$T_{1/2} = \frac{\ln 2}{\lambda}. \quad (15)$$

The decay constant λ can be given by

$$\lambda = S_{2p}\nu P, \quad (16)$$

here S is the spectroscopic factor for the ${}^2\text{He}$ cluster within the parent nucleus, which is derived from the cluster overlap approximation [42, 79]

$$S_{2p} = G^2[A/(A-2)]^{2n}\chi^2, \quad (17)$$

where $G^2 = (2n)!/[2^{2n}(n!)^2]$ [80], and n corresponds to the average principal proton oscillator quantum number, which can be obtained by $n \approx (3Z)^{1/3} - 1$ [81]. Here, χ^2 is the proton overlap function [42]. In the previous work, the value of χ^2 was determined to be 0.0275 by fitting the experimental data of ground-state 2p radioactivity [41].

The assault frequency ν can be obtained by

$$\nu = \frac{\sqrt{2E_{2p}/M_{2p}}}{2R_0}, \quad (18)$$

where E_{2p} and M_{2p} denote the kinetic energy and the mass of the emitted 2p pair, respectively.

Within the WKB approximation, the barrier penetrating probability P is calculated by

$$P = \exp\left[-\frac{2}{\hbar} \int_{r_{in}}^{r_{out}} \sqrt{2B(r)[E(r) - E_{sphere}]dr}\right], \quad (19)$$

where r is the distance between the center of mass of the emitted 2p pair and the daughter nucleus. The classical turning points (r_{in} and r_{out}) can be obtained by solving $E(r_{in}) = Q_{2p}$ and $E(r_{out}) = Q_{2p}$. $B(r) = \mu$ is defined as the reduced mass between the emitted 2p pair and the daughter nucleus.

III. RESULTS AND DISCUSSION

Unlike ground-state 2p radioactivity, excited-state 2p emission is typically characterized by high decay energy and non-zero angular momentum transfer [2, 23-28]. This poses significant challenges for theoretical models in describing the corresponding half-lives. The present work examines the capability of the GLDM with the improved proximity energy Prox. 77-Set 13 parametrization to describe excited-state 2p radioactivity, and systematically investigates the impact of these features on the half-life of 2p radioactivity.

In this work, the GLDM with the improved proximity energy is extended to calculate the 2p radioactivity half-lives for 14 O* [30], 17 Ne* [28, 34], 18 Ne* [31, 32], 29 S* [58], and 94 Ag* [36, 59]. presents the calculations obtained from this work together with those taken from the ELDM [24], the UFM [25], and the CPPM [27]. In Table 1, the first column represents the parent nucleus undergoing excited-state 2p radioactivity and the corresponding daughter nucleus. The second column lists the spin and parity values for the parent nuclei and the daughter nuclei. Based on spin-parity conservation, the calculated angular momentum taken away by the emitted 2p pair is listed in the third column except for S and 94 Ag*. For 29 S*, due to its undetermined spin and parity [58], the angular momentum taken away by the emitted 2p pair is set to $l = 0$ in this work. For Ag*, l is taken from the Ref. [36]. The fourth column contains the experimental decay energy of excited-state 2p radioactivity. The fifth column shows the logarithmic 2p radioactivity half-lives of experimental data. The sixth to ninth columns present the calculated logarithmic half-lives of excited-state 2p radioactivity obtained from this work, the ELDM [24], the UFM [25], and the CPPM [27], respectively.

As shown in Table 1, the present calculations exhibit a consistent trend with those obtained from the ELDM [24], the UFM [25], and the CPPM [27], indicating that the GLDM with the Prox. 77-Set 13 parametrization can be extended to calculate excited-state 2p radioactivity half-lives. For 14 O* , the excited-state at 7.77 MeV, corresponding to a decay energy of $Q_{2p} = 1.20$ MeV, was populated via the N + p reaction [30]. Despite the absence of a distinct signature for 2p emission, an upper limit on its decay width of $\Gamma < 6$ eV was determined, corresponding to a lower limit on the experimental 2p radioactivity half-life of $\log_{10} T_{1/2} > -16.12$ s for 14 O* [30]. From Table 1, it can be observed that the calculations of 2p radioactivity for O ($Q_{2p} = 1.20$ MeV) given by this work, the ELDM [24], and the UFM [25] are all longer than -16.12 s. Meanwhile, this work also calculates the half-lives for 2p emission from its higher excited-state at 9.72 MeV ($Q_{2p} = 3.15$ MeV) and 9.92 MeV ($Q_{2p} = 3.35$ MeV). The calculations taken from this work and other three theoretical models agree with each other. Moreover, for O , as the decay energy of 2p radioactivity increases from 1.20 to 3.15 MeV while the angular momentum remains fixed at $l = 2$, the 2p radioactivity half-lives calculated by the present work and three other theoretical models decrease by approximately three orders of magnitude. This reveals a strong sensitivity of the 2p radioactivity half-life to the decay energy. However, a contrasting trend emerges as the decay energy increases from 3.15 MeV ($l = 2$) to 3.45 MeV ($l = 4$), with the half-life increasing by nearly two orders of magnitude. The increase in angular momentum from 2 to 4 enhances the centrifugal barrier, thereby reducing the tunneling probability and resulting in a longer half-life. Conversely, the increase in decay energy increases the tunneling probability, which shortens the half-life. This contrasting trend reflects the competition between the effects of decay energy and angular momentum on excited-state 2p radioactivity half-lives.

For 17 Ne* , following population of the first excited-state ($3/2^-$) via Ne+197

Au Coulomb excitation, a lower limit of experimental 2p radioactivity half-life $\log_{10} T_{1/2} > -10.59$ s for 2p emission was obtained [28]. Notably, Table 1 shows that the logarithmic half-lives for $\text{Ne}^* (3/2^-)$ calculated by this work and other three theoretical models exceed this limit. Furthermore, for Ne^* , half-lives for the higher excited-state ($5/2^-$ and $1/2^+$) are calculated, and the present results agree well with calculations from the three other theoretical models. Notably, for Ne , a mere 0.47 MeV increase in Q_{2p} from first the second ($5/2^-$) excited-state at fixed $l = 2$ leads to an decrease in the four types of calculated 2p radioactivity half-lives by nearly five orders of magnitude, demonstrating the high sensitivity of the 2p radioactivity half-life to decay energy.

Regarding 18Ne^* , combining the measurements from Gomez del Campo et al. [31] and Raciti et al. [32], the experimental logarithmic half-life for excited-state 2p radioactivity is -16.15 s [31, 32]. As shown in Table 1, the calculated logarithmic 2p radioactivity half-life given by the present work for $\text{Ne}^* (Q_{2p} = 1.63 \text{ MeV}$ [31]) agrees well with the results yielded by the three other theoretical models, further confirming the reliability of the GLDM with the Prox. 77-Set 13 parametrization. For S [58], owing to the undetermined spin and parity of the parent nucleus, this work assumes zero angular momentum transfer for the emitted proton pair and calculates the corresponding half-life. The present calculations are in excellent agreement with those obtained from the three other theoretical models.

In 2006, Mukha et al. reported evidence for 2p radioactivity via simultaneous emission from the 21^+ state of 94Ag^* , for which the experimental logarithmic 2p radioactivity half-life and decay energy are 1.90 s and 1.90 MeV, respectively [36]. Furthermore, based on the analysis of the daughter nucleus spin and considering the possible angular momentum carried by unassigned γ rays, the angular momentum transfer was conservatively constrained to the range of $6 \leq l \leq 10$ [36]. Subsequently, Kankainen et al. measured the masses of Rh and 94Pd , from which the Q_{2p} for the 94Ag high-spin isomer are deduced to be 1.98, 2.05, and 3.45 MeV, respectively [59]. As experimental half-lives for 94Ag^* at $Q_{2p} = 1.98, 2.05,$ and 3.45 MeV remain undetermined, the measured experimental data of $\log_{10} T_{1/2} = 1.90$ s at $Q_{2p} = 1.90 \text{ MeV}$ is adopted as the reference for comparison with the present calculations and three other theoretical models. As shown in Table 1, the present calculations agree well with those taken from the three other theoretical models. Furthermore, for Ag , the calculations given by this work and the three other models reproduce the experimental data well when Q_{2p} is assumed to be 3.45 MeV and angular momentum is in the range of $6 \leq l \leq 10$. This suggests that the Q_{2p} for Ag^* may be 3.45 MeV, with angular momentum in the range of $6 \leq l \leq 10$.

For a clearer comparison, the experimental data of 2p radioactivity half-lives [28, 30-32, 36] and the calculations obtained from this work, the ELDM [24], the UFM [25], and the CPPM [27] are presented in [Figure 1: see original paper]. In this figure, the subfigure (a) represents the comparative analysis of logarithmic half-lives for O, Ne, Ne^* , and S obtained by this work (red open spheres), the

ELDM (magenta solid triangles) [24], the UFM (blue open triangles) [25], the CPPM (orange open stars) [27], and experimental data (black solid spheres) [28, 30–32, 36], respectively. The subfigure (b) is the same as (a), but it depicts the logarithmic half-lives for Ag^* . The black open squares represent the possible experimental data suggested by this work for Ag^* . From Fig. 1, it can be seen that the results calculated by this work are nearly coincident with those taken from the three other theoretical models. For $^{14}\text{O}^*$ [30], ^{20}Ne [28, 34], and ^{20}Ne [31, 32], the calculations yielded by the four theoretical models can well reproduce the experimental data. For Ag^* [36, 59], the calculations given by the four theoretical models can reproduce the experimental data when Q_{2p} is 3.45 MeV and l is 10. This indicates that the GLDM with the improved proximity energy can be successfully extended to calculate the excited-state 2p radioactivity half-lives.

Compared with ground-state 2p radioactivity, excited-state 2p emission typically exhibits high decay energy [2], rendering the 2p radioactivity half-life highly sensitive to variations in Q_{2p} , as shown in Table 1. To further investigate this dependence, the variation of the logarithmic 2p radioactivity half-life with decay energy for Ag^* , calculated by using the GLDM with Prox. 77-Set 13 parametrization, is plotted in [Figure 2: see original paper]. In this figure, the blue stars, the cyan diamonds, the orange pentagons, the magenta squares, and the red circles denote the calculated logarithmic half-lives of Ag^* as a function of Q_{2p} for $l = 6 \sim 10$, respectively. As evident from this figure, the logarithmic half-life for Ag^* [36, 59] exhibits a linear increase with $Q_{2p}^{-1/2}$. This trend indicates a sharp decrease in the half-life as the decay energy increases. Notably, as the decay energy rises from 2.05 to 3.45 MeV, for fixed angular momentum ($l = 6$), the logarithmic half-life calculated in this work decreases from 7.28 to -1.77 s. This can be attributed to the enhanced tunneling probability at higher decay energy, which shortens the half-life. This clearly demonstrates the high sensitivity of excited-state 2p radioactivity half-lives to decay energy. Furthermore, a linear relationship is found between the logarithmic half-life and $Q_{2p}^{-1/2}$. This supports the interpretation that 2p radioactivity proceeds via quantum tunneling.

Excited-state 2p emission typically involves non-zero angular momentum transfer [24, 28]. Therefore, investigating excited-state 2p radioactivity provides insights into the sensitivity of the half-life to angular momentum [24, 28], and enables the identification of the single-particle states occupied by the emitted protons [28]. For $^{94}\text{Ag}^*$ ($Q_{2p} = 1.90$ MeV) [36], [Figure 3: see original paper] is plotted to clearly illustrate the influence of angular momentum on the total interaction potential between the daughter nucleus and the emitted 2p pair. The figure shows that as the angular momentum increases, the barrier peak of the total interaction potential shifts to the left and the potential curves are significantly raised. This phenomenon indicates that as the angular momentum increases, the centrifugal potential rises significantly, thereby raising the total interaction potential between the daughter nucleus and the emitted

2p pair. This suppresses the tunneling probability, prolonging the half-life of excited-state 2p radioactivity.

To further investigate the relationship between 2p radioactivity half-life and angular momentum at fixed decay energy, the logarithmic half-lives for Ag^* , calculated by using the GLDM with Prox. 77-Set 13 parametrization, are plotted as a function of angular momentum for $Q_{2p} = 1.90$ MeV (blue open squares), $Q_{2p} = 1.98$ MeV (purple open diamonds), $Q_{2p} = 2.05$ MeV (cyan open stars), and $Q_{2p} = 3.45$ MeV (red solid circles) in [Figure 4: see original paper]. This figure shows that for fixed decay energy, the logarithmic half-life increases continuously with angular momentum. For $^{94}\text{Ag}^*$ [36], Mukha et al. conservatively estimated the angular momentum transfer to be in the range of 6–10 [36]. Within this range, the GLDM with the Prox. 77-Set 13 parametrization is used to calculate the 2p radioactivity half-life of Ag^* . As shown in Fig. 4, the calculations reproduce the experimental data for $Q_{2p} = 3.45$ MeV and $l = 8 \sim 10$. Furthermore, based on spin-parity conservation and the tentative assignments of 21^+ for the parent nucleus and 11^+ for the daughter nucleus, the angular momentum transfer is deduced to be $l = 10$. By combining the present calculations with experimental data and spin-parity conservation, this work suggests that the 2p emission from Ag carries away angular momentum $l = 10$. Due to these unique decay characteristics, excited-state 2p radioactivity serves as an important tool for extracting nuclear structure information [2, 23–27]. Although its relatively short 2p radioactivity half-life poses experimental challenges, the study of this decay mode remains essential and valuable.

IV. SUMMARY

In summary, excited-state 2p radioactivity is generally characterized by high decay energy and non-zero angular momentum transfer, in contrast to ground-state 2p emission, providing an important probe for exploring the nuclear structure of extremely proton-rich nuclei. Nevertheless, the theoretical description of the corresponding 2p radioactivity half-lives remains challenging. This work examines whether the generalized liquid drop model with the improved proximity energy Prox. 77-Set 13 parametrization can be extended to excited-state 2p radioactivity. Calculations are performed for $^{14}\text{O}^*$, $^{17}\text{Ne}^*$, $^{18}\text{Ne}^*$, $^{29}\text{S}^*$, and $^{94}\text{Ag}^*$, and compared with experimental data and calculations given by the effective liquid drop model, the unified fission model, and the Coulomb and proximity potential model. The present results reproduce the experimental data well and agree with other theoretical results. Taking Ag^* as an example, this work systematically investigates the sensitivity of 2p radioactivity half-lives to variations in 2p decay energy and angular momentum. The 2p radioactivity half-life is found to be highly sensitive to 2p decay energy and exhibits a positive correlation with angular momentum. Furthermore, by comparing with experimental data and incorporating spin-parity conservation, this work suggests that the most probable 2p decay energy and angular momentum for $^{94}\text{Ag}^*$ are $Q_{2p} = 3.45$ MeV and $l = 10$, respectively. This study provides a ref-

erence for future experimental investigations of excited-state $2p$ radioactivity, shedding new light on the properties of exotic proton-rich nuclei.

Note: Figure translations are in progress. See original paper for figures.

Source: ChinaXiv – Machine translation. Verify with original.