



---

Soviet-era science, translated into English

# | | Pages |

|—|—:|

1967

SovietRxiv

---

View the original and related papers at <https://sovietrxiv.org/items/ru-196701.17083>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

## Abstract

## Full Text

	Pages
I. I. Bavrin. Integral representations of functions holomorphic in convex multiple-circular regions, and Taylor' s formula	1217
V. A. Baranova. Interpolation of functions that are regular on closed sets	1221
B. A. Vostretsov, A. V. Ignatieva. Existence of the best approximations of functions by the sums of a finite number of plane waves of given directions in the $L_p$ metric	1225
V. N. Goldberg. The appearance of discontinuities on the continuation of solutions to non-linear mixed problems for the string equation	1229
P. P. Zabreiko, M. A. Krasnosel' skii. A method for the production of new stationary point principles	1233
I. A. Kiprianov. Self-adjoint extensions of certain singular partial differential operators	1236
Iu. S. Kolesov. Periodical solutions to a class of differential equations with hysteretic non-linearity	1240
A. Lelek. On an estimate of the defect of a space	1244
A. M. Olevskii. Growth order of Lebesgue functions of bounded orthonormal systems	1247
E. M. Semenov. Interpolation of linear operators and estimates of Fourier coefficients	1251
I. B. Simonenko. Convolution operators in cones	1255

	Pages
A. B. Sosinskii. Homotopy of complements to knots	1258
M. M. Khapaev. A Liapunov type theorem	1262
E. R. Tsekanovskii. Description of generalized extensions with a unidimensional imaginary component of the differentiation operator without a spectrum	1266
<b>FLUID MECHANICS</b>	
I. N. Gorshkova, K. P. Staniukovich. Initial stage of two-dimensional unsteady gas flow	1270
<b>PHYSICS</b>	
Iu. N. Denisiuk, D. I. Stasel' ko. The possibility of holograms being obtained by means of a referent ray whose wavelength differs from that of the radiation scattered by the object	1274
E. D. Martynov, V. I. Trefilov, S. A. Firstov, B. I. Beresnev, Iu. N. Riabinin. Electron-microscope investigation of chromium and molybdenum deformed under a high pressure	1276
J. J. Shipuk, S. V. Pashkin. Ionization instability of plasma in crossed fields	1278
<b>TECHNICAL PHYSICS</b>	
L. V. Kirenskii, S. G. Rusova, G. I. Rusov, N. I. Sivkov. Some magnetic properties of thin single-crystal films of Fe–Ni–Co alloys	1282
N. M. Mikhin, I. V. Kragel' skii. Variation of the tangency area between solids as they are moving closer together	1285
<b>GEOPHYSICS</b>	

## Corrections

In my article (I. I. Ogievetskii, “Summation of unbounded sequences by linear regular methods” ), published in *Doklady Akademii Nauk*, vol. 167, no. 5, 1966, the first three lines of the statement of Theorem 4 (p. 990) should read as follows:

**Theorem 4.** Whatever  $p$  unbounded linearly independent sequences  $\{s^1\}, \{s^2\}, \dots, \{s^p\}$  and numbers  $x_1, x_2, \dots, x_p$  may be, one can effectively construct a normal regular matrix  $A$  such that: 1) the matrix ...

*I. I. Ogievetskii*

*Note: Figure translations are in progress. See original paper for figures.*

*Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.*